

Do morphological identification and DNA metabarcoding provide different perspectives of insect communities?


Nicole Remmel, Dominik Buchner, Julian Enss, Viktor Hartung, Florian Leese, Ellen Welti, James Sinclair, & Peter Haase



SENCKENBERG
world of biodiversity

Leibniz
Leibniz
Association

Reported declines in insects





Metabarcoding



Trapped



Preserved




Homogenized



DNA extraction
& PCR



Sequencing




Matched

< 50€ per sample
about 3 weeks / 1000 samples

Limitations

1. Presence/absence only
2. Potential biomass bias

Abundance




Body size




Limitations

1. Presence/absence only
2. Potential biomass bias
 - Low-abundance species
 - Smaller-bodied species

Do morphological identification and DNA metabarcoding provide different perspectives of insect communities?




Morphological
(pick out and put back in)




Metabarcoding
(same sample)


Methods



Methods



Methods





Metrics

1. Total richness

→ Apiformes (bees), Heteroptera (true bugs), Syrphidae (hoverflies), Rhopalocera (butterflies), Trichoptera, Ephemeroptera, Plecoptera, Mecoptera

2. Richness of four taxonomic groups

Metrics



Metrics

1. Total richness

→ Apiformes (bees), Heteroptera (true bugs), Syrphidae (hoverflies), Rhopalocera (butterflies), Trichoptera, Ephemeroptera, Plecoptera, Mecoptera

2. Richness of four taxonomic groups

→ Bees, butterflies, hoverflies, and true bugs

3. Species detection (0/1)

Metrics

Morphology



Pieris rapae



Metabarcoding



Pieris rapae

Metrics

1. Total richness


→ Apiformes (bees), Heteroptera (true bugs), Syrphidae (hoverflies), Rhopalocera (butterflies), Trichoptera, Ephemeroptera, Plecoptera, Mecoptera

2. Richness of four taxonomic groups

→ Bees, butterflies, hoverflies, and true bugs

3. Species detection (0/1) ~ Abundance + Body size

Metrics



Metrics

1. Total richness

→ Apiformes (bees), Heteroptera (true bugs), Syrphidae (hoverflies), Rhopalocera (butterflies), Trichoptera, Ephemeroptera, Plecoptera, Mecoptera

2. Richness of four taxonomic groups

→ Bees, butterflies, hoverflies, and true bugs


3. Species detection (0/1) ~ Abundance + Body size



4. Community composition (taxa and body size)

Total richness



Total richness






Morphological richness



Summary

1. Richness



- Missing species at higher diversity, driven by true bugs and bees
- Can detect more hoverflies and butterflies






Detection success (0/1) ~ Abundance + Body size





Detection success (0/1) ~ Abundance + Body size




Detection success (0/1) ~ Abundance + Body size



Detection success (0/1) ~ Abundance + Body size







Summary

1. Richness


- Missing species at higher diversity, driven by true bugs and bees
- Can detect more hoverflies and butterflies

2. Species detection


- Missing rarer hoverflies, bees, and true bugs, and smaller-bodied true bugs




Taxonomic composition




Taxonomic composition




Taxonomic composition – Presence/absence




Taxonomic composition – Abundance weighted



Community body size – Presence/absence




Community body size – Abundance weighted



Summary


1. Richness

- Missing species at higher diversity, driven by true bugs and bees
- Can detect more hoverflies and butterflies




2. Species detection

- Missing rarer hoverflies, bees, and true bugs, and smaller-bodied true bugs



3. Composition

- No strong differences in taxonomic composition, but traits may be affected by missing smaller-bodied species





Take-home messages

- Metabarcoding can provide an accurate overview of insect community differences in a way that is comparable to morphological identification
 - for potentially less time and cost
 - can better resolve some taxonomic groups to species level
- However, metabarcoding can miss some low-abundance and smaller-bodied species
 - These species can provide key functions
 - Can somewhat bias estimates of richness and trait composition

Questions?

DNA metabarcoding and morphological identification reveal similar richness, taxonomic composition and body size patterns among flying insect communities

Nicole Remmel^{1,2} | Dominik Buchner² | Julian Enss^{2,3} | Viktor Hartung⁴ |
Florian Leese^{2,3}  | Ellen A. R. Welti⁵ | James S. Sinclair¹ | Peter Haase^{1,2,3} 

DOI: [10.1111/icad.12710](https://doi.org/10.1111/icad.12710)