

Dissertation 10/2008

**Entwicklung eines Testsystems zur Erfassung der
in situ Biodegradation im Grundwasser**

Nicole Stelzer

Helmholtz-Zentrum für Umweltforschung (UFZ)



00318730

ISSN 1860-0387

Entwicklung eines Testsystems zur Erfassung der *in situ* Biodegradation im Grundwasser

D i s s e r t a t i o n

zur Erlangung des Doktorgrades
der Mathematisch-Naturwissenschaftlichen Fakultät
der Christian-Albrechts-Universität
zu Kiel

vorgelegt von

Diplom-Geologin Nicole Stelzer

geboren am 17.06.1979 in Borna

Archiv

Kiel, Mai 2008

Referent: Prof. Dr. Andreas Dahmke
Koreferent: Dr. habil. Hans-Hermann Richnow

Tag der mündlichen Prüfung: 10.07.2008
zum Druck genehmigt: 10.07.2008

gezeichnet: Prof. Dr. Jürgen Grotzmeyer, Dekan

Inhalt

Tabellenverzeichnis	VI
Abbildungsverzeichnis	VI
Abkürzungsverzeichnis	X
Kurzfassung	XIII
Abstract	XVI
1 Einleitung	1
1.1 Altlastenproblematik	1
1.2 Relevante Schadstoffe	1
1.3 Natürliche Selbstreinigung: <i>Natural Attenuation</i>	3
1.4 Nachweis der <i>in situ</i> Biodegradation	5
1.4.1 Stabile Isotopenfraktionierung	7
1.4.2 Tracerversuche	10
1.4.3 Molekularbiologische Methoden	12
1.5 <i>In situ</i> Mikrokosmen	18
1.6 Zielstellung und Inhalt der Arbeit	20
2 Analyse des anaeroben Benzolabbaus: Vergleich von <i>In situ</i> Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen	23
2.1 Kurzfassung	23
2.2 Abstract	23
2.3 Einleitung	24
2.4 Material und Methoden	26
2.4.1 Isotopen markierte Substanzen und Chemikalien	26
2.4.2 Standortbeschreibung	26
2.4.3 Tiefenorientierte Grundwasserbeprobung	27
2.4.4 Analytik der GW-Proben	28
2.4.5 Herstellung der BACTRAPS	28
2.4.6 Beladung der BACTRAPS	29
2.4.7 Einbau in die Grundwassermessstelle	30
2.4.8 Probenaufbereitung	30
2.4.9 Identifikation der Fettsäuremethylester (FAME) mittels GC-MS	30
2.4.10 Analyse der Kohlenstoffisotopenverhältnisse der FAME mittels GC-C-IRMS	31
2.4.11 Berechnungen	31
2.5 Ergebnisse	33
2.5.1 Geochemie	33
2.5.2 Ergebnisse der tiefenorientierten Untersuchungen mit den BACTRAPS	39

Inhalt

2.6	Diskussion	41
2.6.1	Geochemie	41
2.6.2	BACTRAP	42
2.7	Zusammenfassung	43
2.8	Danksagung	44
3	<i>In situ</i> microcosms to evaluate natural attenuation potentials in contaminated aquifers	45
3.1	Abstract	45
3.2	Introduction	45
3.3	Materials and Methods	47
3.3.1	Chemicals	47
3.3.2	Field Sites	47
3.3.3	Multi level packer systems used for depth discrete experiments	49
3.3.4	Preparation and incubation of <i>in situ</i> microcosms (BACTRAP [®])	49
3.3.5	Extraction and Derivatisation of Fatty Acids	51
3.3.6	Analysis	51
3.3.7	Molecular biological analysis	52
3.4	Results and Discussion	52
3.4.1	Zeitz	52
3.4.2	Gneisenau: Upper quaternary aquifer	56
3.4.3	Gneisenau: Lower cretaceous aquifer	57
3.4.4	Molecular biological analysis of microbial communities on BACTRAPS	59
3.4.5	Investigation of metabolic pathways	64
3.5	Conclusions	65
3.6	Acknowledgements	66
4	Assessment of Microbial <i>In Situ</i> Activity in Contaminated Aquifers	67
4.1	Summary	67
4.2	Introduction	68
4.3	Materials and methods	70
4.3.1	Chemicals	70
4.3.2	Field Sites (Zeitz, Bitterfeld)	70
4.3.3	Isotope fractionation and calculations	71
4.3.4	Preparation and incubation of <i>in situ</i> microcosms	72
4.3.5	Groundwater sampling and analyses	73
4.4	Results and Discussion	76
4.4.1	Benzene and Toluene degradation at the SAFIRA reference site Zeitz	76
4.4.2	Monochlorobenzene degradation at the SAFIRA reference site Bitterfeld	89
4.5	Concluding remarks	93

4.6 Acknowledgements	94
5 Sensitive detection of anaerobic monochlorobenzene degradation using stable isotope tracers	96
5.1 Abstract	96
5.2 Introduction	96
5.3 Materials and Methods	97
5.3.1 Chemicals	97
5.3.2 Site description of the aquifer in Bitterfeld (Saxony-Anhalt, Germany)	97
5.3.3 Preparation and incubation of <i>in situ</i> microcosms	99
5.3.4 Extraction and derivatization of fatty acids (FA)	100
5.3.5 Laboratory microcosms	100
5.4 Results and Discussion	101
5.4.1 <i>In situ</i> microcosms	101
5.4.2 Laboratory microcosms and enrichments	104
5.4.3 Acknowledgements	107
6 Integrative approach to delineate Natural Attenuation of chlorinated benzenes in anoxic aquifers	109
6.1 Abstract	109
6.2 Synopsis	109
6.3 Introduction	109
6.4 Material and Methods	111
6.4.1 Study Site and sampling	111
6.4.2 Chemicals	111
6.4.3 Field study with <i>in situ</i> microcosms	112
6.4.4 Laboratory study with enrichment cultures	112
6.4.5 Determination of $\delta^{13}\text{C}$ values for the total chlorinated benzenes	112
6.4.6 Principal Component Analysis	113
6.5 Results and Discussion	113
6.5.1 <i>In situ</i> microcosms	113
6.5.2 Laboratory enrichment cultures	114
6.5.3 Isotope balance computation	115
6.5.4 Statistical analysis of the field data	119
6.5.5 Implications for environmental studies and site management in terms of natural attenuation	122
6.5.6 Acknowledgment	123
6.5.7 Supporting Information Available	123
7 Assessment of <i>in situ</i> biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland	124
7.1 Abstract	124

Inhalt

7.2	Capsule	124
7.3	Introduction	124
7.4	Materials and methods	125
7.4.1	Design and characteristics of the wetland	125
7.4.2	Sampling	126
7.4.3	Physico-chemical and geochemical parameters of the pore water samples	126
7.4.4	Analysis of benzene, MCB and metabolites	127
7.4.5	<i>In situ</i> microcosms (BACTRAPS)	127
7.4.6	Statistical analysis	129
7.5	Results	129
7.5.1	Characterisation of pore water chemistry	129
7.5.2	Carbon isotopic composition of MCB	132
7.5.3	<i>In situ</i> microcosm experiment	134
7.6	Discussion	138
7.7	Conclusion	141
7.8	Acknowledgements	141
8	Zusammenfassung	142
8.1	Beurteilung der <i>in situ</i> Mikrokosmen zum Nachweis der Biodegradation	142
8.2	Vergleich mit anderen Methoden	147
8.3	Implementierung in die Altlastenpraxis	150
8.4	Ausblick	153
9	Literatur	154
10	Acknowledgements	173
11	Eidesstattliche Erklärung	174
12	Persönlicher Werdegang	175
13	Publikationsliste	176
13.1	Artikel	176
13.2	Konferenzvorträge	176
13.3	Poster	177
14	Anhang (siehe CD)	179

- A N. Stelzer, A. Fischer, M. Kästner, H.-H. Richnow (2006): Analyse des anaeroben Benzolabbaus: Vergleich von *In situ* Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen. Grundwasser 11(4), S. 247-258.
- B A. Fischer, N. Stelzer, H. Eisenmann, H.-H. Richnow (2006): Nachweis des mikrobiellen Schadstoffabbaus in Grundwasserleitern. TerraTec 15(1-2), S. 14-17.

- C N. Stelzer, C. Büning, F. Pfeifer, C.C. Tebbe, I. Nijenhuis, M. Kästner, H.-H. Richnow (2006): *In situ* microcosms to evaluate natural attenuation potentials in contaminated aquifers. *Organic Geochemistry* 37(10), S. 1394-1410.
- D M. Kästner, A. Fischer, I. Nijenhuis, R. Geyer, N. Stelzer, P. Bombach, C.C. Tebbe, H.-H. Richnow (2006): Assessment of Microbial *In Situ* Activity in Contaminated Aquifers. *Engineering in Life Science* 6(3), S. 234-251.
- E I. Nijenhuis, N. Stelzer, M. Kästner, H.-H. Richnow (2007): Sensitive Detection of Anaerobic Monochlorobenzene Degradation Using Stable Isotope Tracers. *Environmental Science and Technology* 41(11), S. 3836-3842.
Supporting Information
- F N. Stelzer, G. Imfeld, M. Thullner, J. Lehmann, A. Poser, H.-H. Richnow, I. Nijenhuis (2008): Integrative approach to delineate Natural Attenuation of chlorinated benzenes in anoxic aquifers. *Environmental Science and Technology*, submitted.
Supporting Information
- G M. Braeckeveldt, H. Rokadia, G. Imfeld, N. Stelzer, H. Paschke, P. Kuschk, M. Kästner, H.-H. Richnow, S. Weber (2007): Assessment of *in situ* biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland. *Environmental Pollution* 148(2), S. 428-437.
- H M. Braeckeveldt, H. Rokadia, G. Mirschel, S. Weber, G. Imfeld, N. Stelzer, P. Kuschk, M. Kästner, H.-H. Richnow (2007): Biodegradation of chlorobenzene in a constructed wetland treating contaminated groundwater. *Water Science and Technology* 56(3), S. 57-62.
- I A. Fischer, K. Theuerkorn, N. Stelzer, M. Gehre, M. Thullner, H.-H. Richnow (2007): Applicability of Stable Isotope Fractionation Analysis for the Characterization of Benzene Biodegradation in a BTEX-contaminated Aquifer. *Environmental Science and Technology* 41(10), S. 3689-3696.
Supporting Information

Tabellenverzeichnis

Tabellenverzeichnis

Tab. 1-1: Relative Häufigkeiten [%] der wichtigsten stabilen Isotope ausgewählter Elemente und Angabe der internationalen Standards (Hoefs, 1997)	8
Tab. 1-2: Übersicht ausgewählter Biomarker bestimmter Organismen, verändert nach (Boschker and Middelburg, 2002; Green and Scow, 2000; Kaur et al., 2005; Zelles, 1999). Für die Nomenklatur der Fettsäuren siehe Kapitel 2.4.9.	15
Tab. 2-1: Berechnung des biologischen Abbaus [B%] für sulfatreduzierende ($\alpha_S = 1.0036$) und methanogene Bedingungen ($\alpha_M = 1.0019$) unter Berücksichtigung der Schwankungsbreite der Isotopensignatur von Benzol in der Quelle ($R_0 = -28,5 \text{ ‰}$ bis $-29,5 \text{ ‰}$).....	35
Tab. 3-1: Carbon isotope composition of fatty acids extracted from <i>in situ</i> microcosms incubated with $^{13}\text{C}_6$ labelled benzene at monitoring well 23/00 (Zeitz, Germany) for 6 weeks. A to H refers to incubation depth indicated in Fig. 3-2.	55
Tab. 3-2: Carbon isotope composition of fatty acids extracted from <i>in situ</i> microcosms incubated with $^{13}\text{C}_6$ labelled benzene and $^{13}\text{C}_7$ labelled toluene in the upper Quaternary monitoring well (P10F) and in the lower Cretaceous monitoring wells (P8T, P11T) at the field site Gneisenau (Germany) for 12 weeks.	58
Tab. 3-3: Characterization of rRNA genes isolated from BACTRAPS incubated in the lower (P11T) and upper (P10F) aquifer at the field site Gneisenau (Germany).	61
Tab. 4-1: Concentrations and ^{13}C abundance of contaminants in the <i>in situ</i> microcosms (BACTRAP®'s) before and after exposure in well Saf Zz 32/02 (modified from Geyer et al., 2005).....	82
Tab. 4-2: Characterization of rRNA genes isolated from groundwater and microcosms loaded with toluene or benzene and controls after exposure in well Saf Zz 32/02 (see also Fig. 4-6)	88
Tab. 5-1: Carbon isotope composition of benzene and MCB extracted from blank, unloaded, and [$^{13}\text{C}_6$] MCB loaded <i>in situ</i> microcosms incubated for 7 weeks at different depths at the multilevel monitoring well SAF4 in Bitterfeld. n.a.= not analysed. The standard deviation for the analysis of the carbon stable isotope composition was $<1\text{ ‰}$ with $\delta^{13}\text{C} < 500 \text{ ‰}$ and between 5-10 ‰ with $\delta^{13}\text{C} > 500 \text{ ‰}$	102
Tab. 6-1: Carbon isotope signature of TLFA methyl esters $\delta^{13}\text{C}_{\text{FAME}}$ [%] extracted from <i>in situ</i> microcosms amended with [$^{13}\text{C}_6$] MCB and incubated in the contaminated aquifer at the study site.	114
Tab. 7-1: Geochemical characterisation, MCB and benzene concentrations of samples collected from (a) unplanted plot and (b) planted plot. Values represent the depth profiles average in the soil compartments (0.5 - 4 m) and average (0 m and pond) over the study period (13.04.05-29.09.05). Standard deviation is indicated in parentheses.....	130
Tab. 7-2: Carbon isotope composition of fatty acids extracted from <i>in situ</i> microcosms incubated for 6 weeks with $^{13}\text{C}_6$ labelled monochlorobenzene at 1.5, 2.5 and 4.5 m from the inflow point as well as at the pond.....	137
Tab. 8-1: Anwendungsbeispiele von <i>in situ</i> Mikrokosmen (BACTRAP®)	152

Abbildungsverzeichnis

Fig. 1-1: Ländervergleich Deutschland und USA über das Vorkommen und die Häufigkeit organischer Schadstoffe im Abstrom von Schadensfällen aus (Teutsch et al., 1997).	2
Fig. 1-2: Übersicht kultivierungsabhängiger und -unabhängiger Methoden zur Charakterisierung mikrobieller Gemeinschaften (Weiss and Cozzarelli, 2008).	13
Fig. 1-3: Konzept der Stabilen Isotopenbeprobung von Nukleinsäuren am Beispiel des RNA-SIP (Friedrich, 2006a)	14

Fig. 1-4: Konzept der <i>in situ</i> Mikrokosmen (BACTRAP [®]) basierend auf ¹³ C-markierten Substraten zum Nachweis mikrobieller Aktivität anhand der Inkorporation der isotopischen Markierung in die Biomarkermoleküle.....	19
Fig. 2-1: Verteilung der BTEX-Konzentrationen ($\mu\text{g L}^{-1}$) im oberen Aquifer im Bereich des ehemaligen Hydrierwerkes bei Zeitz (Sachsen-Anhalt).	27
Fig. 2-2: Links, Schichtenprofil und Ausbauplan der Grundwassermessstelle Saf ZZ 23/00. Tiefe 1 bis 8 bezeichnet die Position der Miniatur-Doppelventilpumpen und BACTRAPs. Rechts, Foto des MLPS mit Edelstahlkäfigen für BACTRAPs vor dem Einbau in die Messstelle.	28
Fig. 2-3: Schematischer Aufbau eines BACTRAP (links) und Foto der Befestigung des Edelstahlkäfiges mit BACTRAP an der Packermembran (rechts).	29
Fig. 2-4: Links vertikale Verteilung der Eisen(II)-(○), Eisen(III)-(●), Nitrat-(*)-, Mangan-(▲), Sulfat-(■), und rechts der Benzolkonzentrationen (●) sowie der Kohlenstoffisotopenverhältnisse des Benzols (■).	35
Fig. 2-5: Vertikale Verteilung der Kohlenstoffisotopenverhältnisse von Methan [■], DIC [●] und den Anreicherungsfaktoren [▼] (links) sowie der Methan- [■] und DIC-Konzentrationen [●] (rechts) in der Messstelle Saf ZZ 23/00.....	37
Fig. 2-6: Übersicht der Isotopensignaturen der Fettsäuren $\delta^{13}\text{C}_{\text{FA}}$ [%] in den einzelnen Beprobungshorizonten. Die Abkürzungen und Nomenklatur der Fettsäuren sind im Kapitel Material und Methoden (2.4.9) erklärt.	40
Fig. 3-1: Concentration of BTEX [$\mu\text{g L}^{-1}$] in the upper aquifer in the area of the former hydrogenation plant in Zeitz (Saxony-Anhalt, Germany). The locations of the <i>in situ</i> microcosm experiments are indicated. At well 23/00 the multilevel experiment was carried out and well 32/00 was used for the <i>in situ</i> microcosm study by Geyer et al. (2005). The arrow depicts the general groundwater flow direction.	48
Fig. 3-2: Left: Geological profile at monitoring well 23/00 (Zeitz, Germany). The location of the filter screen is illustrated. The groundwater table was located at 9.05 m below ground. <i>In situ</i> microcosms were installed at A, C, D, F, G, H and samples for geochemical analyses were taken from all depths except A. Right: The picture shows the multi level sampling system with stainless steel containers for <i>in situ</i> microcosms before subjection into the well.....	50
Fig. 3-3: Left: Concentration of sulphate (○), nitrate (*), sulphide (▲), ferrous iron (■), ferric iron (□) and manganese (x) in [$\mu\text{mol L}^{-1}$]. Right: benzene (■) and methane (●) concentration in [$\mu\text{mol L}^{-1}$] at Zeitz (Germany) monitoring well 23/00.	53
Fig. 3-4: The composition of the total fatty acid fraction (m/z 74) obtained from <i>in situ</i> microcosm experiment deployed at various depths (A to H) in well 23/00 at the field site Zeitz (Germany). The depths of the microcosm (A to H) and the position in the aquifer are shown in Fig. 3-2.	54
Fig. 3-5: Comparison of the composition of the total fatty acid fraction (m/z 74) obtained from <i>in situ</i> microcosms loaded with [¹³ C ₆] benzene (A) and [¹³ C ₇] toluene (B). The <i>in situ</i> microcosms were both exposed in the upper quaternary aquifer (P10F) at the field site Gneisenau (Germany) over 12 weeks.	57
Fig. 3-6: SSCP Gel of PCR amplified 16S rDNA isolated from BACTRAPs loaded with ¹³ C ₇ toluene or ¹³ C ₆ benzene after exposure (84 d) in a contaminated aquifer. The bands (numbers) are sequenced for identification (see Tab. 3-3).....	60
Fig. 3-7: Phylogenetic analysis (maximum likelihood method) of PCR-amplified partial 16S rRNA genes from the group Deltaproteobacteria, retrieved from BACTRAPs exposed at the field site Gneisenau. Clones are indicated by their source of isolation (P10F for upper and P11T for lower aquifer) followed by their clone number. * indicates sequences which were found at two subsequent sampling dates in the same well.....	63
Fig. 3-8: GC-C-IRMS-chromatogram of the carboxylic acid fraction extracted from a [¹³ C]- α -toluene amended BACTRAP incubated at the field site Zeitz (Germany). The metabolites gave very strong signals in the 45/44 ratio plot (A) although their absolute concentration displayed by the amount of CO ₂ (m/z 44) was relatively low (B).....	64
Fig. 3-9: Mass spectra of benzylsuccinate extracted from <i>in situ</i> microcosms incubated 92d within an anaerobic aquifer at the field site Zeitz (Germany). (A) BACTRAP loaded with non labelled toluene, (B) BACTRAP loaded with [¹³ C]- α -toluene. The fragmentation pattern of ¹³ C labelled	

Abbildungsverzeichnis

benzylsuccinate shows that the ^{13}C atom was found at the position of the former methyl group (C).....	65
Fig. 4-1: Approach I: Assessment of the cumulated biodegradation in a contaminated aquifer by use of stable isotope fractionation analysis (SIFA) at natural abundance; for details see text.....	77
Fig. 4-2: A) Spatial distribution of the benzene contaminations in the wells of the Zeitz aquifer in 2002 and 2004 (X = location of sampling wells; Saf Zz 32/02 well of the <i>in situ</i> microcosm experiment). B) Spatial distribution of the carbon isotope composition of benzene in the wells of the Zeitz aquifer.....	79
Fig. 4-3: Percentage of benzene degradation in the wells of the contamination plume calculated by use of equation 3 (isotopic composition of the source $[R_0] = -28.5\text{\textperthousand}$, $\alpha = 1.0020$ (methanogenic) – 1.0036 (sulfate reducing); modified from (Fischer et al., 2004)).....	80
Fig. 4-4: Approach II (A): Assessment of biodegradation potentials in a contaminated aquifer by use of <i>in situ</i> microcosms (BACTRAPs) with tracer compounds labeled with stable isotopes; (B): microcosm containing Bio-Sep® beads. (Picture: Copyright (2003) from Environmental Forensics, 2003, 4(1) by White et al.; reproduced by permission of Taylor & Francis Group, LLC., http://www.taylorandfrancis.com).....	81
Fig. 4-5: GC-MS profiles and isotope enrichment of fatty acids in total lipid extracts obtained from the $[^{13}\text{C}_1]$ - α -toluene (A) and $[^{13}\text{C}_6]$ benzene (B) amended microcosms deployed in well Saf Zz 32/02 in the Zeitz aquifer for 32 days. Major fatty acids with retention time in brackets given in minutes were 14:0 (13.83), i15:0 (15.46), a15:0 (15.66), 15:0 (16.42), i16:0 (17.98), 16:1w7c (18.36), 16:0 (18.91), 18:1w9c (22.96), 18:1w7c (23.10), 18:0 (23.59), br16:0 (19.92, 20.41, 20.60) with undetermined position of methyl-branching, * phthalate (21.31) from extraction process. (The $\delta^{13}\text{C}$ values for the isotopic composition of the respective methyl esters represent the isotopic shift compared to the PDB standard.).....	83
Fig. 4-6: SSCP Gel (A) and digital image analyses of SSCP-profiles (B) of PCR amplified 16S rDNA isolated from groundwater and microcosms loaded with toluene or benzene and controls (blanks) after exposure for 24 d (T1), 49 d (T2), and 92 d (T3) in well Saf Zz 32/02. Selected bands of these profiles were sequenced for identification, numbers indicated in the gel image can be found in Tab. 4-2.....	87
Fig. 4-7: Monitoring wells and distribution of the monochlorobenzene plume in Bitterfeld modified from (Kaschl et al., 2005). Concentrations are given by the dotted lines in mg L^{-1} . The arrows indicate the groundwater flow in November 2002. The solid line (A to B) represents the position of the cross-section perpendicular to the monochlorobenzene plume at the location of the SAFIRA research station. (Saf 4 = monitoring well for the vertical analysis).....	89
Fig. 4-8: Monochlorobenzene concentrations and isotopic signatures of the cross-section A – B, depth of 17-21 m (left) and the vertical profile in Saf 4 (right); (modified from Kaschl et al., 2005); ● = MCB concentrations; □ and ○ = $\delta^{13}\text{C}$ of MCB	90
Fig. 4-9: Rayleigh plot illustrating field-derived and laboratory-derived fractionation factors: aerobic degradation of MCB in batch experiments ($\alpha=1.0001-1.00028$), the field fractionation factor for the whole MCB plume ($\alpha_f=1.00043$), the (▲) left (Saf11, Saf12, Saf27, Saf28) and (●) right fringe (Saf7, Saf30, Saf31, Saf23) of the plume at the cross-section A - B ($\alpha_f=1.00083$ and 1.00106, respectively) and, for comparison, batch experimental data for the reductive dechlorination of TCBs ($\alpha=1.0031-1.0037$ (Griebler et al., 2004a) and anaerobic degradation of benzene ($\alpha=1.0019-1.0036$ (Mancini et al., 2003)).....	91
Fig. 4-10: Total ion current of the fatty acid extracts from a BACTRAP amended with $[^{13}\text{C}_6]$ monochlorobenzene exposed for 49 d in an anaerobic aquifer in well Saf 4/97 at the Bitterfeld field site. Identified fatty acids and their isotope compositions are indicated in the boxes.	93
Fig. 5-1: A). Multilevel sampling system (MLPS), preventing groundwater circulation and penetration of oxygen from the surface, used for the <i>in situ</i> microcosm experiment in well SAF4. The black rectangles indicate the location of the <i>in situ</i> microcosms (m1-m6) in general between the filter screen of the well and the membrane of the MLPS, whereas the small rectangles (I-IV) show the detailed experimental set up with $[^{12}\text{C}]$ MCB (I), Blank (II), $[^{13}\text{C}_6]$ MCB (III) and Blank (IV) <i>in situ</i> microcosms as it was performed in every depth. The black arrows show the groundwater flow direction, gs: ground surface and wt: water table. B). Carbon isotope composition of fatty acid methyl esters (FAME) extracted from $[^{13}\text{C}_6]$ MCB	

amended <i>in situ</i> microcosms incubated for 7 weeks at different depth (m1-m6) at the multilevel monitoring well SAF4 in Bitterfeld. The grey shaded area indicates the natural abundance of $\delta^{13}\text{C}_{\text{FAME}}$. C) Relative abundance of total lipid fatty acids [FAME] extracted from <i>in situ</i> microcosm loaded with $^{13}\text{C}_6$ -labelled MCB incubated in different depths (m1-m6) at the monitoring well SAF4 in Bitterfeld.	99
Fig. 5-2: Carbon isotope signature of CO_2 (♦) and methane (○) in A) a microcosm prepared with groundwater from SAF11 and B) the enrichment culture. Based on the CO_2 content and isotope signature, the amount of $[^{13}\text{C}_6]$ MCB degraded was calculated in μmol per microcosm (○) and % remaining of the total added $[^{13}\text{C}_6]$ MCB (■) for C) the microcosm and D) the enrichment culture. The shaded area in panel A indicates the observed isotope signature of CO_2 in the control microcosms.	103
Fig. 5-3: Total ion current of fatty acid methyl esters derived from microcosms prepared with aquifer water of well SAF30. Shown are samples from a killed control, natural abundance MCB amended and $[^{13}\text{C}_6]$ MCB amended microcosms. Identified peaks are labeled and underlined (—). FAME were identified via GC-MS and the isotope signature in the various samples are shown in $\delta\text{‰}$ [PDB].....	105
Fig. 6-1: Concentrations [$\mu\text{g L}^{-1}$] and carbon isotope signatures [%] of total chlorinated benzenes (numbers) at the study site for sampling in 2005. The letters indicate the name of the wells and the rectangle in diagonal crosses surrounds the area of the containment, which is hydraulically disconnected from the plume. Additionally the dashed (---) black line shows the area where the ratio between MCB and DCB is < 0.3 and the solid (—) line indicates where the ratio between MCB and DCB is > 0.9 . The symbols indicate the samples which belong to a common cluster as obtained by PCA: ▲-cluster I (A, B, C, D); ■-cluster II (G, H, N, O); ●-cluster III (E, F, I, J); ♦-cluster IV (J, M) (see Fig. 6-3). Wells which were not considered for PCA are indicated by ⊗.....	118
Fig. 6-2: Correlation between concentration [$\mu\text{g L}^{-1}$] and isotopic enrichment [%] of total chlorinated benzenes. The symbols indicate the sampling locations which belong to a common cluster as obtained by PCA: ▲-cluster I (A, B, C, D); ■-cluster II (G, H, N, O); ●-cluster III (E, F, I, J); ♦-cluster IV (J, M) (see Fig. 6-3).	119
Fig. 6-3: PCA ordination plot of geochemistry, contaminant concentrations, carbon isotope composition of MCB and total CBs in groundwater samples obtained during sampling in 2005. Values on the axes indicate % of total variation explanation by the corresponding axis (PC1, principal component axis 1; PC 2, principal component axis 2). The symbols indicate the samples which belong to a common cluster: ▲-cluster I (A, B, C, D); ■-cluster II (G, H, N, O); ●-cluster III (E, F, I, J); ♦-cluster IV (J, M). The figure in the upper right corner of the plot depicts the relative contribution of the descriptors in the reduced space. Description vectors correspond to: $\delta^{13}\text{C}_{\Sigma\text{CB}}$, isotope balance of total CBs; $\delta^{13}\text{C}_{\text{MCB}}$, carbon isotope signature of MCB; Fe^{2+} , ferrous iron; EC, electric conductivity; TOC, total organic carbon; CH_4 , methane; Cl^- , chloride; MCB, monochlorobenzene; ΣCB , total chlorobenzenes; NH_4^+ , ammonium; Eh, redox potential; SO_4^{2-} , sulphate; ΣMn , total manganese.....	120
Fig. 7-1: Ordination plot generated by principal component analysis representing the relationship between the sampling locations and the average geochemical parameters measured in both the planted (a) and the unplanted (b) experimental wetlands over the study period. Description vectors correspond to: Fell = ferric iron; Tot.Fe = Total iron ($\text{Fe}(\text{II})+\text{Fe}(\text{III})$); MCB = Monochlorobenzene; MnII = Manganese(II); NH4. = Ammonium; T = Temperature. Objects correspond to: 0.5 to 4 = distance [m] from the inflow point; u = upper depth (30cm); m = medium depth (40cm); l = lower depth (50cm). Values on the axes indicate % of total variation explained by the axes. PC 1 = principal component axis 1; PC 2 = principal component axis 2.....	132
Fig. 7-2: Concentration and isotopic composition of MCB (diamond = concentration; squares = isotopic composition) in both planted (a) and unplanted (b) segments and for both sampling dates (line = 14.04.2005, dashed = 09.05.2005). Error bars show the standard deviation.	133
Fig. 7-3: GC-MS chromatogram (total ion current) of the <i>in situ</i> microcosm extract showing the ^{13}C -labelled benzene metabolite at retention time of 1.73 min and corresponding mass spectrum (a) with the molecule mass of $m/z = 84$ for $[^{13}\text{C}_6]$ benzene and $m/z 78$ for $[^{12}\text{C}]$ benzene (b).	135
Fig. 7-4: Absolute abundance of extracted fatty acids [$\mu\text{g FA}$ per gram Bio-Sep [®] beads] from <i>in situ</i> microcosms exposed in the soil compartment at different distance from the inflow and ponds at the unplanted (a) and planted (b) segments of the constructed wetland.	136

Abkürzungsverzeichnis

Abkürzungsverzeichnis

ALA	Altlastenausschuss
B [%]	prozentuale Biodegradation
BACTRAP®	Bakterienfalle (<i>BACterial TRAP</i>), Testsystem zum Nachweis der <i>in situ</i> Biodegradation
BAME	bakterielle Fettsäuremethylester (<i>bacterial acid methyl ester</i>)
BBodSchG	Bundes-Bodenschutzgesetz
BGR	Bundesanstalt für Geowissenschaften und Rohstoffe
BMBF	Bundesministerium für Bildung und Forschung
BTEX	Benzol, Toluol, Ethylbenzol, Xylole
C ₀ , C _t	Konzentration einer Verbindung zu Beginn (t = 0) oder zu einer bestimmten Zeit (t) der mikrobiellen Umsetzung
C _B	an einer Messstelle zu erwartende Schadstoffkonzentration, die durch mikrobiellen Abbau beeinflusst wurde
C _(x)	Konzentration einer Verbindung (x)
C _n	Anzahl der Kohlenstoffatome einer Fettsäure
C _{Schadstoff}	mineralisierte Schadstofffraktion
CB	Chlorbenzole
CKW	chlorierte Kohlenwasserstoffe
CSIA	Substratspezifische Isotopenfraktionierungsanalyse (<i>compound-specific stable isotope analysis</i>)
DBU	Deutsche Bundesstiftung Umwelt
DCB	Dichlorbenzol
DECHEMA	Gesellschaft für Chemische Technik und Biotechnologie
DEHP	Diethylhexylphthalat
DEV	Deutsches Einheitsverfahren
DGGE	denaturierende Gradientengelektrophorese
DIC	anorganischer, gelöster Kohlenstoff (<i>dissolved inorganic carbon</i>)
DIN	Deutsches Institut für Normung
DNA	Desoxyribonukleinsäure (<i>deoxyribonucleic acid</i>)
DNAPL	Phasen mit einer Dichte > 1 (<i>dense non aqueous phase liquid</i>)
DOC	gelöster, organischer Kohlenstoff (<i>dissolved organic carbon</i>)
ENA	Stimulation der natürlichen Selbstreinigung (<i>Enhanced Natural Attenuation</i>)
Eq.	Gleichung (<i>equation</i>)
FA	Fettsäure (<i>fatty acid</i>)
FAME	Fettsäuremethylester (<i>fatty acid methyl ester</i>)
FD	Filmdicke der GC-Säule

FISH	Fluoreszenz- <i>in-situ</i> -Hybridisierung
GC-C-IRMS	Gaschromatographie-Verbrennungs-Isotopenverhältnis-Massenspektroskopie (<i>Gas chromatography-combustion-isotope ratio mass spectrometry</i>)
GC-FID	Gaschromatographie-Flammenionisationsdetektor
GC-IRMS	Gaschromatographie-Isotopenverhältnis-Massenspektroskopie (<i>Gas chromatography-isotope ratio mass spectrometry</i>)
GC-MS	Gaschromatographie-Massenspektroskopie
GI.	Gleichung
GSF	Helmholtz Zentrum München, Deutsches Forschungszentrum für Umwelt und Gesundheit
GW	Grundwasser
GWL	Grundwasserleiter
HCH	Hexachlorcyclohexan (Lindan)
IAEA	Internationale Atomenergieorganisation (<i>International Atomic Energy Agency</i>)
ID	Innendurchmesser
ITVA	Ingenieurtechnischer Verband Altlasten
IMW	Innovative Messtechnik Weiss
KORA	Kontrollierter natürlicher Rückhalt und Abbau von Schadstoffen bei der Sanierung kontaminiert Böden und Grundwässer
LABO	Bund-/Länderarbeitsgemeinschaft Bodenschutz
LHKW	leichtflüchtige, halogenierte Kohlenwasserstoffe
MCB	Monochlorbenzol
MKW	Mineralölkohlenwasserstoffe
MNA	Überwachung der natürlichen Selbstreinigung (<i>Monitored Natural Attenuation</i>)
MLPS	Multilevel-Packersystem
m uGOK	Meter unter Geländeoberkante
MPN	Wahrscheinlichste Keimzahl Methode (Most-Probable-Number-Methode)
MTBE	Methyl- <i>tert</i> -butylether
NA	Natürliche Selbstreinigung (<i>Natural Attenuation</i>)
n.d.	nicht bestimmt (not determined)
p. A.	Pro analysi (zur Analyse, Reinheitsgrad der Chemikalien)
PAC	Aktivkohle (<i>powdered activated carbon</i>)
PAH	polyzyklische aromatische Kohlenwasserstoffe (<i>polycyclic aromatic hydrocarbon</i>)
PAK	polyzyklische aromatische Kohlenwasserstoffe
PCA	<i>Principal Component Analysis</i>
PC1/2	<i>Principal Component 1/2</i>
PCE	Tetrachlorethen (Perchlorethen)

Abkürzungsverzeichnis

PCR	Polymerase-Kettenreaktion (<i>polymerase chain reaction</i>)
REACH	Verordnung zur Registrierung, Bewertung, Zulassung und Beschränkung chemischer Stoffe (<i>Registration, Evaluation, Authorisation of Chemicals</i>)
PLFA	Phospholipid-Fettsäure (<i>phospholipid-derived fatty acid</i>)
RDH	reduktive Dehalogenierung
RNA	Ribonukleinsäure (<i>ribonucleic acid</i>)
rRNA	ribosomale Ribonukleinsäure (<i>ribosomal ribonucleic acid</i>)
R_0, R_t	Isotopenverhältnis einer Verbindung zu Beginn ($t = 0$) oder zu einer bestimmten Zeit (t) der mikrobiellen Umsetzung
$R_{(x)}$	Isotopenverhältnis einer Verbindung (x)
SAFIRA	Sanierungsforschung in regional kontaminierten Aquiferen
SI	<i>Supporting Information</i>
SIFA	Analyse der Fraktionierung stabiler Isotope (<i>stable isotope fractionation analysis</i>)
SIP	Stabile Isotopenbeprobung (<i>stable isotope probing</i>)
SSCP	<i>Single Strand Conformation Polymorphism</i> (molekulares Nachweisverfahren)
Tab.	Tabelle
TCB	Trichlorbenzol
TEA	terminaler Elektronenakzeptor (<i>terminal electron acceptor</i>)
TLFA	<i>Total Lipid Fatty Acid</i>
TMCS	Trimethylchlorosilane
TOC	gesamter organischer Kohlenstoff (<i>total organic carbon</i>)
UFZ	Helmholtz-Zentrum für Umweltforschung
US-EPA	US-amerikanische Umweltbehörde (United States Environmental Protection Agency)
V-CTD	Vienna Canyon Diabolo Troilite
V-PDB	Vienna PeeDee Belemnite
V-SMOC	Vienna Standard Mean Ocean Chloride
V-SMOW	Vienna Standard Mean Ocean Water
WHO	Weltgesundheitsorganisation (<i>World Health Organization</i>)
$\gamma_{x/s}$	Ertragskoeffizient
Z_c	Anzahl der Kohlenstoffatome des Schadstoffmoleküls
α	Isotopenfraktionierungsfaktor
$\delta^{13}\text{C}_{(x)}$	Kohlenstoffisotopenverhältnis einer Verbindung (x)
ε	Anreicherungsfaktor
$\varepsilon_{\text{DIC-CH}_4}$	Anreicherungsfaktor aus der Differenz zwischen der Kohlenstoffisotopensignatur des DIC und Methans

Kurzfassung

Die natürlichen Abbau- und Rückhalteprozesse (*Natural Attenuation*) gewinnen in der Altlastenpraxis zunehmend an Bedeutung. Insbesondere an großräumig kontaminierten Standorten (*Megasites*), wo Kosten und Nutzen herkömmlicher Sanierungsmaßnahmen oft in keinem angemessenen Verhältnis stehen, wird *Monitored Natural Attenuation* (MNA) immer häufiger zur Überwachung und Sicherung von Altlastenstandorten in Betracht gezogen. Neben den abiotischen Prozessen ist hierbei vor allem die *in situ* Biodegradation von wesentlicher Bedeutung, da der mikrobielle Abbau zu einer nachhaltigen Reduzierung der Schadstoffe im Grundwasser beiträgt und oft die wichtigste Schadstoffsenke darstellt. Die Implementierung von MNA als Standortsicherungskonzept erfordert eine umfangreiche Charakterisierung des natürlichen Selbstreinigungspotenzials von Grundwasserleitern, wobei der Nachweis der mikrobiellen *in situ* Aktivität von zentraler Bedeutung ist.

Zur Erfassung der *in situ* Biodegradation in kontaminierten Aquiferen gibt es eine Vielzahl verschiedener Methoden, wobei der Großteil dieser konventionellen Verfahren nur einen indirekten Nachweis des mikrobiellen Abbaupotenzials ermöglicht. Mit der Entwicklung eines neuen Testsystems, sogenannten *in situ* Mikrokosmen (BACTRAP®), steht neben der substratspezifischen Isotopenanalyse (CSIA) und *in situ* Tracerversuchen ein weiteres Verfahren für den direkten Nachweis des mikrobiellen Schadstoffabbaus zur Verfügung.

Die Aufgaben dieser Promotion bestanden insbesondere in (I) der Validierung des Testsystems zur Beurteilung der *in situ* Biodegradation, (II) der Anwendung von *in situ* Mikrokosmen an verschiedenen Feldstandorten und für unterschiedliche Schadstoffe zur Evaluierung des Verfahrens mithilfe alternativer Methoden, (III) der Untersuchung des intrinsischen Abbaupotenzials persistenter Schadstoffe, (IV) der Weiterentwicklung der Methode zur Analyse der am Schadstoffabbau beteiligten mikrobiellen Konsortien und (V) zur Rekonstruktion von Abbauwegen mithilfe von Metabolitenuntersuchungen. Darüber hinaus stellte die (VI) Implementierung des neuen Testsystems in die Altlastenpraxis zum Nachweis der *in situ* Biodegradation im Kontext von *Natural Attenuation* einen weiteren Schwerpunkt dieser Arbeit dar.

Durch den Einsatz ^{13}C -markierter Substrate als Tracer in *in situ* Mikrokosmen konnte die Biodegradation persistenter Verbindungen direkt im kontaminierten Aquifer innerhalb weniger Monate nachgewiesen werden. Das Testsystem wurde im Rahmen dieser Promotion erfolgreich an verschiedenen Standorten zum Nachweis des anaeroben Abbaus von Benzol, Toluol, Monochlorbenzol (MCB) sowie verschiedener polzyklischer aromatischer Kohlenwasserstoffe (PAK) eingesetzt. Ein Großteil der Feldversuche erfolgte in enger Zusammenarbeit mit Kooperationspartnern aus der Industrie.

Für die Charakterisierung des anaeroben Benzolabbaus sowie zur Validierung der *in situ* Mikrokosmen wurde am Standort eines ehemaligen Hydrierwerks die vertikale Struktur einer

Kurzfassung

Benzolfahne im Rahmen eines Multilevel-Experimentes mithilfe geochemischer und isotopenchemischer Methoden untersucht (Kapitel 2). Die *einzelnen* Ergebnisse der verschiedenen Methoden zeigten generell gute Übereinstimmung hinsichtlich einer Abschätzung des *in situ* Abbaupotenzials. Im hoch kontaminierten Bereich der Fahne konnte der Benzolabbau mit der Methode der Isotopenfraktionierung jedoch nicht nachgewiesen werden. Nur der Einsatz ¹³C-markierter Substrate in *in situ* Mikrokosmen ermöglichte einen sensitiven Nachweis der Biodegradation in allen Zonen der Fahne.

Unter zu Hilfenahme von DNA Fingerprinting Methoden konnte gezeigt werden, dass die mikrobielle Gemeinschaft auf den *in situ* Mikrokosmen im Wesentlichen der Grundwassermikroflora des BTEX kontaminierten Aquifers entspricht (Kapitel 3, 4). Weitere Feldversuche zur Charakterisierung der Benzol- und Toluol abbauenden Mikroorganismen verdeutlichen, dass *in situ* Mikrokosmen geeignet sind, um in Kombination mit Nukleinsäure basierenden molekularbiologischen Methoden empfindlich die mikrobielle Gemeinschaft in Übereinstimmung mit der regionalen Geochemie des Aquifers abzubilden. Die Analyse der Fettsäremuster hingegen ist für taxonomische Fragestellungen ungeeignet.

Ein weiterer Indikator für *in situ* Abbauaktivität ist der Nachweis charakteristischer ¹³C-markierter Metabolite, anhand derer Aussagen bezüglich der Abbaumechanismen möglich sind. In diesem Zusammenhang konnte Benzylsuccinat, als typischer Metabolit des anaeroben Toluolabbaus (Kapitel 3) sowie Benzol als Intermediat der reduktiven Dehalogenierung von MCB (Kapitel 7) von *in situ* Mikrokosmen extrahiert werden. Durch die Verwendung sorptiver Trägermaterialien wie Aktivkohle können sich Metabolite auf dem *in situ* Testsystem anreichern. Die Analyse des Metabolitenpektrums bietet daher das Potenzial, neben dem Nachweis des Schadstoffabbaus auch bisher noch unbekannte Abbauwege zu erforschen.

Mithilfe von Isotopenmethoden konnte unter anoxischen Bedingungen erstmals die vollständige Mineralisierung von MCB in Laborabbauversuchen nachgewiesen werden (Kapitel 5, 6). Des Weiteren ermöglichen *in situ* Mikrokosmen den Nachweis des anaeroben MCB Abbaus unter *in situ* Bedingungen an verschiedenen Standorten (Kapitel 5-7). MCB diente dabei als Modellsubstanz für schwer abbaubare Verbindungen. Die Untersuchungen am Standort Bitterfeld erfolgten in einem Multilevel-Experiment (Kapitel 5) sowie in einer Pflanzenkläranlage, wobei hier die *in situ* Mikrokosmen erstmals direkt im Sedimentkörper inkubiert wurden (Kapitel 7). Daraus lässt sich ableiten, dass das BACTRAP-Testsystem auch direkt in Böden und Sedimenten eingesetzt werden kann.

Der direkte Nachweis des *in situ* Abbaus einzelner Schadstoffe ist mithilfe stabiler Isotopenmethoden wie CSIA und *in situ* Mikrokosmen auch an Standorten mit komplexem Kontaminationsmuster möglich. Für die Charakterisierung des *Natural Attenuation* Potenzials an einem Chlorbenzol kontaminierten Standort wurde hierfür ein integratives Untersuchungsverfahren entwickelt (Kapitel 6). Da die Anwendung des Rayleigh Konzepts zur

Abschätzung der *in situ* Biodegradation anhand der Änderung der Isotopensignaturen für sequenzielle Abbauwege wie die reduktive Dehalogenierung limitiert ist, wurde eine Isotopenbilanz erstellt. Neben den innovativen Isotopentechniken diente die statistische Auswertung umfangreicher Datensätze zur Charakterisierung der heterogenen hydrogeochemischen Bedingungen. Trotz der komplexen Standortbedingungen konnten mithilfe dieses integrativen Ansatzes entsprechend bestehender nationaler und internationaler Richtlinien (Hessisches Landesamt für Umwelt und Geologie, 2004; ITVA, 2004; KORA, 2007; US-EPA, 1999) mehrere Beweislinien für den Nachweis der *in situ* Abbauprozesse erbracht werden.

Mit dieser Arbeit wurde gezeigt, dass *in situ* Mikrokosmen geeignet sind, um die Aktivität, Struktur und Funktion von am Schadstoffabbau beteiligten Mikroorganismen unter Einsatz ^{13}C -markierter Tracerverbindungen mit hoher Sensitivität nachzuweisen. Die Methode empfiehlt sich insbesondere für die Analyse der sehr langsam ablaufenden anaeroben Abbauprozesse persistenter Verbindungen wie z.B. MCB und PAKs, da es kaum alternative Nachweisverfahren gibt. Damit steht neben der komponentenspezifischen Isotopenanalyse (CSIA) und *in situ* Tracerversuchen eine weitere innovative Methode für die Erfassung der *in situ* Biodegradation bei der Implementierung von *Natural Attenuation* als Standortsicherungskonzept in der Altlastenpraxis zur Verfügung. Der Einsatz des Testsystems ist in konventionell ausgebauten und Multilevel Messstellen sowie direkt im Sedimentkörper von zum Beispiel Pflanzenkläranlagen möglich. Die Methode der *in situ* Mikrokosmen wurde bereits im KORA-Leitfaden sowie der Methodensammlung über mikrobiologische NA-Untersuchungsverfahren (KORA, 2007; KORA, 2008) integriert und im Einverständnis mit den verantwortlichen Umweltbehörden an verschiedenen Standorten für den geforderten Nachweis der *in situ* Biodegradation im Rahmen von MNA Vorhaben angewendet.

Abstract

Abstract

The natural ability to remove contaminants by microbial degradation is increasingly used as remedial alternative especially at heavily contaminated field sites (megasites) where active remediation technologies are often technically and economically not feasible. The implementation of Monitored Natural Attenuation (MNA) requires a careful evaluation of the ongoing biogeochemical processes under *in situ* conditions. The Natural Attenuation processes including various chemical-, physical- and biological processes may result in the reduction of the mass, toxicity, mobility, volume, or concentration of contaminants in the aquifer system. Beside the abiotic processes such as sorption, volatilization or dilution, *in situ* biodegradation is the only process leading to a sustainable removal of contaminants and therefore, microbial degradation is the most important sink of contaminants at contaminated field sites. Consequently, the implementation of Natural Attenuation as remediation strategy requires a careful site characterization with special attention regarding the *in situ* biodegradation processes.

Today, many different methods to determine the *in situ* biodegradation are available. Most of these conventional techniques only allow to indirectly assess the microbial degradation potential. Since the development of *in situ* microcosms (BACTRAP®), one further method is now available to directly provide evidence of *in situ* degradation beside compound-specific stable isotope analysis (CSIA) and *in situ* tracer tests.

Main objectives of this thesis were (I) to validate the new *in situ* microcosm test system for the assessment of *in situ* biodegradation, (II) the application of *in situ* microcosms at different field sites with different contaminants to evaluate the method with alternative techniques, (III) the investigation of the intrinsic degradation potential of persistent contaminants, (IV) the further development of the method to enable the analyses of the microbial consortia involved in contaminant degradation and (V) the reconstruction of degradation pathways by means of metabolite analyses. Additionally, (VI) the acceptance and implementation of the test system to determine *in situ* biodegradation within the context of Natural Attenuation as management strategy for contaminated field sites was addressed.

The application of *in situ* microcosms amended with ¹³C-labelled substrates enabled the direct detection of *in situ* biodegradation of persistent compounds inside the contaminated aquifer within only a few months. During this thesis, the test system was successfully applied at different field sites to determine the anaerobic degradation of benzene, toluene, monochlorobenzene (MCB) as wells as of different polycyclic aromatic hydrocarbons (PAH). Many field studies were performed in close collaborations with industrial companies.

To characterise the anaerobic benzene degradation and to validate the *in situ* microcosm test system the vertical structure of a benzene plume at a former hydrogenation plant was analysed by means of geochemical and isotope chemical methods (chapter 2). All results were in good

agreement. At the highly contaminated zone of the plume the results of the CSIA did not show benzene degradation. Only the use of ¹³C-labelled tracers in *in situ* microcosms enabled a very sensitive detection of the anaerobic degradation processes in all parts of the plume.

Using DNA fingerprinting methods it was shown that the microbial community colonizing the *in situ* microcosms mainly represents the groundwater microflora (chapter 3, 4). Further field studies to characterise the benzene and toluene degrading microorganisms revealed that the test system in combination with nucleic acid based molecular methods is appropriate to sensitively reflect the microbial community in accordance with the local geochemistry. The analysis of fatty acids alone was not sufficient for taxonomic interpretation.

Another indicator of *in situ* biodegradation is the proof of characteristic ¹³C-labelled metabolites, which can also be used to elucidate degradation mechanisms. In this context, benzylysuccinate, a well known metabolite of the anaerobic toluene degradation (chapter 3) and benzene as intermediate of the reductive dehalogenation of MCB (chapter 7) were extracted and identified from *in situ* microcosms. The use of activated charcoal-like material allows the enrichment of metabolites on the test system. Analysing metabolite spectra will facilitate the reconstruction of metabolic pathways.

Using stable isotope techniques, for the first time complete mineralisation of MCB under anoxic conditions was proven (chapter 5, 6). Furthermore, anaerobic microbial degradation of MCB under *in situ* conditions was shown at different field sites applying the *in situ* microcosm test system (chapter 5-7). During these experiments MCB was considered as model compound representing persistent contaminants. The investigations at the field site in Bitterfeld were performed using a multilevel sampling approach (chapter 5) and inside a constructed wetland system, where for the first time *in situ* microcosms were directly incubated inside the sediment (chapter 7). Accordingly, it can be concluded that the test system is suitable for direct applications inside soils or sediments.

The direct proof of microbial degradation of single contaminants inside a multi component mixture is feasible using stable isotope techniques such as CSIA or *in situ* microcosms, even at highly complex field sites. To characterise the Natural Attenuation potential at a chlorobenzene contaminated aquifer an integrated approach was developed (chapter 6). Because the application of the Rayleigh concept to estimate the *in situ* biodegradation on the basis of the change in the isotope composition is limited for sequential degradation pathways, an isotope balance was determined. Beside the innovative isotope techniques multivariate statistics were used to analyse the data for the interpretation of the heterogeneous hydrogeochemical aquifer conditions. Despite the highly complex site conditions the integrative approach enabled to provide multiple lines of evidence for the ongoing *in situ* degradation processes as required by the authorities if Natural Attenuation will be considered as remediation option.

Abstract

To summarize, in this thesis it is shown that the *in situ* microcosm test system, based on the use of ¹³C-labelled tracer compounds, can be applied to sensitively determine the activity, structure and function of the microbial community involved in contaminant degradation within a reasonable time frame. The method is in particular recommended for the analysis of the very slow degradation processes of persistent compounds such as MCB and PAHs, since almost no alternative methods are available. Beside the CSIA and *in situ* tracer studies, now a further innovative method exists to prove *in situ* biodegradation according existing national and international guidelines (Hessisches Landesamt für Umwelt und Geologie, 2004; ITVA, 2004; KORA, 2007; KORA, 2008; US-EPA, 1999) for the implementation of Natural Attenuation as monitoring strategy for contaminated field sites. The test system can be used in conventional wells and in combination with multilevel sampling devices as well as directly inside the sediment as shown in investigations inside a constructed wetland. The concept of *in situ* microcosms to assess *in situ* biodegradation is already implemented in the "KORA guideline for Natural Attenuation investigation methods" (KORA, 2007; KORA, 2008) and was applied in agreement with the responsible environmental agencies at several field sites to provide the required information concerning the *in situ* degradation potential within the framework of Natural Attenuation.

1 Einleitung

1.1 Altlastenproblematik

Wasser gilt als Lebenselixier und die Verfügbarkeit von Wasser bildet einen entscheidenden Faktor für die Verbesserung der Lebensqualität sowie der ständig voranschreitenden wirtschaftlichen Entwicklung der Menschheit (WHO, 2006). Daher stellt die Versorgung der Weltbevölkerung mit hygienisch und toxikologisch unbedenklichem Trink- und Nutzwasser eine der größten Herausforderungen für die Zukunft dar. Die Nutzung von Grundwasservorkommen ist in vielen Gebieten der Erde wesentlicher Bestandteil der Wasserversorgung. In Deutschland stellt das Grundwasser sogar die Hauptquelle für die Trinkwassergewinnung dar. Jährlich werden rund 4 km^3 aus den Grundwasserleitern gefördert (BGR, Stand 2004). Seit der Industrialisierung im 19. Jahrhundert wurden zunehmend umweltrelevante Stoffe in den Untergrund eingetragen. Ein ausgeprägtes Umweltbewusstsein existierte zu dieser Zeit in der Öffentlichkeit noch nicht. Vor allem in Bereichen von Bergbaufolgelandschaften, ehemaligen Standorten der chemischen Industrie und Metallurgie, landwirtschaftlich intensiv genutzte Flächen sowie in Gebieten mit militärischer Nutzung gelangten eine Vielzahl von Umweltschadstoffen durch unsachgemäßen Umgang, Leckagen, Havarien oder Kriegsschäden in den Boden und schließlich in das Grundwasser. In Folge dessen finden wir heute im Bereich ehemaliger Industriestandorte eine Vielzahl von Grundwasserschadensfällen. Allein in Deutschland sind laut Umweltbundesamt rund 272.240 Altlastenverdachtstflächen registriert. Dabei handelt es sich um Standorte, bei denen der Verdacht schädlicher Bodenveränderungen oder sonstiger Gefahren für den Einzelnen oder die Allgemeinheit besteht (Umweltbundesamt, 2007). Eine spezielle Problematik stellen dabei die Standorte dar, die durch sehr großräumige Kontaminationen und umfangreiches Schadstoffinventar, so genannte Megasites, charakterisiert sind (Schirmer et al., 2006). Die Altlastenproblematik ist nicht nur ein deutsches oder europäisches Problem, sondern von globaler Relevanz.

1.2 Relevante Schadstoffe

Die bedeutendsten Grundwasserkontaminanten stellen die organischen Schadstoffe dar. Wie Arneth et al. in ihrer Studie gezeigt haben, gehören die chlorierten Kohlenwasserstoffe (CKW), im speziellen die aliphatischen, leichtflüchtigen halogenierten Kohlenwasserstoffe (LHKW) zu den am häufigsten vorkommenden Verunreinigungen in Deutschland (Arneth et al., 1989). Daneben sind vielfach auch Aromaten, wie z.B. Benzol, Toluol, Ethylbenzol und Xylool (BTEX) aber auch Chlorbenzole (CB) im Grundwasserabstrom von ehemaligen Produktionsstätten wie Hydrierwerken, Chemiefabriken, chemischen Reinigungen,

Einleitung

Imprägnierwerken u.ä. zu finden (Fig. 1-1). Des Weiteren sind polyzyklische aromatische Kohlenwasserstoffe (PAK) (Kanaly and Harayama, 2000) sowie Methyl-*tert*-butylether (MTBE) (Schmidt et al., 2002) relevante Grundwasserschadstoffe. Eine generell ähnliche Situation finden wir auch in den USA, wie die Studie von Plumb & Pitchfork (Plumb JR and Pitchford, 1985) zeigt und wie in der vergleichenden Gegenüberstellung der Daten von Deutschland und den USA zu entnehmen ist (Arneth et al., 1989; Teutsch et al., 1997).




Fig. 1-1: Ländervergleich Deutschland und USA über das Vorkommen und die Häufigkeit organischer Schadstoffe im Abstrom von Schadensfällen aus (Teutsch et al., 1997).

Ein Großteil dieser xenobiotischen Verbindungen ist durch hohe Toxizität, Persistenz sowie karzinogene Eigenschaften gekennzeichnet, womit sich ihre hohe Umweltrelevanz begründet. Neben den organischen Schadstoffen gibt es auch Verunreinigungen mit anorganischen Verbindungen (Schwermetalle, Nitrat, etc.). Den Schwerpunkt der vorliegenden Arbeit bilden jedoch die organischen Kontaminanten. Grundwasserschadensfälle mit anorganischen Verbindungen werden nicht näher behandelt. Die physiko-chemischen Eigenschaften und die Geologie eines Standortes sowie die damit im Zusammenhang stehenden Prozesse Advektion, Dispersion, Diffusion, Sorption, Verflüchtigung und Abbau kontrollieren das Ausbreitungsverhalten der Schadstoffe im Grundwasser. In Abhängigkeit von der Wasserlöslichkeit, dem Sorptionsverhalten und der biologischen Abbaubarkeit unterscheidet sich das Transportverhalten der verschiedenen Schadstoffe. Kontaminanten mit sehr guter Wasserlöslichkeit, geringer Retardation und

hoher Persistenz sind oft durch große Transportweiten gekennzeichnet, so dass sich die Schadstoffe über weite Strecken im Untergrund entlang der Grundwasserfließrichtung ausbreiten. Zu dieser Stoffgruppe zählen beispielsweise die CKW, die Fahnenlängen von mehreren Kilometern aufweisen können. Die Transportweite der BTEX hingegen ist, trotz relativ guter Wasserlöslichkeit und geringer Sorption, infolge guter Abbaubarkeit mit Fahnenlängen von meist weniger als 500 m wesentlich geringer. Die PAK hingegen zeichnen sich durch eine weitaus geringere Wasserlöslichkeit und damit verbundene stärkere Sorption an der Aquifermatrix aus, die in Abhängigkeit der Anzahl der Benzolringe variiert. Fahnenlängen von PAK Schadensfällen sind daher abhängig vom PAK-Inventar und können um mehrere Größenordnungen variieren (Teutsch et al., 1997).

Da von kontaminiertem Grundwasser ein hohes Gefährdungspotenzial für den Mensch und die Umwelt ausgehen kann, ist die Überwachung und gegebenenfalls Sanierung von großem öffentlichem Interesse. Die *in situ* Biodegradation kann wesentlich zur nachhaltigen Entfernung der Schadstoffmengen im kontaminierten Grundwasser beitragen. Deshalb ist die Charakterisierung der biogeochemischen Prozesse im Aquifer für ein differenziertes Prozessverständnis ein erster wichtiger Schritt, bevor Entscheidungen bezüglich geeigneter Sanierungsmaßnahmen bzw. Standortmanagementstrategien getroffen werden können.

1.3 Natürliche Selbstreinigung: *Natural Attenuation*

Die Berücksichtigung der natürlichen Rückhalte- und Abbauprozesse gewinnt in der Altlastenbearbeitung zunehmend an Bedeutung (Grandel and Dahmke, 2004; Rittmann, 2004). Die rechtliche Grundlage hierfür bildet in Deutschland das am 1. März 1999 in Kraft getretene Bundes-Bodenschutzgesetz (BBodSchG). Insbesondere im Fall von Schadensfällen, bei denen die Kosten und der Nutzen einer Sanierung im herkömmlichen Sinn (z.B. *Pump & Treat*, Reaktive Wände sowie *Funnel & Gate*) in keinem günstigen Verhältnis stehen und zu keiner nachhaltigen Dekontamination des Standortes führen, werden immer häufiger die natürlichen Selbstreinigungs Kräfte des Aquifers in Betracht gezogen, um eine langfristige Sicherung des Standortes zu erreichen. An eine zunächst aktive Sanierung (z.B. Quell- oder Hot-Spot Sanierung) kann sich eine passive Überwachung und Sicherung des Standortes durch *Monitored Natural Attenuation* (MNA) anschließen.

Natürliche Abbau- und Rückhalteprozesse werden unter den Begriff *Natural Attenuation* (NA) zusammengefasst und verlangsamen die Schadstoffausbreitung im Grundwasserleiter, weshalb Schadstofffahnen nur eine begrenzte Ausdehnung im Aquifer erreichen (Wiedemeier et al., 1999). Die amerikanische Umweltbehörde (US-EPA) definiert *Natural Attenuation* als *Eigenschaft eines Aquifers durch Prozesse wie biologischer Abbau, Dispersion, Verdünnung, Sorption, Verflüchtigung und/oder chemischer oder biochemischer Stabilisierung die Masse, Toxizität, Mobilität oder das Volumen der Schadstoffe so weit zu*

Einleitung

reduzieren, dass die menschliche Gesundheit und das Ökosystem nicht gefährdet sind (US-EPA, 1999).

Den Schwerpunkt von NA als Sicherungsmaßnahme bilden gezielte Untersuchungen zum Nachweis der natürlichen Abbau- und Rückhalteprozesse am jeweiligen Standort in Verbindung mit einem sich anschließenden Langzeitmonitoring (MNA). Der mikrobielle Schadstoffabbau (*intrinsic biodegradation*) ist ein wesentlicher Prozess im Zusammenhang mit NA, da sich infolge der Biodegradation nicht nur die Konzentrationen sondern auch die Schadstofffrachten im Grundwasser reduzieren. Die gezielte Stimulation des Abbaupotenzials (*Enhanced Natural Attenuation - ENA*) durch Zugabe von z.B. Elektronenakzeptoren kann die Effektivität der NA-Prozesse steigern, wie zum Beispiel an verschiedenen Tetrachloreten (PCE) kontaminierten Standorten gezeigt werden konnte (Major et al., 2002; Ramsburg et al., 2004). Eine umfangreiche Zusammenfassung zu aktuellen Entwicklungen bezüglich NA und ENA Maßnahmen wurde anhand verschiedener Standorte mit chlorierten Ethenen, aromatischen Kohlenwasserstoffen bzw. MTBE als Hauptkontaminanten in Übersichtsartikeln veröffentlicht (Farhadiana et al., 2008; Jørgensen, 2007; Scow and Hicks, 2005).

In den USA hat sich MNA als Alternative zu aktiven Sanierungsmaßnahmen speziell für Schadensfälle mit MKW bereits etabliert, was sich auch in den bestehenden Handlungsempfehlungen der amerikanischen Umweltbehörde US-EPA zur Anwendung von NA widerspiegelt (Doll, 1998; US-EPA, 1999; US-EPA, 2004; Wöstmann, 2007). In Deutschland hingegen fehlten in der Vergangenheit die rechtlichen und fachlichen Grundlagen für eine sachgerechte Implementierung und gezielte Berücksichtigung der natürlichen Selbstanreinigungs Kräfte in der Altlastensanierung (Wöstmann, 2007). NA war und ist daher Inhalt verschiedener deutscher Forschungsprojekte, wie zum Beispiel der vom BMBF geförderte Förderschwerpunkt „KORA - Kontrollierter natürlicher Rückhalt und Abbau von Schadstoffen bei der Sanierung kontaminierten Böden und Grundwässer“ oder auch verschiedene Langzeituntersuchungen des Umweltbundesamtes (BMBF, 2000). Als Ergebnis dieser umfangreichen Forschungstätigkeiten liegen heute verschiedene Merkblätter, Handlungsempfehlungen und Arbeitshilfen vor, die den verantwortlichen Behörden, Ingenieurbüros und Sanierungspflichtigen als Leitfäden und Arbeitsrichtlinien zur Verfügung stehen. In diesem Kontext sind die folgenden Schriften zu nennen, wobei diese Liste keinen Anspruch auf Vollständigkeit erhebt (Bayrisches Landesamt für Wasserwirtschaft, 2004; Hessisches Landesamt für Umwelt und Geologie, 2004; ITVA, 2004; KORA, 2007; LABO, 2005):

- Bayrisches Landesamt für Wasserwirtschaft (2004): Natürliche Schadstoffminderung bei Grundwasserverunreinigungen durch Altlasten und schädliche

Bodenveränderungen - Natural Attenuation - Merkblatt Nr. 3.8/3 vom 05.11.2004,
München, 17 S.

- Hessisches Landesamt für Umwelt und Geologie (2004): Arbeitshilfe zu überwachten natürlichen Abbau- und Rückhalteprozessen im Grundwasser (Monitored Natural Attenuation MNA), Wiesbaden, Hessisches Landesamt für Umwelt und Geologie. Handbuch Altlasten, Bd.8, T.1
- ITVA (2004): Monitored Natural Attenuation. Ingenieurtechnischer Verband Altlasten e.V., ITVA-Arbeitshilfe –H1-12, Dezember 2004, Berlin, 35 S.
- LABO (2005): Berücksichtigung natürlicher Schadstoffminderungsprozesse bei der Altlastenbearbeitung -Bund-/Länderarbeitsgemeinschaft Bodenschutz (LABO) – Altlastenausschuss (ALA) Ad-hoc Unterausschuss „Natural Attenuation“, vom 01.06.2005
- KORA (2007): Mikrobiologische NA-Untersuchungsmethoden: Fachliche Grundlagen für die Anwendung von Methoden zur Erfassung des natürlichen mikrobiellen Schadstoffabbaus im Aquifer / ProcessNet, Expertengruppe "Intrinsisches Abbaupotenzial" im Arbeitsausschuss "Ressourcenmanagement Boden und Grundwasser". - Stand: März 2007. – Frankfurt M. DECHEMA, 2007. - 73 S.

Diese Schriftreihen dienen als Richtlinien zur Berücksichtigung natürlicher Abbau- und Rückhalteprozesse in kontaminierten Aquifern in der Sanierungspraxis und sie unterstreichen die Bedeutung der mikrobiell initiierten Prozesse, die einen signifikanten Beitrag zur nachhaltigen Reduzierung der Schadstoffmengen leisten können.

1.4 Nachweis der *in situ* Biodegradation

Der Nachweis des biologischen Abbaus organischer Verbindungen ist ein wesentliches Element für die Implementierung von Natural Attenuation zur Überwachung und Sicherung kontaminierte Standorte, da nur der mikrobielle Schadstoffabbau zu einer destruktiven und vor allem nachhaltigen Reduzierung der Schadstoffmengen im Grundwasser beiträgt. Abiotische Prozesse, wie zum Beispiel die Sorption und Verflüchtigung sind für eine Gefahrenabwehr nicht unbedingt akzeptabel, da es hierbei nur zu einer Verlagerung der Kontaminanten in andere Umweltkompartimente kommt und keine tatsächliche Reduzierung der Schadstofffrachten erfolgt. Daher ist es von besonderem Interesse, geeignete Methoden zur Charakterisierung des *in situ* Abbaupotenzials kontaminierte Aquifere zu entwickeln. Einen Überblick zu traditionellen und innovativen NA-Untersuchungsverfahren gibt die Broschüre, die im Rahmen des BMBF geförderten KORA-Forschungsvorhaben entstanden ist (KORA, 2007).

Einleitung

Der Nachweis der *in situ* Biodegradation kann anhand hydrogeochemischer Indikatoren erfolgen. Dabei wird der Rückgang der Schadstoffkonzentrationen mit der Ausbildung spezifischer Redoxzonen relevanter terminaler Elektronenakzeptoren (TEA) in Beziehung gebracht. Ein gleichzeitiges Auftreten von charakteristischen Metaboliten untermauert hierbei die Existenz des *in situ* Abbaupotenzials. Des Weiteren gehören zu den mikrobiologischen Indikatoren für *in situ* Abbau (1) die Bestimmung von stoffwechselspezifischen Keimzahlen mit der Most-Probable-Number-(MPN)-Methode, (2) mikrobiologische Untersuchungen an Standortmaterial im Rahmen von Labormikrokosmenstudien, sowie (3) der Nachweis über das Vorhandensein von abbauaktiven Mikroorganismen mithilfe von „Fingerprinting“ Verfahren. Bei diesen Techniken handelt es sich um indirekte Nachweisverfahren. Der Nachweis erfolgt meist *ex situ*, also in Laborstudien, so dass eine direkte Überführung der Ergebnisse auf die realen komplexen Aquiferbedingungen schwierig bleibt. Auch die Identifizierung von Mikroorganismen von denen bekannt ist, dass sie zum Abbau von bestimmten Kontaminanten befähigt sind, ist noch kein eindeutiger Beweis, dass diese Bakterien auch tatsächlich am jeweiligen Untersuchungsstandort am Abbau beteiligt sind. Insbesondere da die Mehrheit der Mikroorganismen noch nicht kultiviert wurde (Amann et al., 1995) und auch die Abbauwege bestimmter Schadstoffe (z.B. Benzol, Monochlorbenzol) bis heute noch nicht verstanden sind, ist ein zweifelsfreier Nachweis der *in situ* Abbauaktivität mit diesen konventionellen Methoden schwierig bzw. kaum möglich. Deutlich bessere Aussagekraft und einen sehr hohen Informationsgehalt bieten die Isotopenmethoden.

Bei der Analyse der substratspezifischen Isotopenfraktionierung werden anhand der Veränderung der Isotopensignaturen eines Schadstoffes Rückschlüsse zum Abbaupotenzial gezogen und der *in situ* Abbau kann über einen Fließweg erfasst werden. Da Mikroorganismen aus energetischen Gründen bevorzugt isotopisch leichtes Substrat verwerten, kommt es in der noch nicht abgebauten residualen Schadstofffraktion zu einer relativen Anreicherung der schweren Isotope. Anhand dieser Verschiebung der natürlichen Isotopenverhältnisse kann unter Kenntnis der spezifischen Isotopenfraktionierungsfaktoren die *in situ* Biodegradation auch quantitativ erfasst werden (Meckenstock et al., 2004a; Meckenstock et al., 1999) (siehe auch Kapitel 1.4.1).

Die Charakterisierung des mikrobiellen Abbaus im Aquifer kann auch anhand von *in situ* Tracerexperimenten in einer Grundwassermessstelle (Push- und Pull Versuche) (Pombo et al., 2002) oder entlang einer Fließstrecke erfolgen. Dabei wird ein isotopisch markierter Schadstoff (z.B. Deuterium markiertes Toluol) sowie ein konservativer Tracer in das Grundwasser eingespeist. Mithilfe unterschiedlicher Konzepte wie Isotopenfraktionierung, Analyse von Mineralisierungsprodukten und Metaboliten können anschließend quantitative Aussagen hinsichtlich des Abbaus getroffen werden (Fischer et al., 2006; Reusser and Field, 2002) (siehe auch Kapitel 1.4.2).

Da der biologische Abbau die wichtigste und nachhaltigste Schadstoffsenke darstellt, ist der Nachweise der mikrobiellen Abbauprozesse von großer Bedeutung für die Charakterisierung und Bewertung des NA Potenzials eines Standortes. Wie in diesem Kapitel beschrieben wurde, existieren inzwischen eine Vielzahl verschiedener Methoden, um den Nachweis der Biodegradation zu führen und umfangreiche Aussagen hinsichtlich der Schadstoffmindernden Prozesse zu erhalten. Ein wesentlicher Nachteil, der sich mit dem Großteil der bisher beschriebenen Methoden verknüpft, ist jedoch die Tatsache, dass oftmals zwar das Abbaupotenzial analysiert wird, Indikatoren über die tatsächliche Abbauaktivität jedoch nicht geliefert werden können, denn allein das Vorhandensein von zum Abbau befähigten Mikroorganismen sagt nichts über deren tatsächliche Abbauaktivität am entsprechenden Standort aus. Mit Ausnahme der Isotopenmethoden (Isotopenfraktionierung, Tracertests) handelt es sich bei allen anderen Konzepten um so genannte indirekte Nachweisverfahren, die keine spezifischen Informationen über die *in situ* Aktivität der Mikroorganismen liefern. Nur die auf Isotopen basierenden Methoden ermöglichen den direkten Nachweis des mikrobiellen Schadstoffabbaus. Da die Durchführung von *in situ* Tracerversuchen mit enormem Kosten- und Arbeitsaufwand verbunden ist und darüber hinaus eine Genehmigung zur Einleitung von Schadstoffen benötigt wird, ist dieses Verfahren sicher nur in einzelnen Fällen für die Altlastenpraxis geeignet. Auch wenn die Methode der Isotopenfraktionierung seit den letzten Jahren stets an Bedeutung gewonnen hat, sind alternative Methoden notwendig, um den geforderten Nachweis über die *in situ* Abbauaktivität an kontaminierten Standorten zu erbringen.

In diesem Zusammenhang wurde ein weiteres direktes Testverfahren (*in situ* Mikrokosmen – BACTRAP®) entwickelt, welches auf dem Einsatz stabiler Isotope als Tracer basiert (Kapitel 1.5). In den folgenden Kapiteln soll zunächst etwas detaillierter in die Grundlagen ausgewählter Methoden (Isotopenfraktionierung, Tracerversuche, Molekularbiologische Methoden, *in situ* Mikrokosmen) eingeführt werden, bevor anschließend in den einzelnen Kapiteln dieser Promotionsarbeit der Stand der Forschung zur Anwendung von *in situ* Mikrokosmen zum Nachweis des mikrobiellen Schadstoffabbaus präsentiert und diskutiert wird (Kapitel 2 bis 7).

1.4.1 Stabile Isotopenfraktionierung

Eine Möglichkeit zur Untersuchung des *in situ* Abbaus von Schadstoffen bietet die Methode der Isotopenfraktionierung, bekannt als „stabile Isotopenfraktionierungsanalyse“ (SIFA) oder „substratspezifische Isotopenfraktionierungsanalyse“ (*compound-specific stable isotope analysis*, CSIA). Umfangreiche Übersichtsartikel bieten eine fundierte Einleitung in die Thematik, insbesondere zu den Grundprinzipien der stabilen Isotopenchemie, der Analytik sowie zur Quantifizierung des mikrobiellen Abbaus organischer Schadstoffe am Beispiel von

Einleitung

Labor- und Feldstudien (Blessing et al., 2008; Meckenstock et al., 2004a; Schmidt et al., 2004).

Die überwiegende Mehrheit der natürlich vorkommenden Elemente besitzt mehrere stabile Isotope, die sich bei gleicher Ordnungszahl hinsichtlich ihrer Massenzahlen unterscheiden. Diese Massendifferenz beruht auf der unterschiedlichen Anzahl von Neutronen im Atomkern. Für Fragestellungen bezüglich des Nachweises der *in situ* Biodegradation von organischen Schadstoffen sind insbesondere Wasserstoff- (^1H , ^2H) und Kohlenstoffisotope (^{12}C , ^{13}C) von Bedeutung, da organische Kontaminanten im Wesentlichen aus diesen beiden Elementen aufgebaut sind. Die verschiedenen Isotope eines Elementes unterscheiden sich signifikant hinsichtlich ihrer relativen Häufigkeit mit der sie in der Umwelt vorkommen (Tab. 1-1).

Tab. 1-1: Relative Häufigkeiten [%] der wichtigsten stabilen Isotope ausgewählter Elemente und Angabe der internationalen Standards (Hoefs, 1997)

Element	Isotope	Relative Häufigkeit [%]	Internationaler Standard
Wasserstoff	$^1\text{H}/^2\text{H}$ (H/D)	99,98 / 0,02	V-SMOW
Kohlenstoff	$^{12}\text{C}/^{13}\text{C}$	99,89 / 1,11	V-PDB
Stickstoff	$^{14}\text{N}/^{15}\text{N}$	99,64 / 0,36	Luft
Sauerstoff	$^{16}\text{O}/^{18}\text{O}$	99,76 / 0,02	V-SMOW
Schwefel	$^{32}\text{S}/^{34}\text{S}$	94,02 / 4,21	V-CTD
Chlor	$^{35}\text{Cl}/^{37}\text{Cl}$	75,53 / 24,47	V-SMOC

Die Methode der Isotopenfraktionierung beruht darauf, dass der mikrobielle Schadstoffabbau mit einem kinetischen Isotopenfraktionierungseffekt verbunden ist, weshalb das natürliche Isotopenverhältnis (z.B. $^{13}\text{C}/^{12}\text{C}$) eines Substrates im Verlauf der Abbaureaktion verändert wird. Die Ursache der Isotopenfraktionierung liegt in den unterschiedlichen Aktivierungsenergien (Unterschied der Nullpunktschwingungen), die benötigt werden, um chemische Bindungen zu spalten oder aufzubauen: die Spaltung (oder Bildung) einer Bindung mit schweren Isotopen erfordert höhere Aktivierungsenergien, so dass bevorzugt leichte Isotopomere einer Kontaminante während der mikrobiellen Umsetzung verwertet werden, wodurch es zu einer Anreicherung der schweren Isotopomere in der residualen, nicht abgebauten Schadstofffraktion kommt. Die daraus resultierenden Änderungen der Isotopenverhältnisse werden mit Hilfe der Gaschromatographie-Isotopenverhältnismassenspektroskopie (GC-IRMS) analytisch nachgewiesen. Derzeit sind online Messungen mit GC-IRMS für Kohlenstoff, Wasserstoff, Stickstoff und Sauerstoff möglich (Brenna et al., 1997). Die Reproduzierbarkeit der Messungen liegt für Kohlenstoffisotopenanalysen bei 0,2 bis 0,5 ‰ und für Wasserstoffisotopenanalysen bei 2 bis 5 ‰ (Meckenstock et al., 2004a). Die Isotopensignatur einer Verbindung wird in Promille [‰] relativ zu einem internationalen Standard in der so genannten Delta-Notation angegeben (Gleichung 1-1) (Coplen et al., 2006).

Gleichung (1-1)

$$\delta^{13}\text{C} [\%] = \frac{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{sample}} - \left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{VPDB-Standard}}}{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{VPDB-Standard}}} \times 1000$$

Die Rayleigh-Gleichung (Gleichung 1-2) beschreibt den mathematischen Zusammenhang zwischen der Konzentrationsänderung der mikrobiell umgesetzten Substanz in Verbindung mit der Änderung der Isotopensignatur durch den Isotopenfraktionierungsfaktor (α) (Hoefs, 1997).

Gleichung (1-2)

$$\frac{R_t}{R_0} = \left(\frac{C_t}{C_0} \right)^{\left(\frac{1}{\alpha} - 1 \right)}$$

R_0 und R_t geben die Isotopenzusammensetzung, C_0 und C_t die Konzentration des Substrats zum Zeitpunkt 0 und t der mikrobiellen Umsetzung an. Reaktionsspezifische Isotopenfraktionierungsfaktoren werden unter definierten Laborbedingungen bestimmt, um zu garantieren, dass ausschließlich die Biotransformationsreaktion die Veränderung von Konzentration und Isotopensignatur des Substrats verursachen. Eine Übersicht bereits publizierter Fraktionierungsfaktoren ist in der Datenbank der Firma Isodetect im Internet hinterlegt, die regelmäßig aktualisiert wird (www.isodetect.de).

Anhand der Verschiebung der natürlichen Isotopenverhältnisse kann unter Kenntnis der spezifischen Isotopenfraktionierungsfaktoren (α) die *in situ* Biodegradation auch quantitativ erfasst werden (Meckenstock et al., 2004a; Meckenstock et al., 1999). Der prozentuale biologische Abbau B[%] wird dabei über den Zusammenhang der Rayleigh Gleichung berechnet (Gleichung 1-3) (Meckenstock et al., 2002).

Gleichung (1-3):

$$B[\%] = \left(1 - \frac{C_t}{C_0} \right) \times 100 = \left(1 - \left(\frac{R_t}{R_0} \right)^{\left(\frac{1}{\alpha} - 1 \right)} \right) \times 100$$

CSIA wurde bereits erfolgreich an einer Vielzahl von Feldstandorten für BTEX (z.B. Griebler et al., 2004b; Mancini et al., 2002; Vieth et al., 2005), chlorierte Ethene (z.B. Hunkeler et al., 1999a; Imfeld et al., 2008; Nijenhuis et al., 2005; Sherwood Lollar and Slater, 2001) und MTBE (z.B. Kuder et al., 2005; Rosell et al., 2007b; Zwank et al., 2005) zur Charakterisierung des *in situ* Abbaupotenzials angewendet. Für abiotische Prozesse wie z.B. Sorption oder Verdünnung konnte gezeigt werden, dass sie unter den *in situ* Bedingungen nicht mit einem signifikanten Isotopenfraktionierungseffekt assoziiert sind (Kopinke et al.,

Einleitung

2005; Meckenstock et al., 1999; Slater et al., 2000). Des Weiteren ermöglicht die mehrdimensionale Isotopenfraktionierungsanalyse (gleichzeitige Betrachtung von z.B. Kohlenstoff- und Wasserstoffisotopen) Rückschlüsse über die Abbaewege zu ziehen (Elsner et al., 2007; Fischer et al., 2008; Fischer et al., 2007).

Das Ausmaß der Isotopenfraktionierung wird hauptsächlich durch den Reaktionsmechanismus des initialen Angriffs auf das Substrat bestimmt (primärer Isotopeneffekt), wie beispielsweise Meckenstock et al. (1999) für den Toluolabbau gezeigt haben. Sekundäre Isotopeneffekte sind in der Regel ein bis zwei Größenordnungen geringer und beschreiben die Fraktionierung von Isotopen, die nicht direkt in der Abbaureaktion involviert sind.

Ein weiterer Einflussfaktor ist die Molekülgröße. Ein signifikanter Isotopeneffekt findet nur an der Position im Molekül statt, wo die biochemische Reaktion erfolgt. Bei der Analyse der Isotopenverhältnisse einer Verbindung werden allerdings alle Atome eines Moleküls erfasst, weshalb das Ausmaß der Isotopenfraktionierung in Abhängigkeit von der Molekülgröße abgeschwächt wird. Je größer das Molekül ist, umso kleiner ist der nachweisbare Isotopeneffekt. Für Schadstoffe mit großen Molekülen ($> 12\text{--}14$ Kohlenstoffatome), wie zum Beispiel höher molekulare PAKs, ist die Methode der Isotopenfraktionierung daher weniger geeignet, da der messbare Isotopeneffekt zu gering ist, um sensitiv genug nachgewiesen werden zu können (Meckenstock et al., 2004a).

In diesem Zusammenhang unterscheidet man zwischen dem intrinsischen Isotopeneffekt, der die tatsächliche Reaktion im Molekül widerspiegelt und dem kumulativen Isotopeneffekt für das gesamte Molekül, der in der Regel geringer ist (Elsner et al., 2005; Meckenstock et al., 2004a).

Aktuelle Studien haben gezeigt, dass die Isotopenfraktionierung während des mikrobiellen Abbaus organischer Schadstoffe neben dem Reaktionsmechanismus auch von anderen Faktoren, wie zum Beispiel der Temperatur (Morasch et al., 2001), den Enzymen (Morasch et al., 2004; Nikolausz et al., 2006), den am Abbau beteiligten Mikroorganismen sowie ratenlimitierenden Prozessen (z.B. Substrataufnahme und -transport in die Zelle) (Cichocka et al., 2008; Nijenhuis et al., 2005) beeinflusst werden können. Dies führt zu Limitationen für die Anwendung von CSIA zur Erfassung der *in situ* Biodegradation kontaminierten Standorte, da die Wahl eines geeigneten Fraktionierungsfaktors erschwert ist. Eine konservative Abschätzung der *in situ* Biodegradation ist dennoch möglich, insofern die Berechnungen mit den größten Fraktionierungsfaktoren durchgeführt werden, um eine Überschätzung des Abbaupotenzials zu vermeiden (Fischer et al., 2008; Rosell et al., 2007a).

1.4.2 Tracerversuche

Mithilfe von Mikrokosmenstudien im Labor kann der Abbau organischer Schadstoffe unter definierten Bedingungen nachgewiesen werden, in dem die Abnahme der

Schadstoffkonzentrationen und der gleichzeitige Verbrauch von Elektronenakzeptoren sowie die Bildung von Metaboliten als Indikatoren für Biodegradation dienen. Insbesondere für den Nachweis der Biodegradation einzelner Substanzen in einem Mix verschiedener Kontaminanten, die häufig im Bereich ehemaliger Altlasten zu finden sind, ist diese Methode jedoch ungeeignet, da verschiedene Elektronendonator und -akzeptorprozesse gleichzeitig ablaufen, was die Berechnung von Elektronenbilanzen erschwert. Besonderes Interesse besteht darin, die vollständige Mineralisierung eines Substrats beziehungsweise die Transformation in die Biomasse sowie Metabolite nachzuweisen. Hierfür eignen sich Tracerversuche mit isotopisch markierten Substraten (^{13}C , ^{14}C), bei denen anhand der Produktion von markiertem CO_2 die vollständige Mineralisierung eines bestimmten Schadstoffs gezeigt oder anhand der Inkorporation des markierten Kohlenstoffs in die Biomasse der biologische Abbau nachgewiesen werden kann.

Die Analyse von $^{14}\text{CO}_2$ ermöglicht einen sehr sensitiven Nachweis der vollständigen Mineralisierung (Bradley et al., 2002; Chapelle et al., 1996; Fent et al., 2003; Scheunert et al., 1987). Daher sind Tracerversuche besonders für Untersuchungen zum Verhalten und Verbleib von Pestiziden (Mordaunt et al., 2005) sowie persistenten umweltrelevanten Stoffen geeignet. Der Einsatz radioaktiver Tracer birgt jedoch Gefahren für Mensch und Umwelt und strenge Sicherheitsvorkehrungen erschweren die Arbeiten im Labor und im Feldeinsatz. Aufgrund der Umweltbedenken gibt es nur relativ wenige Feldanwendungen mit ^{14}C -markierten Tracerverbindungen (Bianchin et al., 2006; Rügge et al., 1999). Alternativ zu den radioaktiven Markierungen werden zunehmend stabile Isotope (z.B. ^{13}C -markierte Substrate, deuterierte Verbindungen) verwendet (Richnow et al., 1999a). Der wesentliche Vorteil stabiler Isotope liegt in der einfachen Handhabung sowie der Anwendbarkeit in offenen Systemen, so dass für *in situ* Untersuchungen, wie die Erfassung des mikrobiellen Schadstoffabbaus kontaminierten Aquifere, alternativ stabile Isotope als Tracer genutzt werden sollten.

In situ Tracerversuche eignen sich zur Charakterisierung des Transport- und Reaktionsverhaltens eines Grundwasserleiters (Thierrin et al., 1995). Zum Einsatz kommen sowohl konservative (z.B. Uranin, Bromid) als auch reaktive Tracer (der zu untersuchende Schadstoff isotopisch markiert). Prinzipiell können zwei Methoden unterschieden werden: (1) so genannte Push- und Pull Tests, bei denen die Tracer zunächst in einer Messstelle eingeleitet und nach einer gewissen Verweildauer aus der gleichen Messstelle extrahiert werden (Istok et al., 1997; Pombo et al., 2002; Schroth et al., 2001) oder (2) Tracerversuche, bei denen die Einspeisung der Tracer an einem Injektionsbrunnen erfolgt und die Ausbreitung entlang des Fließpfades in so genannten Kontrollbrunnen überwacht wird (Fischer et al., 2006). Mithilfe unterschiedlicher Konzepte wie Isotopenfraktionierung, Analyse von Mineralisierungsprodukten und Metaboliten können anschließend quantitative

Einleitung

Aussagen hinsichtlich des Abbaus getroffen werden (Fischer et al., 2006; Reusser and Field, 2002). Basierend auf dem Einsatz stabiler Isotope können anhand der Inkorporation markierten Kohlenstoffs in die Zellbestandteile der bakteriellen Biomasse Informationen bezüglich der am Schadstoffabbau beteiligten Mikroorganismen gewonnen (Dumont and Murell, 2005; Evershed et al., 2006; Friedrich, 2006b; Madsen, 2006; Radajewski et al., 2003) (siehe auch Kapitel 1.4.3) sowie anhand der Analyse von charakteristischen Metaboliten Abbauwege rekonstruiert werden (Reusser and Field, 2002).

Aufbauend auf dem Konzept stabile Isotope als Tracer zur Erfassung mikrobieller Aktivität zu nutzen, entwickelten Geyer et al. *in situ* Mikrokosmen, die einen direkten Nachweis der Biodegradation im kontaminierten Aquifer anhand der Inkorporation ^{13}C -markierten Kohlenstoffs in die Biomasse ermöglichen (Geyer et al., 2005).

1.4.3 Molekularbiologische Methoden

Mikroorganismen besitzen genetisch und phylogenetisch die größte Diversität unter den Lebewesen. Bisher sind jedoch nur etwa 5000 von weltweit schätzungsweise einer Milliarde existierenden Mikroorganismen kultivierbar (Amann et al., 1995). Das verdeutlicht, welche Schwierigkeiten mit den klassischen Methoden der Kultivierung verbunden sind (Fig. 1-2). Mithilfe von kultivierungsunabhängigen Methoden wie mikroskopischen Untersuchungen, Durchflusszytometrie (*flow cytometry*), Fluoreszenz-*in-situ*-Hybridisierung (FISH), Polymerasekettenreaktion (PCR), DNA Fingerprinting Methoden (z.B. denaturierende Gradientengelektrophorese, DGGE; DNA-Einzelstrang-Konformations-Polymorphismus-Analyse, *single strand conformation polymorphism analysis*, SSCP) sowie Stabile Isotopenbeprobung (SIP) ist es dennoch möglich Informationen über die Struktur und Bedeutung mikrobieller Gemeinschaften zu gewinnen (Fig. 1-2). Eine Vielzahl aktueller Übersichtsartikel geben eine gute Einführung zu den einzelnen Methoden und vergleichen die Vor- und Nachteile, die sich bei den unterschiedlichen Techniken je nach Fragestellung ergeben (Madsen, 2000; Spiegelman et al., 2005; Wagner et al., 2006; Weiss and Cozzarelli, 2008; Widada et al., 2002).

Molekularbiologische Methoden (Fig. 1-2) bieten die Möglichkeit, die Struktur und Diversität der mikrobiellen Gemeinschaft zu erfassen (Imfeld et al., 2008). Darüber hinaus können Gene spezifischer Enzyme auch als funktionelle Marker benutzt werden (Zhou, 2003), um Informationen bezüglich der Abbauwege bestimmter Schadstoffe zu gewinnen und das *in situ* Abbaupotenzial kontaminiert Systeme zu beurteilen (He et al., 2007; Kuntze et al., 2008; Winderl et al., 2007).




Fig. 1-2: Übersicht kultivierungsabhängiger und -unabhängiger Methoden zur Charakterisierung mikrobieller Gemeinschaften (Weiss and Cozzarelli, 2008).

Die vor allem in den letzten Jahren weiterentwickelte Methode der „Stabilen Isotopenbeprobung“ (SIP) von Nukleinsäuren ermöglicht darüber hinaus nicht nur die Diversität der mikrobiellen Gemeinschaften, sondern auch die metabolischen Fähigkeiten der bisher noch unkultivierten Mikroorganismen aufzuklären. Die Methode des SIP bietet daher den Schlüssel, um die Black Box mikrobieller Diversität zu öffnen (Friedrich, 2006a). Die Methode beruht auf dem Einbau von mit stabilen Isotopen (z.B. ^{13}C) markierten Substraten in die Biomasse der aktiven Mikroorganismen einer Lebensgemeinschaft und deren anschließende Identifizierung über den Nachweis der Markierung in Biomarkermolekülen wie Fettsäuren, Aminosäuren, DNA oder RNA (Fig. 1-3). In zahlreichen Übersichtsartikel werden die Konzepte der verschiedenen Methoden vorgestellt, aktuelle Anwendungen beschrieben und Ausblicke zukünftiger Entwicklungen präsentiert (Dumont and Murell, 2005; Friedrich, 2006a; Friedrich, 2006b; Madsen, 2006; Neufeld et al., 2007; Whiteley et al., 2007).

Organische Schadstoffe bestehen hauptsächlich aus Kohlenstoff und Wasserstoff. Kohlenstoff ist neben Wasserstoff und Sauerstoff eines der wichtigsten Elemente der Biosphäre und wird von allen Lebewesen für den Stoff- und Energiewechsel benötigt. Kohlenstoff besitzt zwei stabile Isotope: ^{12}C - und ^{13}C -Kohlenstoff, die natürlicherweise mit ca. 99 % und 1 % Häufigkeit in der Umwelt vorhanden sind (Tab. 1-1). Die in der Regel eine Größenordnung seltener vorkommenden schweren Isotope eines Elementes (z.B. ^{13}C -Kohlenstoff) sind zur Anwendung als Tracer geeignet.

Stabile Isotope werden als Tracer benutzt, um anhand des Verbleibs des Isotops (z.B. ^{13}C -Kohlenstoff) biologische Prozesse in der Umwelt zu untersuchen. Infolge mikrobieller Prozesse wird der isotopisch markierte Kohlenstoff von den Bakterien in ihre Biomasse

Einleitung

inkorporiert. Diese Anreicherung des ^{13}C -markierten Kohlenstoffs in den Biomarkermolekülen der Bakterien dient als eindeutiger Indikator für den Nachweis bestimmter biologischer Prozesse, wie zum Beispiel der Biodegradation organischer Schadstoffe.

Die Anfänge der SIP-Technologie gehen ins Jahr 1958 zurück. Zu dieser Zeit haben Meselson und Stahl erste SIP Experimente mit *Escherichia coli* und ^{15}N -markiertem Substrat (NH_4Cl) durchgeführt (Meselson and Stahl, 1958). Da die Nukleinsäuren zu einem großen Teil auch aus Kohlenstoff und Wasserstoff bestehen, wurde schnell erkannt, dass die DNA bzw. RNA ebenfalls mit den schweren Kohlenstoff und Wasserstoffisotopen markiert werden kann (Schildkraut, 1967).

Zu Beginn des 21. Jahrhunderts führten Radajewski et al. die Methode des DNA-SIP ein (Radajewski et al., 2000). Hierbei wurden Mikroorganismen zunächst in Laborexperimenten mit ^{13}C -markierten Substraten inkubiert. Der ^{13}C -Kohlenstoff der Substrate wird infolge der Assimilation in das Erbgut der aktiven Bakterien inkorporiert. Anschließend erfolgt die Extraktion der DNA aus dem Kultursatz. Die Trennung der isotopisch schweren (^{13}C -DNA) von der isotopisch leichten DNA (^{12}C -DNA) erfolgt mithilfe der Dichtegradientenzentrifugation (Fig. 1-3). Anschließend werden die aktiven Mikroorganismen der mikrobiellen Gemeinschaft anhand der ^{13}C -markierten DNA direkt über Klonierung und Sequenzierung sowie mittels molekularbiologischer Fingerprint-Verfahren identifiziert (Friedrich, 2006b). Bereits wenig später publizierten Manefield et al. erste Anwendungen der RNA-SIP Methode (Manefield et al., 2002a; Manefield et al., 2002b).




Fig. 1-3: Konzept der Stabilen Isotopenbeprobung von Nukleinsäuren am Beispiel des RNA-SIP (Friedrich, 2006a)

Die ersten DNA bzw. RNA-SIP Experimente basierten auf Untersuchungen mit C₁-Verbindungen wie ^{13}C -Methanol ($^{13}\text{CH}_3\text{OH}$), -Methan ($^{13}\text{CH}_4$) oder -Kohlendioxid ($^{13}\text{CO}_2$). Wenig später folgten dann Laborexperimente mit Aromaten und (halogenierten) Aliphaten,

bei denen die Mikroorganismen, die diese Verbindungen assimilieren können, identifiziert wurden (Friedrich, 2006b).

Eine Alternative zur stabilen Isotopenbeprobung von Nukleinsäuren stellt die Analyse der Inkorporation der ^{13}C -markierten Kohlenstoffatome in die Biomarkermoleküle wie Fettsäuren oder Aminosäuren dar. Der wesentliche Vorteil liegt hierbei in der höheren Empfindlichkeit, da die Analyse der Isotopenanreicherung direkt an den Aminosäure- bzw. Fettsäureextrakten mittels *Gas Chromatography-Combustion-Isotope Ratio Mass Spectrometry* (GC-C-IRMS) erfolgen kann (Mottram and Evershed, 2003). Mithilfe dieser Methode können bereits geringe Anreicherungen von schweren Isotopen sehr empfindlich analysiert werden. Eine Auf trennung der Biomasse in schwere und leichte Fraktion ist daher nicht erforderlich und die benötigten Mengen isotopisch markierter Substrate können minimiert werden, so dass die experimentellen Bedingungen den natürlichen Umweltbedingungen angemessen sind (Boschker and Middelburg, 2002; Boschker et al., 1998).

Fettsäuren sind wichtige Zellbestandteile, insbesondere der Zellmembranen (Doppellipidlayer). Der Anteil der Zellmembranen an der Trockenmasse einer Bakterienzelle liegt bei 8-15 % (Schlegel, 1992). Es können drei Hauptgruppen von Membranlipiden unterschieden werden: Phospho-, Glyko- und Neutrallipide. Da Phospholipide (PLFA) die lebende Biomasse charakterisieren, eignen sie sich besonders als Biomarker (Green and Scow, 2000; Kaur et al., 2005). Bestimmte Membranfettsäuren können als Biomarker benutzt werden, weil sie charakteristisch für verschiedene Organismengruppen sind (Tab. 1-2). Zum Teil kann eine weitere Unterscheidung innerhalb dieser Gruppen erfolgen. Bei den sulfatreduzierenden oder methanotrophen Mikroorganismen kann theoretisch sogar zwischen verschiedenen Gattungen differenziert werden (Tab. 1-2). Eingeschränkt kann anhand der Konzentration der Fettsäuren mithilfe von Umrechnungsfaktoren die Menge der Gesamtbiomasse beziehungsweise die Zellzahlen einer Probe abgeschätzt werden (Brinch-Iversen and King, 1990; Kaur et al., 2005; Ludvigsen et al., 1997).

Tab. 1-2: Übersicht ausgewählter Biomarker bestimmter Organismen, verändert nach (Boschker and Middelburg, 2002; Green and Scow, 2000; Kaur et al., 2005; Zelles, 1999). Für die Nomenklatur der Fettsäuren siehe Kapitel 2.4.9.

Organismus	Biomarker	Literatur
Bakterien	i15:0, a15:0, 15:0, 16:1 ω 5, i17:0, 17:0, 18:1 ω 7	(Tunlid and White, 1992)
Gram-positiv	verzweigte PLFAs (br17:0, br18:0, i17:0, i16:0, i16:1, 10Me16:0, 10Me17:0) i15:0, a15:0 einfach ungesättigte Fettsäuren <	(Harwood and Russell, 1984; O'leary and Wilkinson, 1988)

Einleitung

	20% (16:1 ω 9, 16:1 ω 7c, 16:1 ω 5, 18:1 ω 7c, 19:1)	
Gram-negativ	Cyclopropan-Fettsäuren (cy17:0, cy19:0) β -Hydroxy-Fettsäuren der Zellwände einfach ungesättigte Fettsäuren > 20% (16:1 ω 9, 16:1 ω 7c, 16:1 ω 5, 18:1 ω 7c, 19:1)	(Guckert et al., 1985; Parker et al., 1982; White, 1994; Wilkinson, 1988)
Sulfatreduzierer	<i>Desulfobacter</i> (10Me16:0, cy18:0(ω 7,8) <i>Desulfovibrio</i> (i17:1 ω 7c, i15:1 ω 7c, i19:1 ω 7c) <i>Desulfobulbus</i> (17:1 ω 6, 15:1)	(Dowling et al., 1986; Edlung et al., 1985; Parker et al., 1982; Parkes and Calder, 1985; Scheuerbrandt and Bloch, 1962; Taylor and Parkes, 1983)
Methanotrophe	Typ I (16:1 ω 8c, 16:1 ω 6c) Typ II (18:1 ω 8c, 18:1 ω 8t, 18:1 ω 6c)	(Borjesson et al., 1998; Bowman et al., 1993; Makula, 1978; Nichols et al., 1987; Nichols et al., 1985; Sundh et al., 1995)
Aktinomyzeten	(10Me17:0, 10Me18:0)	
Pilze	mehrfach ungesättigte Fettsäuren (18:2 ω 6, 18:3 ω 6, 18:3 ω 3c)	(Federle, 1986; Frostegard and Bååth, 1996; Vestal and White, 1989)
Protozoa	20:2 ω 6, 20:3 ω 6, 20:4 ω 6	(White, 1988; White et al., 1983)
Algen	20:5 ω 3, 18:3 ω 3	

Die Anwendungen, basierend auf der Methode des PLFA-SIP, in der Umweltmikrobiologie sind zahlreich, wie in verschiedenen Übersichtsartikeln zusammengefasst wurde (Boschker and Middelburg, 2002; Evershed et al., 2006; Kreuzer-Martin, 2007; Zhang, 2002). Sehr häufig diente die Methode zur Untersuchung methanotropher Bakterien unter aeroben und anaeroben Umweltbedingungen (Blumenberg et al., 2005; Bull et al., 2000; Crossmann et al., 2005), aber auch Untersuchungen zur Aktivität der mikrobiellen Gemeinschaft im Bereich der Rhizosphäre bilden einen weiteren Anwendungsschwerpunkt dieser Methode (Butler et al., 2003; Lu et al., 2004; Treonis et al., 2004).

Der Fokus dieser Arbeit liegt allerdings auf Anwendungen, die im Zusammenhang mit Umweltschadstoffen stehen. Eine große Anzahl von ^{13}C -markierten Substraten ist bei kommerziellen Anbietern erhältlich. Die verhältnismäßig hohen Preise für ^{13}C -markierte Verbindungen limitieren jedoch maßgeblich die Anwendung im Feldmaßstab. Experimente

zum Nachweis des biologischen Abbaus organischer Schadstoffe sind daher noch verhältnismäßig selten, insbesondere unter *in situ* Bedingungen. Die ersten Untersuchungen zum Nachweis des Toluol-Abbaus erfolgten in Batchkulturen sowie in Boden-, Sediment- und Grundwasser-Mikrokosmen (Hanson et al., 1999; Mauclaire et al., 2003; Pelz et al., 2001a; Pelz et al., 2001b). Weitere Studien beschäftigten sich mit dem PAK-Abbau (Johnsen et al., 2002; Richnow et al., 1998). Ein wesentliches Problem für die Anwendung dieser Methode im Feldmaßstab bestand im Zusammenhang mit der Verfügbarkeit ausreichender Mengen ^{13}C -markierter Substrate. Dieses Problem konnte mit der Entwicklung so genannter Bio-Sep® Beads gelöst werden (Peacock et al., 2004; White et al., 2003). Bio-Sep® ist ein aktivkohlehaltiges Material, das einerseits als Substratpool für die ^{13}C -markierten Verbindungen und andererseits als Aufwuchskörper den Mikroorganismen zur Verfügung steht, die anschließend extrahiert werden können, um die Inkorporation des ^{13}C -Kohlenstoffs in den Zellbestandteilen zu analysieren. Erstmals wurden das Konzept der *in situ* Mikrokosmen mit Bio-Sep® Beads am Standort Zeitz zum Nachweis des Benzol und Toluol Abbaus im kontaminierten Grundwasser eingesetzt (Geyer et al., 2005).

Die Fettsäuren der Membranlipide waren die ersten Biomarker, die in Kombination mit stabilen Isotopenreicherungen untersucht wurden (Boschker et al., 1998; Manefield et al., 2004). Der wesentliche Vorteil liegt in der hohen Empfindlichkeit dieser Methode, ohne dass eine Auftrennung von isotopisch leichter und schwerer Fraktion nötig wird. Daher können selbst geringe Aktivitäten detektiert werden, so dass die Mengen des bereit gestellten ^{13}C -markierten Substrats den natürlich vorkommenden Substratkonzentrationen entsprechen können. Das ist hinsichtlich Reproduzierbarkeit der natürlichen Bedingungen von Bedeutung und nicht zuletzt auch ein finanzieller Vorteil, da folglich bereits geringe Mengen der teuren ^{13}C -markierten Substrate genügen, um eine Anreicherung in den Membranlipiden nachweisen zu können. Aufgrund der extrem hohen Sensitivität ist diese Methode insbesondere für *in situ* Experimente geeignet, bei denen in der Regel nur relativ geringe Mengen an Biomasse extrahiert werden können und auch die ^{13}C -Anreicherung in den Biomarkern meist relativ gering ist.

Fettsäuren, insbesondere die PLFA, bieten zwar ein gewisses taxonomisches Potenzial (Green and Scow, 2000; Kaur et al., 2005; Zelles, 1999), aber Variationen der Gen-Sequenzen verschiedener Mikroorganismen sind deutlich höher und erlauben detailliertere Aussagen hinsichtlich den Veränderungen in der aktiven mikrobiellen Gemeinschaft. Im Gegensatz zu PLFA-SIP ist es mithilfe von DNA-SIP möglich die *in situ* Funktion verschiedener physiologischer Gruppen der mikrobiellen Gemeinschaft anhand der Kohlenstoffflüsse zu bestimmen und damit eine Charakterisierung komplexer Zusammenhänge zwischen Umweltfaktoren und physiologischer Diversität vorzunehmen (Lu et al., 2007). Allerdings sind für die Anwendung von DNA-SIP deutlich höhere ^{13}C -

Einleitung

Anreicherungen von > 30 % notwendig (Radajewski et al., 2003) und es bedarf mindestens zwei Zellteilungen, bis die neu synthetisierte DNA vollständig ^{13}C -markiert ist, was für eine optimale Trennung von schwerer und leichter DNA im Dichtegradienten wichtig ist (Manefield et al., 2002a; Manefield et al., 2002b).

Der Vorteil des RNA-SIP gegenüber dem DNA-SIP liegt in der größeren Sensitivität, da jedes neu synthetisierte RNA-Molekül nach Zugabe ^{13}C -markierter Substrate vollständig markiert sein kann, wodurch die RNA deutlich schneller ^{13}C -markiert wird als die DNA ohne dass es eine Zellteilung erfordert (Manefield et al., 2002a; Radajewski et al., 2003).

DNA und RNA haben ein deutlich größeres taxonomisches Potenzial als die Fettsäuren, jedoch sind relativ große Mengen Biomasse erforderlich und eine ^{13}C -Inkorporation von mindestens 30 % wird benötigt. Die resultierenden langen Inkubationszeiten können zusätzlich zum Problem des „cross-feeding“ innerhalb einer Nahrungskette führen. Außerdem werden größere Mengen der ^{13}C -markierten Substrate für die erfolgreiche Durchführung eines DNA- bzw. RNA-SIP Experimentes benötigt (Radajewski et al., 2003).

1.5 *In situ* Mikrokosmen

Die Methode des SIP ist geeignet, um Aktivität, Diversität und Funktionalität mikrobieller Gemeinschaften unter den verschiedensten Umweltbedingungen zu analysieren. Da die Laborergebnisse nicht ohne weiteres auf die weitaus komplexeren natürlichen Systeme übertragen werden können, besteht seit längerem das Bestreben die verschiedenen Untersuchungen möglichst unter *in situ* Bedingungen durchzuführen (Mandelbaum et al., 1997). Trotzdem gibt es bis heute nur wenige Testsysteme, die Experimente direkt unter Aquiferbedingungen ermöglichen. In einem Übersichtsartikel zu *in situ* Mikrokosmen beschreiben Mandelbaum et al. (1997) zwei verschiedene *in situ* Mikrokosmen Konzepte, um einerseits in isolierten, ungestörten Bereichen des Aquifers Aussagen bezüglich physikalisch-biochemischer Parameter zu erhalten beziehungsweise um Rein- und Mischkulturen aus dem Labor direkt im Grundwasserstrom zu testen (Mandelbaum et al., 1997). In einer Studie von Hendrickx et al. (2005) wurde unkontaminiertes Aquifermaterial in einem Mesokosmos sowohl im kontaminierten als auch im unkontaminierten Bereich des Aquifers inkubiert, um *in situ* die Dynamik der sich entwickelnden mikrobiellen Gemeinschaft zu analysieren.

Um die Methode des SIP direkt im kontaminierten Aquifer zum Nachweis der Biodegradation organischer Schadstoffe anwenden zu können, wird ein Testsystem benötigt, dass einen Substratpool für die isotopisch markierten Substanzen (^{13}C -markierte Kontaminanten) zur Verfügung stellt und als Aufwuchskörper für die Grundwassermikroorganismen dient. Ziel hierbei ist es, entgegen aufwändiger und kostspieliger Tracerexperimente wie zum Beispiel Push-Pull Tests, mit weitaus geringeren Mengen ^{13}C -markierter Substanzen den direkten

Nachweis der *in situ* Biodegradation zu führen und damit perspektivisch eine effiziente Methode für die Anwendung in der Altlastenpraxis zu etablieren.

White et al. (2003) entwickelten zunächst zur Überwachung der Trinkwasserqualität einen Biofilm Sammler, bestehend aus Bio-Sep® Beads. Bei Bio-Sep® (K. L. Sublette, Tulsa, USA) handelt es sich um ein aktivkohlehaltiges Material mit besonders großer innerer Oberfläche und Poren von ca. 2-20 µm Durchmesser, was den Bakterien die Besiedlung des Kugellinneren ermöglicht. In weiteren Anwendungen wurden die Biofilm Sammler genutzt, um Veränderungen innerhalb der mikrobiellen Gemeinschaft des Grundwassers während der Zugabe verschiedener Elektronendonatoren in Push-Pull Tests zu beobachten (Istok et al., 2004; Peacock et al., 2004). Hierfür wurden die Testsysteme für mehrere Wochen in den Grundwassermessstellen inkubiert. Während dieser Zeit besiedelten die Mikroorganismen das Innere der Aufwuchskörper und bildeten Biofilme. Die spätere Extraktion der Biomasse und Analyse der Biomarker (DNA, Lipide) zeigte, dass signifikante Veränderungen innerhalb der mikrobiellen Gemeinschaft infolge der Stimulation mit verschiedenen Elektronendonatoren mithilfe dieser Art der Probenahme detektiert werden konnten.




Fig. 1-4: Konzept der *in situ* Mikrokosmen (BACTRAP®) basierend auf ¹³C-markierten Substraten zum Nachweis mikrobieller Aktivität anhand der Inkorporation der isotopischen Markierung in die Biomarkermoleküle.

In einem ersten Versuch zum Nachweis der *in situ* Biodegradation von organischen Grundwasserkontaminanten wurden Bio-Sep® Bead Sammler (BACTRAP®) zunächst im Labor mit ¹³C-markierten Schadstoffen (¹³C-Benzol oder ¹³C-Toluol) beladen und anschließend in einem BTEX kontaminierten Grundwasserleiter inkubiert (Geyer et al.,

Einleitung

2005). Der Nachweis ^{13}C -markierter Fettsäuren sowohl auf den Toluol- als auch Benzol beaufschlagten BACTRAP®'s diente als direkter Indikator für den anaeroben Abbau von Benzol und Toluol unter *in situ* Bedingungen. Damit wurde erstmalig ein Testsystem beschrieben, welches den Nachweis der *in situ* Biodegradation basierend auf der SIP-Technik direkt im kontaminierten Grundwasserleiter ermöglichte (Fig. 1-4).

1.6 Zielstellung und Inhalt der Arbeit

Die Zielsetzung für diese Arbeit war, das Testsystem der *in situ* Mikrokosmen (BACTRAP®) weiterzuentwickeln, um die Methode in der Altlastenpraxis anwenden und etablieren zu können. Hierfür war es erforderlich die Methode durch den Vergleich mit anderen Techniken zu evaluieren sowie in weiteren Feldanwendungen und mit verschiedenen Schadstoffen zu testen. Diese Arbeit beinhaltet daher Ergebnisse von mehreren Feldversuchen, die an verschiedenen Standorten und mit unterschiedlichen Schadstoffen durchgeführt wurden. Den Schwerpunkt bilden dabei Untersuchungen mit BTEX (insbesondere Benzol, Toluol) sowie Monochlorbenzol (MCB). Obwohl der anaerobe Benzolabbau unter nitrat- (Coates et al., 2001; Ulrich and Edwards, 2003), eisen (III)- (Anderson et al., 1998; Cozzarelli et al., 1990; Lovley, 2000; Lovley et al., 1989; Lovley et al., 1994; Lovley et al., 1996) und sulfatreduzierenden (Anderson and Lovley, 2000; Lovley, 1995; Lovley, 2000; Wiedemeier et al., 1999) sowie methanogenen (Bekins et al., 2001; Grbic-Galic and Vogel, 1987) Bedingungen bereits seit mehreren Jahren bekannt ist, konnten die am Abbau beteiligten Mikroorganismen bisher nicht isoliert werden. MCB galt sogar bis vor kurzem unter anoxischen Bedingungen als persistente, nicht abbaubare Verbindung. Ergebnisse dieser Dissertation (Kapitel 5 bis 7) lieferten den Nachweis, dass unter anaeroben Bedingungen der Abbau von MCB erfolgen kann. MCB wurde hierbei als Modellverbindung für sehr persistente Substrate betrachtet.

Die Arbeitsziele dieser Promotionsarbeit lauteten:

- Validierung des Testsystems zur Beurteilung der *in situ* Biodegradation kontaminiertes Aquifere mithilfe alternativer Methoden
- Anwendung der *in situ* Mikrosomen an verschiedenen Feldstandorten und für unterschiedliche Schadstoffe
- Weiterentwicklung des Testsystems zur Analyse mikrobieller Konsortien und Rekonstruktion von Abbauwegen
- Charakterisierung des intrinsischen Abbaupotenzials persistenter Schadstoffe an komplexen Standorten

- Implementierung in die Altlastenpraxis als Nachweismethode zur Erfassung der *in situ* Biodegradation im Kontext von *Natural Attenuation* als Standortsicherungsmaßnahme

Die vorliegende Arbeit ist eine kumulativ verfasste Dissertationsschrift, deren Hauptteil sich aus sechs Veröffentlichungen zusammensetzt. Davon wurden drei Artikel als Erstautor verfasst, von denen bereits zwei im Jahr 2006 in den Fachzeitschriften *Grundwasser* und *Organic Geochemistry* veröffentlicht werden konnten. Der dritte Artikel wurde 2008 zur Veröffentlichung bei *Environmental Science and Technology* eingereicht. Bei den anderen Publikationen hat die Doktorandin als Co-Autorin maßgeblich mitgewirkt und war sowohl bei Feldversuchen und Laborarbeiten, als auch bei der Erstellung der Manuskripte wesentlich beteiligt. Die Labor- und Feldarbeiten wurden teilweise von Diplomanden und Praktikanten unterstützt, wobei die Autorin aktiv und verantwortlich bei der Betreuung der Arbeiten mitgewirkt hat.

Die Dissertation beginnt nach einer allgemeinen, übergreifenden Einleitung (Kapitel 1) im Hauptteil mit einer Gegenüberstellung geochemischer und isotopenchemischer Methoden zur Charakterisierung des anaeroben Benzolabbaus im Rahmen eines Multilevel-Experimentes (Kapitel 2). Das Kapitel 2, 2006 in der Fachzeitschrift *Grundwasser* (Band 11(4), S. 247-258) veröffentlicht, dient des Weiteren als umfangreiche Methodenbeschreibung zur Herstellung, Installation und Extraktion der *in situ* Mikrokosmen. Der Artikel wurde in einer deutschsprachigen Fachzeitschrift veröffentlicht, um auf diesem Weg möglichst eine Vielzahl der deutschen Sanierungspflichtigen, Behörden und Ingenieurbüros aus dem angewandten Bereich zu erreichen.

Kapitel 3, 2006 in *Organic Geochemistry* (Band 37(10), S. 1394-1410) publiziert, präsentiert weitere Beispiele zum Nachweis der *in situ* Biodegradation von Benzol und Toluol im hoch- und niedrig kontaminierten Grundwasser an zwei verschiedenen Feldstandorten. Außerdem wird der Frage nachgegangen, ob *in situ* Mikrokosmen geeignet sind, Abbauwege anhand von Metaboliten, die sich auf dem Testsystem anreichern, rekonstruieren und die mikrobielle Gemeinschaft des Grundwassers charakterisieren zu können. Hierfür wurde alternativ zur Analyse der Fettsäureprofile die Methode des *Single Strand Conformation Polymorphism* (SSCP) gewählt. Die verschiedenen Untersuchungsergebnisse (Geochemie, *in situ* Mikrokosmen, Molekularbiologie) dienten einer Verifizierung der Methode der BACTRAP®s. Kapitel 4, 2006 als Übersichtsartikel in *Engineering Life Science* (6(3), S. 234-251) veröffentlicht, dokumentiert den gegenwärtigen Stand des Wissens über Verfahren zur Beurteilung der *in situ* Biodegradation im kontaminierten Grundwasser. Es werden die aktuellen Ergebnisse und Fortschritte der Methode der Stabilen Isotopenfraktionierungsanalyse (SIFA) sowie der *in situ* Mikrokosmen vorgestellt und diskutiert. Außerdem wird der Frage nachgegangen, ob die mikrobielle Gemeinschaft, die

Einleitung

das *in situ* Testsystem besiedelt, tatsächlich die Mikroflora des Aquifers repräsentiert, in dem die Biomasse des Grundwassers und des Testsystems mittels SSCP verglichen wurden. Ziel dieser Betrachtung ist es zu prüfen, ob die Ergebnisse der *in situ* Mikrokosmen auf den Grundwasserleiter übertragbar sind, eine wichtige Voraussetzung für die Erfassung des *in situ* Abbaupotenzials an kontaminierten Standorten.

Einen Schwerpunkt dieser Dissertation stellen Untersuchungen zum Nachweis des anaeroben Abbaus von Monochlorbenzol dar. Die diesbezüglichen Ergebnisse sind in den Kapitel 4, 5, 6 und 7 aufgeführt. Kapitel 5, 2007 in *Environmental Science and Technology* (41, S. 3836-3842) publiziert, befasst sich im Wesentlichen mit dem sensitiven Nachweis des anaeroben MCB Abbaus in Tracerversuchen mit ¹³C-markiertem MCB in *in situ* und *ex situ* Mikrokosmen.

Kapitel 6, 2008 zur Veröffentlichung bei *Environmental Science and Technology* eingereicht, zeigt die Entwicklung eines integrativen Untersuchungskonzepts für die Beurteilung des Natural Attenuation Potenzials komplexer Standorte, die durch multiple, schwer abbaubare Schadstoffe wie chlorierte Benzole, sequentielle Abbaupfade und heterogene Aquiferbedingungen charakterisiert sind. Um plausible Aussagen bezüglich der biogeochemischen Prozesse eines Untersuchungsstandortes zu erhalten, müssen neuartige Untersuchungsmethoden wie CSIA und BACTRAP® sowie statistische Auswertungsmethoden miteinander kombiniert werden.

Mit der Frage nach weiteren Anwendungsfeldern für den Einsatz von *in situ* Mikrokosmen beschäftigt sich Kapitel 7, dass 2007 in *Environmental Pollution* (148, S. 428-437) veröffentlicht wurde. Es handelt sich ebenfalls um eine Studie zum MCB Abbau, allerdings in einer Pflanzenkläranlage. Zielsetzung war es die Abbauleistung der Pflanzenkläranlage für MCB zu evaluieren. Erstmalig wurden hierfür *in situ* Mikrokosmen direkt im Sediment inkubiert.

Der Hauptteil der Dissertation endet mit einer kritischen Zusammenfassung und Diskussion aller Ergebnisse in Kapitel 8.

2 Analyse des anaeroben Benzolabbaus:

Vergleich von *In situ* Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen

Nicole Stelzer, Anko Fischer, Matthias Kästner, Hans-H. Richnow

(Dieses Kapitel wurde in der Zeitschrift Grundwasser Band 11(4), S. 247-258 veröffentlicht)

2.1 Kurzfassung

Geochemische und isotopenchemische Methoden wurden zur Charakterisierung des anaeroben Benzolabbaus am Standort eines ehemaligen Hydrierwerkes (Zeitz, Deutschland) genutzt. Im Abstrom des Schadenszentrums wurde hierfür die vertikale Struktur der Benzolfahne in verschiedenen Tiefen untersucht. Durch Einsatz von [$^{13}\text{C}_6$] Benzol in *in situ* Mikrokosmen konnte anhand der Transformation des markierten Kohlenstoffs in die Biomasse eindeutig das Abbaupotenzial nachgewiesen werden. Das Fettsäurespektrum sowie deren Isotopensignaturen deuten auf eine Besiedlung durch komplexe mikrobielle Gemeinschaften hin, die in unterschiedlicher Weise am Benzolabbau beteiligt sind. Die Sulfatsenke in der Schadstofffahne deutet auf überwiegend sulfatreduzierende Abbaubedingungen mit einem Abbaupotenzial von etwa 1,7 mmol L $^{-1}$ Benzol hin. Anhand der Isotopensignaturen und Konzentrationen des DIC wurde in der Schadstofffahne ein Abbau von 2,2 mmol L $^{-1}$ Benzol abgeschätzt. Eine Berechnung des Benzolabbaus anhand der Isotopenfraktionierungsmethode ergibt einen Abbau von etwa 3,0 mmol L $^{-1}$ auf dem Fließweg. Die quantitativen geochemischen und isotopenchemischen Abschätzungen liegen in der gleichen Größenordnung und zeigen einen signifikanten Benzolabbau im oberen Aquifer.

2.2 Abstract

Geochemical and isotope chemical methods were applied to assess the *in situ* biodegradation of benzene in a shallow aquifer (Zeitz, Germany). The vertical structure of the plume was investigated in a multi level sampling approach. Benzene degradation was investigated using *in situ* microcosms incubated with [$^{13}\text{C}_6$]-labelled benzene. The transformation of ^{13}C into fatty acids proved biodegradation and indicated that a complex microbial community is colonizing the *in situ* microcosms.

The sink of sulphate in the course of the plume indicated sulphate reducing conditions which may account for the oxidation of 1.7 mmol L $^{-1}$ benzene. The concentration and isotope composition of dissolved inorganic carbon indicated a degradation of 2.2 mmol L $^{-1}$ benzene within the plume. The isotope composition of benzene suggested an average degradation of

3.0 mmol L⁻¹ benzene on the groundwater flow path. The various approaches to assess the benzene degradation were in the same order and illustrated significant benzene degradation in the anaerobic aquifer.

2.3 Einleitung

Benzol, Toluol, Ethylbenzol und Xylole (BTEX) sind häufige Grundwasserkontaminanten, die eine Gefahr für Ökosysteme und die menschliche Gesundheit darstellen. In einem verunreinigten Aquifer wird der Verbleib dieser Verbindungen durch abiotische (Advektion, Dispersion, Verflüchtigung, Sorption) und biologische Prozesse beeinflusst. Eine nachhaltige Reduzierung der Schadstoffmenge erfolgt allerdings hauptsächlich durch den mikrobiellen Abbau. Bei der Anwendung von Sanierungsstrategien wie *Monitored Natural Attenuation* (MNA) und *Enhanced Natural Attenuation* (ENA) kommt somit dem mikrobiellen Schadstoffabbau eine besondere Bedeutung zu. Es werden daher geeignete Methoden benötigt, um das natürliche Selbstreinigungspotenzial eines kontaminierten Grundwasserleiters bestimmen zu können.

Eine Möglichkeit zur Untersuchung des *in situ* Abbaus von Schadstoffen ist die Nutzung der substratspezifischen Isotopenfraktionierung (Meckenstock et al., 2004a). Diese Methode beruht darauf, dass Bakterien infolge des mikrobiellen Schadstoffabbaus das natürliche Isotopenverhältnis (z.B. ¹³C/¹²C) eines Substrates verändern. Die bevorzugte Verwertung der leichten Isotopomere einer Kontaminante führt infolge der mikrobiellen Umsetzung zu einer Anreicherung der schweren Isotopomere in der residualen Schadstofffraktion, was mit Hilfe der Gaschromatographie-Isotopenverhältnis-Massenspektroskopie (GC-IRMS) analytisch nachgewiesen werden kann.

Zur Abschätzung der Biodegradation in kontaminierten Grundwasserleitern werden verschiedene Verfahren angewendet, die häufig auf der Kultivierung von Mikroorganismen im Labor beruhen. Der anaerobe Schadstoffabbau ist in der Regel ein sehr langsam ablaufender Prozess und viele der am Abbau beteiligten Mikroorganismen sind bisher noch nicht oder nur eingeschränkt kultivierbar. Labortestverfahren erscheinen daher nur begrenzt geeignet zu sein, um die realen, sehr komplexen Vorgänge und anoxischen Milieubedingungen des fließenden Grundwassers abzubilden. Darüber hinaus sind anaerobe Abbauversuche sehr langwierig und generell mit einem relativ großen Zeitaufwand verbunden.

Hieraus resultiert ein großer Bedarf an der Entwicklung geeigneter *in situ* Methoden zur Bestimmung der im Grundwasserleiter stattfindenden Biodegradation, insbesondere für anoxische Milieubedingungen.

Der mikrobielle Abbau aller BTEX-Verbindungen konnte unter oxischen und anoxischen Bedingungen nachgewiesen werden. Aufgrund der Sauerstofflimitation findet in

kontaminierten Grundwasserleitern meist eine anaerobe Biodegradation der monoaromatischen Kohlenwasserstoffe statt (Cerniglia, 1984; Chakraborty and Coates, 2004; Gibson and Subramanian, 1984; Wiedemeier et al., 1999). Dabei ist Benzol im Vergleich zu Toluol, Ethylbenzol und Xylolen oft schwerer abbaubar, so dass folglich bis heute nur wenige anaerob benzolabbauende Konsortien und eine Reinkultur isoliert wurden (Chakraborty and Coates, 2004; Coates et al., 2001; Wiedemeier et al., 1999). Das begründet, warum in Laborstudien der anaerobe Benzolabbau bisher nur relativ selten eindeutig nachgewiesen werden konnte und sich in Kulturversuchen nur schwer reproduzieren lässt.

Tracerexperimente mit isotopisch markierten Substanzen stellen eine alternative Methode zur Untersuchung der Biodegradation im Grundwasser dar. Radioaktiv markierte Substanzen (^{14}C) sollten aufgrund ökologischer Risiken nicht in offenen Systemen eingesetzt werden. Ersatzweise dienen stabile Isotope (D, ^{13}C) als Tracer. Während der Assimilation wird der isotopisch markierte Kohlenstoff von den Mikroorganismen zur Synthese von Biomasse verwendet. Durch den Einsatz von ^{13}C -markierten Substraten kann der mikrobielle Schadstoffabbau anhand der Anreicherung des ^{13}C -markierten Kohlenstoffs in verschiedenen Zellkomponenten nachgewiesen werden. Dabei stehen als Zellmembranbestandteile häufig Phospholipid-Fettsäuren (**phospholipid-derived fatty acids**, PLFA) im Mittelpunkt der Untersuchungen (Abraham et al., 1998; Boschker and Middelburg, 2002; Taylor and Parkes, 1983). Es sind nur wenige Arbeiten zur Nutzung ^{13}C -markierter Substrate für die Bestimmung des mikrobiellen Schadstoffabbaus anhand der ^{13}C -Anreicherung in PLFA bekannt. Die Anreicherung von ^{13}C in verschiedenen PLFA diente zum Nachweis des Toluolabbaus. Dabei wurde Aquifermaterial in Labormikrokosmen mit [^{13}C]- α -Toluol als Substrat inkubiert, die Transformation des Isotopensignals in die Biomasse nachgewiesen, und die am anaeroben Toluolabbau beteiligten Mikroorganismen identifiziert (Pelz et al., 2001a; Pelz et al., 2001b). In ähnlichen Studien mit [$^{13}\text{C}_6$]-markiertem Toluol konnten anhand des Einbaus von ^{13}C in PLFA die mikrobiellen Populationen in aeroben Bodenmikrokosmen identifiziert werden (Hanson et al., 1999). In einem "Push & Pull"-Tracerexperiment mit [2- ^{13}C]-Acetat als Substrat und Nitrat als Elektronenakzeptor ist anhand der Transformation der ^{13}C -Markierung in die verschiedenen PLFA die mikrobielle Abbauaktivität im Aquifer nachgewiesen worden (Pombo et al., 2002). Demgegenüber wurden spezielle *in situ* Mikrokosmen (BACTRAP[®]) entwickelt, die mit ^{13}C -markierten Testsubstanzen beladen und direkt *in situ* im Aquifer inkubiert werden können. Anhand der Transformation der isotopischen Markierung in die Biomasse (Membranfettsäuren) kann der Schadstoffabbau nachgewiesen werden. Zur Erprobung dieses Testsystems wurden verschiedene Experimente durchgeführt (Büning et al., 2005; Geyer et al., 2005).

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Die vorliegende Arbeit stellt die Ergebnisse eines Feldversuches vor, bei dem BACTRAPS zur Analyse des Benzolabbaus genutzt wurden. Dabei sind die Mikrokosmen mit [¹³C₆] Benzol als Substrat beladen und mit Hilfe eines Multilevel-Packersystems (MLPS) in acht verschiedenen Tiefen einer Grundwassermessstelle eines BTEX-kontaminierten Aquifers während 51 Tagen inkubiert worden. Der Einbau der ¹³C-Atome aus dem Benzol in mikrobielle Fettsäuren wurde im Anschluss mittels GC-IRMS nachgewiesen und als Beweis für den *in situ* Benzolabbau genutzt. Hydrogeo- und isotopenchemische Parameter dienten zur Charakterisierung des Abbaumilieus. Darüber hinaus wurde der *in situ* Benzolabbau mit Hilfe der Isotopenfraktionierungsmethode und anhand von Elektronenbilanzen abgeschätzt. Die Diskussion der Ergebnisse erfolgte im Hinblick auf den Einsatz verschiedener Methoden zur Kennzeichnung des *Natural Attenuation* Potenzials.

2.4 Material und Methoden

2.4.1 Isotopen markierte Substanzen und Chemikalien

Es wurden ausschließlich Chemikalien in analytischer Qualität (p. A.) verwendet. Isotopisch markiertes [¹³C₆] Benzol wurde von Sigma-Aldrich (St. Louis, USA) mit einer chemischen Reinheit von 99 % bezogen.

2.4.2 Standortbeschreibung

Bei dem Untersuchungsstandort handelt es sich um einen BTEX-Schadensfall im Bereich eines ehemaligen Hydrierwerkes bei Zeitz (Sachsen-Anhalt, Deutschland). Am Standort sind im Wesentlichen zwei Grundwasserleiter ausgebildet, die durch einen eozänen Braunkohle-führenden Ton-Schluff-Komplex voneinander getrennt sind, jedoch lokal teilweise infolge Erosion des Stauerkomplexes in direktem hydraulischem Kontakt stehen. Die Geschichte, Hydrogeologie und Geochemie des Standortes wurden ausführlich beschrieben (Dahmke et al., 2004; Fischer et al., 2004; Gödeke et al., 2004a; Gödeke et al., 2004b; Schirmer et al., 2006; Vieth et al., 2001; Vieth et al., 2005). Benzol stellt mit einem Anteil von mehr als 99 % die Hauptkontaminante dar, wobei im Zentrum der Schadstofffahne Konzentrationen von > 1000 mg L⁻¹ bestimmt wurden. Der mikrobielle BTEX-Abbau ist überwiegend durch anaerobe Prozesse wie Sulfatreduktion geprägt (Fischer et al., 2004; Vieth et al., 2005).

Die hier beschriebenen Untersuchungen wurden im oberen, ca. 2-10 m mächtigen jungtertiären und quartären Aquifer (GWL I) durchgeführt, der hauptsächlich aus Terrassenschottern der Weißen Elster besteht. Die vertikale Struktur der Kontaminationsfahne wurde im Bereich der Grundwassermessstelle Saf ZZ 23/00, die sich im nördlichen Abstrom des Kontaminationsherdes befindet, untersucht (Fig. 2-1).




Fig. 2-1: Verteilung der BTEX-Konzentrationen ($\mu\text{g L}^{-1}$) im oberen Aquifer im Bereich des ehemaligen Hydrierwerkes bei Zeitz (Sachsen-Anhalt).

2.4.3 Tiefenorientierte Grundwasserbeprobung

Die tiefenorientierte Grundwasserbeprobung wurde mit Hilfe eines Multilevel-Packersystems (MLPS) durchgeführt (Schirmer et al., 1995). Die Förderung des Grundwassers erfolgte dabei mit pneumatisch regulierbaren Miniatur-Doppelventil-Pumpen (Innovative Messtechnik Weiß), die eine gleichmäßige und blasenfreie Probennahme gewährleisteten. Bei geringer Förderleistung (ca. 100 ml min^{-1}) sowie einem Abstand zwischen den einzelnen Pumpen von etwa einem Meter werden bei der Entnahme der Grundwasserproben vertikale hydrogeochemischen Gradienten nicht durch Vermischungseffekte beeinflusst. Für die Erzeugung des benötigten Pumpendruckes wurde Stickstoff aus einer Druckflasche benutzt, um Sauerstoffeintrag während der Probenahme auszuschließen. Mit Ausnahme des Horizontes in 9,40 m Tiefe konnte aus allen verbleibenden 7 Tiefen das benötigte Probenvolumen für die Analysen entnommen werden (Fig. 2-2). Die Grundwasserproben wurden bis zur weiteren Analyse bei 4°C kühl und dunkel gelagert.

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen




Fig. 2-2: Links, Schichtenprofil und Ausbauplan der Grundwassermessstelle Saf ZZ 23/00. Tiefe 1 bis 8 bezeichnet die Position der Miniatur-Doppelventilpumpen und BACTRAPs. Rechts, Foto des MLPS mit Edelstahlkäfigen für BACTRAPs vor dem Einbau in die Messstelle.

2.4.4 Analytik der GW-Proben

Die Benzolkonzentrationen sind gaschromatographisch nach DIN 38407-F9-1 bestimmt worden. Die ionenchromatographische Analyse der Sulfat- und Nitrat-Konzentrationen basiert auf DIN EN ISO 10304-1 D19. Die Bestimmung der Konzentration des gelösten anorganischen Kohlenstoffs (DIC) erfolgte nach DEV D8, der Eisen(II)-Konzentrationen nach DIN 38406-E1-1 und der Eisen(III)- sowie Mangan-Konzentrationen nach DIN EN ISO 11885-E22.

Zur Bestimmung der Kohlenstoffisotopensignaturen des Benzols ($\delta^{13}\text{C}_{\text{Benzol}}$), des Methans ($\delta^{13}\text{C}_{\text{CH}_4}$) und des DIC ($\delta^{13}\text{C}_{\text{DIC}}$) sind erprobte Methoden verwendet worden (Fischer et al., 2004; Vieth et al., 2005).

2.4.5 Herstellung der BACTRAPs

Bei der Herstellung der BACTRAPs dienten Bio-Sep®-Kugeln (K. L. Sublette; Universität Tulsa, USA) als Trägermaterial für die Testsubstanzen [^{12}C]- und [$^{13}\text{C}_6$] Benzol und als Aufwuchskörper für die Mikroorganismen. Dabei handelt es sich um Kugeln mit 2-3 Millimeter Durchmesser, bestehend aus 75 % pulverisierter Aktivkohle eingebettet in einer Matrix aus thermostabilem Nomex®. Die Oberfläche der Kugeln weist Poren mit einem

Durchmesser von 2-10 µm auf. Bei einer Dichte von ca. 0,16 g cm⁻³ zeichnen sich die Kugeln besonders durch ihre hohe innere Oberfläche von > 600 m² g⁻¹ Bio-Sep®-Material aus (White et al., 2003). Die Kugeln wurden zur Entfernung residualer organischer Kohlenstoffverbindungen bei 300 °C im Muffelofen für ca. 3 Stunden ausgeheizt.

Für die Herstellung der BACTRAPS ist ein Teflonrohr (VWR Darmstadt, Deutschland) mit 10 Millimeter Durchmesser in jeweils 40 Millimeter lange Stücke zerschnitten und anschließend perforiert worden, um einen Grundwasserdurchfluss gewährleisten zu können (Fig. 2-3). Pro BACTRAP wurden jeweils 0,4 g Bio-Sep® genutzt. Ein Glaswollestopfen am oberen und unteren Ende verhinderte das Herausfallen des Bio-Sep®-Materials.




Fig. 2-3: Schematischer Aufbau eines BACTRAP (links) und Foto der Befestigung des Edelstahlkäfiges mit BACTRAP an der Packermembran (rechts).

2.4.6 Beladung der BACTRAPS

Die BACTRAPS wurden im Autoklav sterilisiert und hydratisiert. Im Anschluss erfolgte die Beladung der Mikrokosmen mit [¹²C]- bzw. [¹³C₆] Benzol. BACTRAPS ohne Benzolaufschlag dienten als Kontrolle zur Analyse der mikrobiellen Besiedlung ohne zusätzliches Substrat.

Die BACTRAPS wurden in verschließbare Glasgefäße überführt und jeweils mit etwa 76 mg Benzol pro Gramm Bio-Sep® bei einem Unterdruck von ca. 60 mbar beladen. Um einen Gleichgewichtszustand in der Gasphase herzustellen und eine einheitliche Beladung der Aktivkohle mit dem Benzol gewährleisten zu können, verblieben die BACTRAPS zur Inkubation bei Raumtemperatur für mindestens 48 Stunden in dem verschlossenen Gefäß. Die Beladung ist mit einer Standardabweichung von ± 5 % reproduzierbar (Geyer et al., 2005). Die BACTRAPS wurden im Anschluss in einer Anaerobbox mit sterilem, anaeroben Wasser überschichtet, kurz evakuiert und dann unter anoxischer Atmosphäre entspannt.

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Dadurch füllte sich ein Teil des Porenraums der Kugeln mit sauerstofffreiem Wasser und der Kontakt mit Luftsauerstoff beim Einbau der BACTRAPS wurde vermieden. Die BACTRAPS wurden in diesem Wasser bis zur Installation in den Brunnen gelagert.

2.4.7 Einbau in die Grundwassermessstelle

Die BACTRAPS wurden mittels MLPS in acht Tiefen der Messstelle Saf ZZ 23/00 (Fig. 2-2) eingebbracht. Um eine gleichmäßige Durchströmung der BACTRAPS mit dem Grundwasser zu gewährleisten, wurden sie in Edelstahlkäfigen eingefasst (Fig. 2-3). Die Edelstahlkäfige dienten dabei als Abstandhalter um direkte Kontakte mit dem Pegelrohr sowie der Packermembran zu vermeiden.

2.4.8 Probenaufbereitung

Nach einer Expositionszeit von 51 Tagen wurden die Mikrokosmen aus der Messstelle entnommen und zur Konservierung bei -15 °C gelagert.

Die Extraktion der Fettsäuren erfolgte nach einer modifizierten Methode von Bligh und Dyer (1959). Dabei wurden 2 ml Methanol und Dichlormethan im Verhältnis 1:3 zugegeben. Für die Herstellung der Fettsäuremethylester (FAME) dienten 300 µl eines Derivatisierungsreagens aus Chlortrimethylsilan und wasserfreiem Methanol im Verhältnis 1:8. Die Veresterung erfolgte bei einer Temperatur von 70 °C und einer Reaktionsdauer von zwei Stunden. Die FAME wurden in Hexan gelöst und anschließend mittels GC-MS bzw. GC-IRMS analysiert.

2.4.9 Identifikation der Fettsäuremethylester (FAME) mittels GC-MS

Die Bestimmung der FAME erfolgte mit einem Gaschromatographen (HP 6890 Series, Agilent Technology, USA) mit gekoppeltem Massenspektrometer (HP 5973 Mass Selective Detector, Agilent Technologie, USA). Zur gaschromatographischen Trennung der FAME wurde eine BPX 5-Säule (30m x 0,32 mm x 0.25 µm FD; Agilent Technologie, USA) eingesetzt. Das Temperaturprogramm beginnt bei 120 °C. Nach 4 Minuten steigt die Temperatur mit 4 °C min⁻¹ auf 250 °C und schließlich mit 20 °C min⁻¹ auf 300 °C, die abschließend 10 Minuten konstant gehalten wurden. Die FAME sind mit Hilfe des *Bacterial Acid Methyl Ester*-Standards (BAME, Supelco) durch den Vergleich der Retentionszeiten und Massenspektren identifiziert und auf ihre Reinheit geprüft worden. Die Fettsäuren werden nach folgender Nomenklatur benannt: A:BwC, wobei A die Anzahl der Kohlenstoffatome, B die Anzahl der Doppelbindungen und C die Entfernung der nächsten Doppelbindung vom aliphatischen Ende des Moleküls repräsentiert (ω -Nomenklatur). Durch die Vorsilben *iso* (i-) und *anteiso* (a-) wird die Position der Methylverzweigung gekennzeichnet.

2.4.10 Analyse der Kohlenstoffisotopenverhältnisse der FAME mittels GC-C-IRMS

Zur Analyse der Isotopenverhältnisse der FAME wurde ein Gas *Chromatography-Combustion Isotope-Ratio-Monitoring Mass Spectrometer System* (GC-C-IRMS) verwendet. Das System besteht aus einem Gaschromatograph (HP 6890 Series, Agilent Technology, USA), der an einen Verbrennungsofen, ein GC-C-III-Interface (Combustion III, Finnigan MAT, Deutschland), eine Wasserfalle (Nafion®-Membran) und ein Massenspektrometer (IRMS MAT 252, Finnigan MAT, Deutschland) gekoppelt ist. Eine Beschreibung zur verwendeten Gerätekonfiguration geben Richnow et al. (2003a).

Zur Auftrennung der FAME wurde eine BPX 5-Säule (50m x 0,32 mm x 0,5 µm FD; SGE GmbH, Deutschland) mit folgendem Temperaturprogramm verwendet: eine Minute 70 °C isotherm, 20 °C min⁻¹ auf 170 °C, 2 °C min⁻¹ auf 280 °C und mit 20 °C min⁻¹ auf 300 °C, die für 5 Minuten isotherm gehalten wurden.

2.4.11 Berechnungen

Die Isotopensignaturen werden als δ-Notation [‰] relativ zu dem internationalen VPDB-Standard (**Vienna PeeDee Belemnite** mit ¹³C/¹²C = 0,0112372) (Clark and Fritz, 1997; Hoefs, 1997) angegeben (Gleichung 1-1).

$$\text{Gleichung 1-1: } \delta^{13}\text{C} = \frac{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{Probe}} - \left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{VPDB-Standard}}}{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{VPDB-Standard}}} \times 1000$$

Während der Methylierung der Fettsäuren (FA) zu Fettsäuremethylestern (FAME) wird ein zusätzliches Kohlenstoffatom in das Produkt eingefügt, so dass die ursprüngliche Isotopensignatur aller FA konstant verändert wird. Die gemessenen δ¹³C-Werte der FAME wurden deshalb nach der Gleichung 1-2 korrigiert, um die Isotopenverhältnisse der Fettsäuren (δ¹³C_{FA}) zu ermitteln (Abraham et al., 1998).

$$\text{Gleichung 1-2: } \delta^{13}\text{C}_{\text{FA}} = \frac{[(C_n + 1) \times \delta^{13}\text{C}_{\text{FAME}} - \delta^{13}\text{C}_{\text{MeOH}}]}{C_n}$$

C_n entspricht der Anzahl der Kohlenstoffatome in den Fettsäuren, δ¹³C_{FAME} ist die Isotopensignatur der Fettsäuremethylester (FAME) und δ¹³C_{MeOH} die Isotopensignatur des für die Methylierung benutzten Methanols [-38,15 ‰].

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Mit der Rayleigh-Gleichung (Gleichung 1-3) wird die Beziehung zwischen der Konzentrationsänderung der mikrobiell umgesetzten Substanz in Verbindung mit einer Änderung der Isotopensignatur durch den Isotopenfraktionierungsfaktor α dargestellt (Hoefs, 1997).

Gleichung 1-3:
$$\frac{R_t}{R_0} = \left(\frac{C_t}{C_0} \right)^{\left(\frac{1}{\alpha} - 1 \right)}$$

R_0 und R_t geben die Isotopenzusammensetzung (Gleichung 1-4), C_t und C_0 die Konzentration des Substrats zum Zeitpunkt t und 0 der mikrobiellen Umsetzung an.

Gleichung 1-4:
$$\frac{R_t}{R_0} = \frac{\delta_t^{13}\text{C} \times 1000}{\delta_0^{13}\text{C} \times 1000}$$

Der prozentuale biologische Abbau der residualen Schadstofffraktion wird nach Gleichung 1-5 berechnet (Meckenstock et al., 2002). Dabei wird (C_t/C_0) durch Umstellen der Rayleigh-Gleichung ersetzt (Gleichung 1-6), so dass der prozentuale biologische Abbau unter Verwendung von Isotopenwerten quantifiziert werden kann.

Gleichung 1-5:
$$B[\%] = \left(1 - \frac{C_t}{C_0} \right) * 100$$

Gleichung 1-6:
$$B[\%] = \left[1 - \left(\frac{R_t}{R_0} \right)^{\left(\frac{1}{\alpha} - 1 \right)} \right] * 100$$

Die Veränderung der Kohlenstoffisotopensignatur und der Konzentration des gelösten anorganischen Kohlenstoffs (DIC) innerhalb einer Schadstofffahne dienen als biogeochemischer Indikator und können zur Abschätzung der Schadstoffmineralisierung nach Gleichung 1-7 verwendet werden (Fischer et al., 2004).

Gleichung 1-7
$$A \times \delta^{13}\text{C}_{\text{DIC(GA)}} + (1 - A) \times \delta^{13}\text{C}_{\text{DIC(Q)}} = \delta^{13}\text{C}_{\text{DIC(M)}}$$

Dabei bezeichnet $\delta^{13}\text{C}_{\text{DIC(GA)}}$ das Kohlenstoffisotopenverhältnis des DIC im Grundwasseranstrom der Kontamination. $\delta^{13}\text{C}_{\text{DIC(Q)}}$ steht für das Kohlenstoffisotopenverhältnis des während der Schadstoffmineralisierung gebildeten DIC. Dabei wird angenommen, dass $\delta^{13}\text{C}_{\text{DIC(Q)}}$ die Isotopensignatur des nicht abgebauten

Schadstoffs (Ausgangssubstrats) in der Quelle trägt (Hunkeler et al., 1999b). $\delta^{13}\text{C}_{\text{DIC}(\text{M})}$ entspricht dem $^{13}\text{C}/^{12}\text{C}$ -Verhältnis des DIC, das innerhalb der Schadstofffahne gemessen wurde. A und (1-A) stellen die Mischungsanteile der beiden Ausgangsisotopenverhältnisse ($\delta^{13}\text{C}_{\text{DIC}(\text{GA})}$ und $\delta^{13}\text{C}_{\text{DIC}(\text{Q})}$) dar, aus denen $\delta^{13}\text{C}_{\text{DIC}(\text{M})}$ resultiert. Anhand der DIC-Konzentration im Grundwasseranstrom der Schadstofffahne ($C_{\text{DIC}(\text{GA})}$) und mit Hilfe von A kann auf die Menge des in der Schadstofffahne gebildeten DIC ($C_{\text{DIC}(\text{B})}$) geschlossen werden (Gleichung 1-8).

$$\text{Gleichung 1-8} \quad C_{\text{DIC}(\text{B})} = C_{\text{DIC}(\text{GA})} \frac{(1 - A)}{A}$$

Schließlich ist eine Abschätzung der mineralisierten Schadstofffraktion ($C_{\text{Schadstoff}}$) anhand des gebildeten DIC ($C_{\text{DIC}(\text{B})}$) möglich (Gleichung 1-9).

$$\text{Gleichung 1-9} \quad C_{\text{Schadstoff}} [\text{M}] = \frac{C_{\text{DIC}(\text{B})}}{Z_c}$$

Z_c bezeichnet die Anzahl der Kohlenstoffatome des Schadstoffmoleküls, die bei der Mineralisierung umgesetzt werden.

2.5 Ergebnisse

Die Messstelle Saf ZZ 23/00 im nordöstlichen Abstrom der Schadstofffahne (Fig. 2-1) wies in Monitoringkampagnen der letzten Jahre BTEX-Konzentrationen in einer Schwankungsbreite von ca. 40 bis 700 mg L⁻¹ auf. Im Dezember 2004 wurde mit einem MLPS die vertikale hydrogeochemische Struktur der BTEX-Schadstofffahne untersucht und der *in situ* Benzolabbau anhand der Isotopensignatur des Benzols und DIC charakterisiert. Um den mikrobiellen Benzolabbau innerhalb der verschiedenen geochemischen Zonen untersuchen zu können, wurde parallel ein BACTRAP-Experiment in acht Tiefen durchgeführt.

2.5.1 Geochemie

Benzol stellt mit einem Anteil von mehr als 99 % die wesentliche BTEX Komponente dar. Toluol, Ethylbenzol sowie die Xyole überschreiten als Summenparameter in keinem Beprobungshorizont Gesamtkonzentrationen von 2 mg L⁻¹ (nicht dargestellt). Die Benzolfahne ist vertikal strukturiert (Fig. 2-4). Der maximale Benzolgehalt von > 6.400 µmol L⁻¹ (> 500 mg L⁻¹) wurde in 14 m Tiefe bestimmt. Zum oberen und unteren Schadstofffahnenrand nehmen die Benzolkonzentrationen bis auf 415 bzw. 1670 µmol L⁻¹ kontinuierlich ab.

Analyse des anaeroben Benzolabbaus:

Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen

Die Isotopensignatur des Benzols weist in den mittleren Tiefen $\delta^{13}\text{C}$ -Werte zwischen -28,0 ‰ und -29,0 ‰ auf (Fig. 2-4), die im wesentlichen den Werten des Benzols im Schadenszentrum (-28,5 bis -29,5 ‰) entsprechen (Fischer et al., 2004; Vieth et al., 2005). In dieser Zone der Kontaminationsfahne kann daher über die Isotopenfraktionierungsmethode kein bzw. nur ein geringer biologischer Abbau von Benzol nachgewiesen werden (Tab. 2-1). Im oberen und unteren Bereich des Vertikalprofils der Messstelle Saf ZZ 23/00 ist eine ^{13}C -Anreicherung in der residualen Benzolfraktion um 3 bzw. 6 ‰ festzustellen, die offensichtlich auf Abbauprozesse hinweist. Eine Quantifizierung des prozentualen biologischen Schadstoffabbaus (B %) wurde unter Annahme sulfatreduzierender bzw. methanogener Bedingungen durchgeführt. Die Berechnung basiert auf den von Mancini et al. (2003) für den Benzolabbau bestimmten Fraktionierungsfaktoren (α), mit α von 1,0036 für sulfatreduzierende und α von 1,0019 für methanogene Abbaubedingungen. Der mikrobielle Benzolabbau unter der Verwendung der schweren Isotopensignatur der Quelle von -28,5 und dem höheren Isotopenfraktionierungsfaktor für sulfatreduzierende Bedingungen (1,0036) zeigt Abbaugrade bis zu 59 % am oberen und bis 83 % am unteren Fahnenrand (Tab. 2-1). Die Berechnung mit dem Fraktionierungsfaktor für methanogene Bedingungen (1,0019) unter Annahme einer relativ leichten Benzolquelle (-29,5 ‰) ergibt einen entsprechend höheren Abbau von 89 bis 98 % für die Fahnenränder. Mit Hilfe der hier berechneten Variabilität der Abbaugrade und den gemessenen Benzolkonzentrationen (C_t) in den einzelnen Tiefen der Messstelle Saf ZZ 23/00 konnte auf die theoretischen Schadstoffkonzentrationen der Kontaminationsquelle (C_0) zu Beginn eines biologischen Abbaus ($t=0$) zurückgerechnet werden (Tab. 2-1). Die Berechnung des Benzolabbaus für sulfatreduzierende Bedingungen zeigte, dass diese theoretisch bestimmten Konzentrationen von > 10.000 $\mu\text{mol L}^{-1}$ im unteren Aquiferbereich in etwa mit den Benzolkonzentrationen der Schadstoffquelle (>12.800 $\mu\text{mol L}^{-1}$) vergleichbar sind. Unter der Annahme methanogener Abbaubedingungen ergeben sich hypothetische Benzolkonzentrationen für die Schadstoffquelle von > 90.000 μmol , die die physikalische Löslichkeit des Benzols im Wasser überschreiten und daher unrealistisch sind. Das unterstreicht frühere Ergebnisse, nach denen die Methanogenese im Vergleich zur Sulfatreduktion am Standort keinen wesentlichen Anteil am Benzolabbau hat (Fischer et al., 2004). Unter sulfatreduzierenden Bedingungen ergibt sich ein Benzolabbau von bis zu 9.000 $\mu\text{mol L}^{-1}$ (ca. 700 mg L^{-1}) auf dem Fließweg zwischen Quelle und Messpunkt im unteren Fahnenbereich, welcher aufgrund der hohen Benzolkonzentrationen im Schadenszentrum durchaus möglich sein könnte. Im oberen Rand der Fahne ergibt sich ein entsprechend geringerer Benzolabbau zwischen 200 bis 700 $\mu\text{mol L}^{-1}$ (16-55 mg L^{-1}).

In unbelasteten Bereichen des Aquifers werden Sulfatkonzentrationen von > 10.000 $\mu\text{mol L}^{-1}$ nachgewiesen, so dass Sulfat den wichtigsten Elektronenakzeptor für mikrobielle

Abbauprozesse darstellt (Fischer et al., 2004; Vieth et al., 2005). Im Vertikalprofil schwanken die Sulfatkonzentrationen zwischen ca. $20 \mu\text{mol L}^{-1}$ in einer Tiefe von 14 m und $> 5.000 \mu\text{mol L}^{-1}$ am oberen Fahnenrand (Fig. 2-4). Verglichen mit unbelasteten Bereichen des Aquifers deutet die Abnahme des Sulfats infolge einer mikrobiellen Sulfatreduktion auf eine deutliche Sulfatsenke in der Fahne hin.

Tab. 2-1: Berechnung des biologischen Abbaus [B%] für sulfatreduzierende ($\alpha_S = 1.0036$) und methanogene Bedingungen ($\alpha_M = 1.0019$) unter Berücksichtigung der Schwankungsbreite der Isotopensignatur von Benzol in der Quelle ($R_0 = -28,5 \text{ ‰}$ bis $-29,5 \text{ ‰}$)

Tiefe [m]	$C_{\text{Benzol}} [\mu\text{mol L}^{-1}]$	$\delta^{13}\text{C}_{\text{Benzol}} [\text{‰}]$	$B \%(\text{S})^1$ [%]	$B \%(\text{M})^2$ [%]	$C_0(\text{S})^3 [\mu\text{mol L}^{-1}]$	$C_0(\text{M})^4 [\mu\text{mol L}^{-1}]$
10,4	415	$-25,4 \pm 0,4$	59	89	1013	3857
11,4	2557	$-28,2 \pm 0,2$	9	51	2795	5208
12,2	4901	$-28,0 \pm 0,4$	13	55	5608	10886
13,2	6159	$-28,6 \pm 0,2$	-4	37	5916	9825
14,2	6419	$-27,4 \pm 0,2$	26	68	8725	19743
15,6	1840	$-22,3 \pm 0,2$	83	98	10871	91177
16,7	1668	$-22,3 \pm 0,4$	83	98	9916	83647

¹ Berechnung des biologischen Abbaus für sulfatreduzierende Bedingungen unter Annahme von $R_0 = -28,5 \text{ ‰}$

² Berechnung des biologischen Abbaus für methanogene Bedingungen unter Annahme von $R_0 = -29,5 \text{ ‰}$

³ Berechnung der Benzolkonzentration der Quelle auf Grundlage von $B \%(\text{S})^1$

⁴ Berechnung der Benzolkonzentration der Quelle auf Grundlage von $B \%(\text{M})^2$




Fig. 2-4: Links vertikale Verteilung der Eisen(II)-(◊), Eisen(III)-(●), Nitrat-(★), Mangan-(▲), Sulfat-(■), und rechts der Benzolkonzentrationen (●) sowie der Kohlenstoffisotopenverhältnisse des Benzols (■).

Der höchste Nitratgehalt von $244 \mu\text{mol L}^{-1}$ wurde am oberen Rand der Fahne im Übergangsbereich zur ungesättigten Zone festgestellt (Fig. 2-4). In tieferen Zonen konnte

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

kein Nitrat nachgewiesen werden. Im Vergleich zum Sulfat steht für Abbauprozesse Nitrat als Elektronenakzeptor also nur in der oberen Zone des Aquifers in relativ geringen Konzentrationen zur Verfügung.

Die Konzentrationen von Mangan betragen etwa $180 \mu\text{mol L}^{-1}$ am oberen Rand der Fahne und reduzieren sich mit zunehmender Tiefe kontinuierlich auf Konzentrationen um $44 \mu\text{mol L}^{-1}$ (Fig. 2-4). Eine Manganreduktion könnte in der oberen Zone der Kontaminationsfahne einen Beitrag zum Schadstoffabbau liefern. Insgesamt sind die Mangankonzentrationen für einen signifikanten Schadstoffabbau jedoch zu gering.

Die Konzentrationen von Eisen(III) liegen im Bereich von ca. $20-60 \mu\text{mol L}^{-1}$ mit Ausnahme von ca. $170 \mu\text{mol L}^{-1}$ in einer Tiefe von 11,5 m. Die Eisen(II)-Werte schwanken zwischen etwa $80-470 \mu\text{mol L}^{-1}$ (Fig. 2-4). Erhöhte Fe(II)-Konzentrationen zwischen 11-14 m weisen auf eine Fe(II)-Mobilisierung hin, die durch Abbauprozesse verursacht sein könnte. Auffällig ist, dass die Bereiche erhöhter Fe(II)-Gehalte (11-14 m) durch die geringsten Sulfid-Konzentrationen gekennzeichnet sind. Diese liegen zwar generell bei $\leq 6 \mu\text{mol L}^{-1}$ (nicht dargestellt), aber die Fällung von Fe(II) durch Sulfid könnte zu einer Unterschätzung der Eisenreduktion beitragen. Gefälltes Eisensulfid verbleibt an der Aquifermatrix und kann im Grundwasser nicht quantitativ nachgewiesen werden, so dass der Anteil der Eisenreduktion anhand der Eisenmobilisierung kaum quantifizierbar ist. Es kann aber grundsätzlich davon ausgegangen werden, dass Eisen(III) am Standort keinen bedeutenden Elektronenakzeptor für Abbauprozesse darstellt (Wachter, 2004).

Die Sauerstoffkonzentrationen liegen in der Messstelle mit Ausnahme des Übergangsbereichs zur ungesättigten Zone unterhalb von $40 \mu\text{mol L}^{-1}$, so dass Sauerstoff als Elektronenakzeptor in tieferen Zonen der Fahne nicht zur Verfügung steht (nicht dargestellt).

Die DIC-Konzentrationen verhalten sich proportional zum Schadstoff (Fig. 2-5). Die höchsten Konzentrationen von $> 17.000 \mu\text{molC L}^{-1}$ wurden für den am stärksten kontaminierten Bereich zwischen 11-14 m registriert und nehmen am oberen und unteren Fahnenrand ab. Die maximal gemessenen DIC-Werte des Vertikalprofils entsprechen in der Größenordnung den DIC-Konzentrationen, die für die Hauptschadenszonen der BTEX-Fahne ermittelt wurden (Fischer et al., 2004). Im unteren Bereich der Fahne geht die DIC-Konzentration bis auf rund $1.300 \mu\text{molC L}^{-1}$ zurück. Die Verteilung der verschiedenen Spezies des DIC im Grundwasser ist pH-Wert abhängig. Im Bereich der Schadstofffahne wurden pH-Werte von 6,5 - 7,0 bestimmt, so dass entsprechend des Kalk-Kohlensäure-Gleichgewichts davon auszugehen ist, dass Hydrogencarbonat die dominierende Spezies des DIC am Standort darstellt und Fällungsreaktionen ausgeschlossen werden können (Clark and Fritz, 1997). Dieser Trend wird auch im Tiefenprofil der Messstelle Saf ZZ 23/00 beobachtet, wo neutrale pH-Werte von ca. 7,0 für alle Tiefen bestimmt wurden.

Unter der Annahme, dass die Sulfatreduktion der dominierende Prozess des mikrobiellen Schadstoffabbaus am Standort ist (Wachter, 2004), kann folgende Abschätzung vorgenommen werden. Nach Wiedemeier et al. (1999) werden bei der Mineralisierung von 1.000 µmol Benzol unter der Bildung von 6.000 µmol CO₂ 3.750 µmol Sulfat verbraucht (Wiedemeier et al., 1999). Eine mikrobielle Oxidation von rund 1.300 µmol L⁻¹ Benzol unter Bildung von etwa 8.000 µmolC L⁻¹ DIC kann somit theoretisch eine Sulfatsenke von ca. 5.000 µmol L⁻¹ bewirken, wenn der Aufbau zelleigener Biomasse unberücksichtigt bleibt. Verglichen mit der DIC-Konzentration von ca. 8.300 µmolC L⁻¹ im Anstrom der BTEX-Fahne konnte eine maximale Konzentrationszunahme um ca. 10.000 µmolC L⁻¹ im stark kontaminierten Bereich der Messstelle Saf ZZ 23/00 beobachtet werden. Aus der maximalen Differenz der DIC-Konzentrationen im Grundwasseranstrom und im Bereich der Messstelle Saf ZZ 23/00 berechnet sich ein theoretisches Abbaupotenzial von ca. 1.700 µmol L⁻¹ Benzol. Aus den beiden Berechnungsansätzen resultiert ein Unterschied von etwa 400 µmol Benzol.




Fig. 2-5: Vertikale Verteilung der Kohlenstoffisotopenverhältnisse von Methan [■], DIC [●] und den Anreicherungsfaktoren [▼] (links) sowie der Methan- [■] und DIC-Konzentrationen [●] (rechts) in der Messstelle Saf ZZ 23/00.

In kontaminierten Aquiferen gilt die Zunahme von Methan als ein Indiz für Biodegradation unter methanogenen Milieubedingungen (Wiedemeier et al., 1999). Die CH₄-Konzentrationen im Grundwasser können außerdem durch Methanoxidation beeinflusst werden (Breukelen and Griffioen, 2004; Grossmann et al., 2002). Die Methanogenese führt zur Bildung von isotopisch leichtem Methan während die aerobe Methanoxidation mit einer Anreicherung schwerer Isotope verbunden ist. Um Methanbildungs- oder Abbauprozesse voneinander zu unterscheiden, kann der Anreicherungsfaktor $\varepsilon_{\text{DIC-CH}_4}$, die Differenz zwischen der Kohlenstoffisotopensignatur des DIC und Methans, genutzt werden (Fischer et al., 2004).

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Geringe $\varepsilon_{\text{DIC-CH}_4}$ -Werte von < 20 ‰ gelten als Indiz für Methanoxidation, höhere Anreicherungsfaktoren deuten auf Methanbildung hin (Acetogenese 10-50 ‰; > 50 ‰ CO₂-Reduktion) (Fischer et al., 2004; Whiticar, 1999). Die berechneten Anreicherungsfaktoren für das Tiefenprofil der Messstelle Saf ZZ 23/00 variieren von 17-26 ‰ (Fig. 2-5). Im Beprobungshorizont bei 10,4 m wurden sowohl die geringsten Methankonzentrationen als auch die geringsten Anreicherungsfaktoren bestimmt, so dass im Übergang zur ungesättigten Zone des Aquifers Methanoxidation denkbar wäre. Mit zunehmender Tiefe trägt wahrscheinlich die Methanogenese zum Abbau der BTEX bei und führt zu einer Erhöhung der CH₄-Konzentration auf ca. 350 µmol L⁻¹, was etwa einem Abbau von 93 µmol (7 mg L⁻¹) Benzol entspricht (Wiedemeier et al., 1999). Wie anhand der Anreicherungsfaktoren abgeleitet werden kann, ist die Acetatgärung hierbei vermutlich der dominierende Methanbildungsprozess. Die relativ niedrigen Methankonzentrationen insgesamt zeigen jedoch, dass die Methanogenese keinen signifikanten Beitrag am Schadstoffabbau liefert.

Bei der Mineralisation von Benzol entsteht Kohlendioxid, das die gleiche Isotopensignatur wie das Ausgangssubstrat trägt (Fischer et al., 2004; Hunkeler et al., 2001). Im Anstrom der Fahne liegt die natürliche DIC-Isotopensignatur im Mittel bei -19,5 ‰ (Fischer et al., 2004), im Bereich der Messstelle Saf ZZ 23/00 sind die $\delta^{13}\text{C}_{\text{DIC}}$ -Werte hingegen durchschnittlich um ca. 5,5 ‰ leichter (-24,5 bis -25,6 ‰) und zeigen keine markanten Unterschiede über die Tiefe (Fig. 2-5). Werte dieser Größenordnung sind generell im stärker kontaminierten Bereich der BTEX-Fahne beobachtet worden (Fischer et al., 2004). Anhand dieser Veränderung des Isotopensignals über die Fließstrecke wurde die Umsetzung von mineralisiertem Benzol berechnet (Gleichung 1-7 - Gleichung 1-9). Als Kohlenstoffisotopenverhältnis ($\delta^{13}\text{C}_{\text{DIC(GA)}}$) bzw. Konzentration ($C_{\text{DIC(GA)}}$) des DIC im Grundwasseranstrom wurde ein mittlerer Wert von -19,5 ‰ für $\delta^{13}\text{C}_{\text{DIC(GA)}}$ bzw. 8.300 µmol C L⁻¹ für $C_{\text{DIC(GA)}}$ angenommen (Fischer et al., 2004). Das durchschnittliche Isotopenverhältnis des DIC im Profil der Messstelle Saf ZZ 23/00 ($\delta^{13}\text{C}_{\text{DIC(M)}}$) betrug ca. -25,0 ‰. Unter der Annahme identischer Isotopensignaturen des Benzols im Schadenszentrum und des DIC, das infolge einer Mineralisierung des Schadstoffs entsteht, wurde für die Quantifizierung ein $\delta^{13}\text{C}_{\text{DIC(Q)}}$ Wert von -28,5 ‰ angenommen, da im Fall einer vollständigen Mineralisierung die Isotopenbilanz geschlossen ist. Die Abschätzung ergibt für die Fließstrecke zwischen dem Zentrum der Schadstofffahne und der Messstelle Saf ZZ 23/00 eine Mineralisierung von etwa 2.160 µmol L⁻¹ Benzol (169 mg L⁻¹).

Beim aeroben Benzolabbau kann eine Isotopenfraktionierung zwischen nicht abgebautem Benzol und gebildetem DIC von bis zu 4 - 6 ‰ auftreten (Hunkeler et al., 2001), wenn Biomasse gebildet wird oder der Abbau unvollständig ist. Die Fraktionierung zwischen Benzol und DIC beim anaeroben Abbau ist noch nicht untersucht. Würde eine Fraktionierung

zwischen nicht abgebauten Benzol und gebildetem DIC von maximal 6 ‰ ($\delta^{13}\text{C}_{\text{DIC}(\text{Q})} = -34,5\text{ ‰}$) bei der Quantifizierung des anaeroben Benzolabbaus berücksichtigt werden, so verringert sich die Menge des mineralisierten Benzols (808 µmol L⁻¹ bzw. 63 mg L⁻¹) erheblich. Die Abschätzung des Abbaus setzt voraus, dass die Quellen organischen Materials für das DIC isotopisch charakterisiert sind. Am Standort Zeitz steht vorherrschend Benzol zur Verfügung. Wenn andere Quellen wie natürlicher gelöster Kohlenstoff (DOC) oder andere Schadstoffe zur DIC-Bildung beitragen, ist eine Abschätzung schwierig. Es ist daher offensichtlich, dass die Abschätzung des Abbaupotenzials über eine Isotopenbilanz mit Unsicherheiten verbunden ist.

2.5.2 Ergebnisse der tiefenorientierten Untersuchungen mit den BACTRAPs

Die BACTRAPs mit und ohne Benzolbeaufschlagung (¹²C-Benzol, ¹³C-Benzol, Kontrolle) zeigen für alle Beprobungshorizonte vergleichbare Fettsäremuster. Dominierende Fettsäuren sind C16:0 (> 40 %) und C18:0 (> 20 %) sowie die ungesättigten Fettsäuren C16:1 und C18:1 mit einem prozentualen Anteil von jeweils ca. 10 %. Untergeordnet, in geringen Konzentrationen (< 10 %), konnten die Fettsäuren C14:0, C15:0, i-C15:0, a-C15:0, C18:2 sowie eine weitere C18:1 Fettsäure mittels GC-MS identifiziert werden.

Da sich die Fettsäremuster der mit Benzol beladenen und unbeladenen BACTRAPs nicht wesentlich voneinander unterscheiden, konnte weder ein Substrateinfluss noch eine Abhängigkeit des mikrobiellen Konsortiums vom geochemischen Milieu in den einzelnen Tiefen festgestellt werden. Die Fettsäuren liefern daher keinen Hinweis auf eine unterschiedliche Besiedlung der BACTRAPs in den verschiedenen Beprobungshorizonten. Verglichen mit einem BACTRAP-Experiment, das in einer Messstelle im weiter entfernten Grundwasserabstrom durchgeführt wurde (Geyer et al., 2005), sind die Fettsäremuster im Bereich der Messstelle Saf ZZ 23/00 homogener und weniger komplex. Das lässt auf eine geringere Biodiversität auf dem BACTRAP schließen.

Die Kohlenstoffisotopensignaturen der Fettsäuren ($\delta^{13}\text{C}_{\text{FA}}$) aus dem Kontrollexperiment variieren zwischen -23 und -34 ‰. Die Isotopenanalysen der Proben aus dem Markierungsexperiment zeigten die höchsten ¹³C-Anreicherungen (zwischen 28 und 468 ‰) für die C16:1 Fettsäure (Fig. 2-6). Eine etwas geringere Markierung wurde für die Fettsäuren i-C15:0, C14:0, C16:0 und C18:1 (-20 ‰ bis 145 ‰) bestimmt. Die signifikante ¹³C-Anreicherung von Fettsäuren ist auf die Transformation von ¹³C aus dem Benzol in die Biomasse zurückzuführen. Keine signifikante Transformation der ¹³C-Markierung des Benzols wurde in die a-C15:0, C15:0 und C18:0 mit $\delta^{13}\text{C}_{\text{FA}}$ -Werten von -14 bis -25 ‰ beobachtet (Fig. 2-6).

Die Isotopensignatur der Fettsäuren einer Spezies spiegelt, abgesehen von geringen Unterschieden, die auf Isotopenfraktionierungsprozesse beruhen, im Wesentlichen das

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Isotopenverhältnis des Wachstumssubstrates wider (Abraham et al., 1998). Auch in Experimenten mit markierten Kohlenstoffsubstraten ist die Isotopensignatur bei PLFA verschiedener Kettenlänge etwa gleich (Pelz et al., 2001a) und auch kaum vom Wachstumsstadium beeinflusst (Abraham et al., 1998). Der Kohlenstoff wird also etwa gleichermaßen zum Aufbau von Membranlipiden verwendet. Die unterschiedlich starke Markierung zeigt daher die Besiedlung des BACTRAPS durch mikrobielle Konsortien, welche ^{13}C -Benzol unterschiedlich effektiv als Kohlenstoffquelle nutzen. Zum Beispiel scheinen Organismen, die unmarkierte C18:0, a-C15:0 oder C15:0 Fettsäuren besitzen, das markierte Benzol nicht oder nur kaum zu metabolisieren und bevorzugen andere Kohlenstoffquellen zur Synthese. Als zusätzliche Kohlenstoffquellen, die den Mikroorganismen zur Verfügung stehen können und die durch eine natürliche Isotopensignatur gekennzeichnet sind, kommen aromatische Kohlenwasserstoffe wie zum Beispiel BTEX aus der Kontamination sowie auch DOC in Frage.




Fig. 2-6: Übersicht der Isotopensignaturen der Fettsäuren $\delta^{13}\text{C}_{\text{FA}} [\text{\textperthousand}]$ in den einzelnen Beprobungshorizonten. Die Abkürzungen und Nomenklatur der Fettsäuren sind im Kapitel Material und Methoden (2.4.9) erklärt.

Anhand der Intensität der ^{13}C -Anreicherung in den Fettsäuren konnten Zonen unterschiedlicher mikrobiologischer Aktivität identifiziert werden. Im obersten Bereich der untersuchten Messstelle (1. und 3. Tiefe) zeigt die isotopische Anreicherung in der C16:1 von >350 ‰ demnach die größte mikrobielle Aktivität an. Im mittleren Tiefenintervall (12 m) wurden die geringsten Markierungen (40 ‰) für C16:1 nachgewiesen. Im unteren Bereich der Messstelle (14-15 m) zeigt die C16:1 mittlere ^{13}C -Anreicherung von ca. 94 ‰ (14,2 m) bzw. 188 ‰ (15,6 m). Die Ergebnisse verdeutlichen, dass nicht nur in den Randbereichen

der Schadstofffahne sondern auch in den stark benzolbelasteten Zonen des Aquifers ein mikrobieller Benzolabbau stattfindet.

2.6 Diskussion

2.6.1 Geochemie

Die Abschätzung des Benzolabbaus anhand der Isotopensignaturen des Benzols bzw. DIC und Methan hat gezeigt, dass bevorzugt im oberen und hauptsächlich im unteren Randbereich der Fahne ein *in situ* Abbau stattfindet. Im Tiefenintervall von ca. 12-14 m geben die $\delta^{13}\text{C}_{\text{Benzol}}$ -Werte keinen Hinweis auf mikrobiellen Abbau, da sie ungefähr das Isotopensignal des Quellbereichs (-28,5‰) tragen. Vermutlich ist im Bereich der sehr hohen BTEX-Konzentrationen der Anteil des mikrobiell abgebauten Benzols im Verhältnis zum nicht umgesetzten zu gering, um eine Veränderung der Isotopensignatur eindeutig nachweisen zu können.

Die konservative Abschätzung des Abbaus auf dem Fließweg zwischen Quelle und Messpunkt anhand der Isotopenmethode ergab für sulfatreduzierende Bedingungen einen Benzolabbau von bis zu 9.000 $\mu\text{mol L}^{-1}$ (ca. 700 mg L^{-1}). Der gemittelte Abbau über alle Tiefen liegt bei ca. 3.000 $\mu\text{mol L}^{-1}$ (ca. 230 mg L^{-1}). Mit Hilfe der Konzentrationen und Isotopensignaturen des DIC wurde ein Benzolabbau von etwa 2.160 $\mu\text{mol L}^{-1}$ (ca. 169 mg L^{-1}) abgeschätzt. Aus der Elektronenbilanz über die Sulfatsenke ergab sich ein Abbaupotenzial von 1.700 $\mu\text{mol L}^{-1}$ (ca. 133 mg L^{-1}) Benzol. Etwa 350 $\mu\text{mol L}^{-1}$ (ca. 28 mg L^{-1}) Benzol könnten zusätzlich über methanogene Prozesse abgebaut worden sein. In der Summe errechnet sich ein möglicher Benzolabbau infolge Sulfatreduktion und Methanogenese von 2.050 μmol (ca. 160 mg). Dieser Wert entspricht etwa der Abschätzung der Mineralisation über das DIC. Die Quantifizierung des Abbaus anhand der Benzolisotope liegt etwa um ein Drittel höher aber in der gleichen Größenordnung. Die verschiedenen Berechnungsansätze liefern in der Größenordnung übereinstimmende Ergebnisse, so dass die Werte für den berechneten Benzolabbau realistisch erscheinen.

Eine Diskussion der Abweichungen der verschiedenen vorgestellten Ansätze zur Abschätzung des anaeroben Benzolabbaus gestaltet sich schwierig. Das Karbonatsystem (DIC) könnte durch Fällungen beeinflusst werden. Dies scheint aufgrund neutraler pH-Werte am Standort jedoch unwahrscheinlich zu sein. Ausgasungen in erheblichem Umfang sind nur im Bereich der Zonen mit direktem Kontakt zur ungesättigten Bodenzone zu erwarten und über die Mächtigkeit des gesamten Aquifers eher als gering anzusehen. In der Elektronenbilanz fehlt im Wesentlichen die Eisenmobilisierung, weil Fe(II) unter sulfatreduzierenden Bedingungen gefällt wird und für quantitative Abschätzungen nicht erfasst werden kann.

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Die Beurteilung des mikrobiellen Schadstoffabbaus über die Methode der Isotopenfraktionierung gibt vermutlich eine realistische Größe für den Abbau zwischen Quelle und Messpunkt. Die Methode führt eher zu einer konservativen Abschätzung, wobei der mikrobielle Abbau nicht wesentlich überschätzt werden sollte. Mischungsprozesse auf dem Fließweg von Grundwasserströmen mit einerseits kaum und andererseits stark abgebauten Schadstoffmengen können zu einer Unterschätzung des tatsächlichen Abbaus führen (Fischer et al., 2004; Richnow et al., 2003a). Zu einer Überschätzung des Abbaus führt die Isotopenmethode in der Regel nicht.

2.6.2 BACTRAP

Die Auswertung des BACTRAP-Experiments hat gezeigt, dass in allen untersuchten Tiefen ein Einbau von ^{13}C -Kohlenstoff in die Fettsäuren erfolgte. Es konnte daher nachgewiesen werden, dass über den gesamten Bereich des untersuchten Vertikalprofils Benzolabbau stattfindet. Wie aufgrund der unterschiedlichen Intensität der Markierung abgeleitet werden kann, scheint der mikrobielle Benzolabbau offensichtlich in den Bereichen der maximalen Schadstoffbelastungen geringer als in den Randbereichen der Fahne zu sein. Da mit den BACTRAPS der mikrobielle Benzolabbau auch in stark kontaminierten Zonen nachgewiesen werden konnte, ist die Sensitivität dieses Testsystems wesentlich höher als beim Nachweis der Biodegradation über die Methode der Isotopenfraktionierung.

Die Quantifizierung des Benzolabbaus und Erstellung von Stoffumsatzbilanzen mit der Methode der BACTRAPS gestaltet sich problematisch. BACTRAPS verlieren unter vergleichbaren Bedingungen zu diesem Experiment etwa 80 bis 85 % der Menge des beaufschlagten Substrats (Benzol bzw. Toluol) (Geyer et al., 2005). Ein Teil des Substrats wird im Wasser gelöst und mit dem Grundwasserstrom abtransportiert. Dieser Verlust ist nur sehr schwer abschätzbar. Qualitativ konnte in anderen Multilevel-Versuchen eine Desorption des ^{13}C -markierten Substrats aus den BACTRAPS nachgewiesen werden. BACTRAPS mit nicht markierten Substraten, die in ca. 25 cm Entfernung zu ^{13}C -markierten BACTRAPS exponiert waren bzw. Grundwasserproben aus ähnlicher Tiefe zeigten eine ^{13}C -Anreicherung (unveröffentlichte Ergebnisse). Zukünftig sind Referenzversuche mit inerten, nicht abbaubaren Substraten zur Bestimmung der Desorptionskinetik, zur Berechnung des den Mikroorganismen tatsächlichen zur Verfügung stehenden markierten Substrats sowie zur Abschätzung von Verlusten nötig.

Ein weiterer Teil des Substrats wird offensichtlich mineralisiert und als CO_2 bzw. Methan aus dem BACTRAP in den Grundwasserstrom abgegeben und steht somit zur Quantifizierung einer geschlossenen Massenbilanz nicht zur Verfügung. Da Mikroorganismen in der Regel einen bestimmten Teil des Substrates, auf dem sie wachsen, zur Synthese eigener Biomasse verwenden, könnten zur Abschätzung der Schadstofftransformation

Ertragskoeffizienten dienen. Ertragskoeffizienten beschreiben das Verhältnis zwischen Mineralisierung und Bildung von Biomasse und könnten in geschlossenen Modellsystemen bestimmt werden. *Ex situ* Referenzversuche im Labor sind allerdings schwierig, da anaerob Benzol abbauende Mikroorganismen gegenwärtig nur schwer kultivierbar sind. Alternativ könnte eine Abschätzung der Ertragskoeffizienten für die verschiedenen terminalen Elektronenakzeptoren mittels Referenzkulturen in Versuchen mit leichter abbaubaren Substraten wie Toluol oder Acetat bestimmt werden, die dann zunächst als Richtwerte für verschiedene biogeochemische Bedingungen zur Verfügung stehen.

Zukünftig soll anhand der Isotopensignatur der Anteil des in die Biomasse transformierten Substrats quantifiziert werden, so dass eine Abschätzung der Mineralisierung perspektivisch möglich scheint. Methoden zur Extraktion der gesamten Biomasse aus den *in situ* Mikrokosmen werden gegenwärtig entwickelt.

BACTRAPs stellen ein bedeutendes Testsystem zum direkten Nachweis der im Grundwasserleiter stattfindenden Biodegradation dar. Der entscheidende Vorteil des Testsystems ist, dass der Nachweis des mikrobiellen Schadstoffabbaus unter realen Aquiferbedingungen innerhalb nur weniger Monate erfolgen kann. Ein Nachteil zu Laborversuchen besteht derzeit in der eingeschränkten Möglichkeit zur Bilanzierung des Abbaus. Laborsysteme können geschlossene Stoffbilanzen einfach erfassen. Das ist, wie bereits dargestellt, mit *in situ* Mikrokosmen zurzeit nicht möglich. Da Testverfahren für Sedimente und Böden im Labor in der Regel nicht die Komplexität der Umweltbedingungen reproduzieren können, ist eine Übertragung der quantitativen Laborergebnisse auf den Aquifer ein kontrovers diskutiertes, ungelöstes Problem. Außerdem erfolgt der anaerobe Abbau meist sehr langsam, weshalb Laborverfahren häufig relativ aufwendig und zeitintensiv sind. Deshalb besitzen im Vergleich zu diesen herkömmlichen Systemen sowohl die Untersuchung der Isotopenfraktionierung als auch *in situ* Mikrokosmen mit isotopisch markierten Substanzen ein erhebliches Potenzial zur Charakterisierung des mikrobiellen *in situ* Schadstoffabbaus in kontaminierten Grundwasserleitern.

2.7 Zusammenfassung

Eine tiefenorientierte Untersuchung von Kontaminationsfahnen erlaubt die Analyse hydrogeochemischer Gradienten im Aquifer. Konzentrationen von Elektronenakzeptoren geben Aufschluss über vorherrschende biogeochemische Prozesse. Die vertikale Analyse der Kontaminationsfahne anhand der Konzentration und Isotopensignatur von Kontaminanten liefert qualitative und quantitative Indikatoren für den *in situ* Schadstoffabbau. Anhand der Methode der Isotopenfraktionierung konnten Abbauprozesse an den vertikalen Fahnenrändern des untersuchten Tiefenprofils analysiert werden. Eine messbare Veränderung der Isotopensignatur kann meist erst nach einem relativ hohen

Analyse des anaeroben Benzolabbaus:
Vergleich von In situ Mikrokosmen, Elektronenakzeptorbilanzen und
Isotopenfraktionierungsprozessen

Schadstoffumsatz nachgewiesen werden. Bei geringem Abbau ist die Isotopenmethode relativ unempfindlich. Zusätzliche Untersuchungen mit den BACTRAPS konnten hier weiterhelfen, da mit dieser Methode der Schadstoffabbau mit einer höheren Empfindlichkeit nachgewiesen werden konnte. Mit der Isotopenmethode ist eine quantitative Abschätzung des Abbaus auf dem Fließweg zwischen Quelle und Messpunkt möglich. Der Einsatz von *in situ* Mikrokosmen mit isotopisch markierten Substanzen kann als ein empfindliches Testsystem für den eindeutigen Nachweis des Abbaus genutzt werden. Es werden qualitative Informationen zum Abbaupotenzial in Bereich der untersuchten Messstelle gewonnen. Verglichen mit Abbauproben im Labor geben *in situ* Mikrokosmen direkte Aussagen zur Biodegradation im Grundwasserleiter. Schwierigkeiten bestehen derzeit noch in der quantitativen Interpretation der Daten. Die vertikale Analyse von geochemischen Gradienten und Isotopensignaturen von Schadstoffen, DIC sowie Methan verbunden mit dem Einsatz von *in situ* Mikrokosmen liefert Informationen zu biogeochemischen Abbauprozessen, die für einen Schadensfall relevant sind. Der Einsatz von MLPS in Verbindung mit modernen geochemischen und isotopenchemischen Verfahren ermöglicht somit mit vertretbarem Aufwand die Gewinnung entscheidender Informationen für die Planung von *Natural Attenuation*-Vorhaben bzw. für die Überwachung von aktiven Sanierungs- bzw. *Enhanced Natural Attenuation*-Verfahren.

2.8 Danksagung

Die Bio-Sep®-Beads wurden von K.L. Sublette (Universität Tulsa, USA) für die Experimente zur Verfügung gestellt. Wir bedanken uns bei U. Günther und M. Gehre (Department für Isotopenbiogeochemie, UFZ) für ihre Hilfe während der Isotopenanalysen, bei R. Trabitzsch und H. Weiß (Department für Grundwassersanierung, UFZ) für ihre Unterstützung bei der Planung und Koordination der Feldarbeiten und bei H. Weiß (IMW, Tübingen) für die Hilfe bei der Installation des MLPS. Das Vorhaben wurde finanziell durch den Projektbereich SAFIRA des UFZ unterstützt. N. Stelzer wird von der Deutschen Bundesstiftung Umwelt (DBU) durch ein Promotionsstipendium (AZ 20004/751) finanziell gefördert.

3 In situ microcosms to evaluate natural attenuation potentials in contaminated aquifers

Nicole Stelzer, Christian Büning, Frank Pfeifer, Anja B. Dohrmann, Christoph C. Tebbe, Ivonne Nijenhuis, Matthias Kästner, Hans-H. Richnow

(This chapter was published in Organic Geochemistry 37(10), p. 1394-1410)

3.1 Abstract

In situ microcosm (BACTRAP[®]) experiments with ¹³C-labelled toluene and benzene were conducted to investigate the *in situ* biodegradation in contaminated aquifers at field sites with different geological and geochemical conditions. The transformation of the carbon (¹³C), derived from labelled substrates, into fatty acids verified the assimilation of the pollutants with formation of biomass. This clearly demonstrated that the *in situ* microcosm system is a useful culture-independent method to investigate *in situ* biodegradation in the aquifer. In addition, metabolites such as benzylsuccinic acid were found on toluene amended BACTRAPS indicating that toluene was degraded anaerobically. This result corresponded to the geochemical conditions found at the field site and the approach enabled to analyse the metabolic pathway governing *in situ* toluene biodegradation in the polluted aquifer.

Phospholipid fatty acids (PLFA) of living cells make up a significant part of the total lipid fatty acid (TLFA) fraction. Comparing BACTRAPS exposed at different geochemical zones of the aquifers, the fatty acid composition was found to be relatively similar indicating that the composition of the TLFA was of low taxonomic value and not sensitive enough for a community analysis. Therefore the composition of the microbial communities were analysed by genetic profiling and sequencing of partial 16S rRNA genes PCR-amplified from total DNA, extracted directly from the BACTRAPS. Sequences retrieved from the BACTRAPS indicated a dominance of not-yet cultivated bacteria, with several of them phylogenetically closely related to those with an iron and sulphate reducing capacity, typically found at BTEX and mineral oil polluted sites.

3.2 Introduction

The fate of pollutants in contaminated aquifers is mainly governed by microbial processes. Abiotic processes such as dispersion, dilution and volatilisation may contribute to a decrease in concentration but do not lead to a significant reduction of the overall amount of contaminants. Therefore, the evaluation of *in situ* biodegradation is crucial for the implementation of Natural Attenuation (NA) concepts in groundwater management strategies. Conventionally, complex calculations of decreasing contaminant concentrations related to the depletion of electron acceptors have been used to identify biodegradation

processes. However, this method is difficult to apply in order to evaluate the fate of a single pollutant in contaminant mixtures on the field scale, particularly in the very heterogeneous natural environment with several competing electron acceptor-donor interactions. Except for the stable isotope fractionation technique (Meckenstock et al., 2004a) and relatively labour intensive tracer experiments (Fischer et al., 2005; Reusser and Field, 2002) tools to quantify the actual *in situ* biodegradation of organic contaminants in the aquifer are scarce.

Culture-dependent laboratory microcosm studies are often used to investigate the potential of *in situ* biodegradation. However, this approach is not reliable because the majority of microorganisms have not been cultured yet and even the reproducible cultivation of anaerobic bacteria degrading typical contaminants such as BTEX and PAH is not an easy task. Laboratory microcosm studies as well as percolation column experiments, simulating the contaminant degradation under controlled water flow conditions, are time consuming and the conditions in the laboratory are certainly different to the actual natural environment complicating the interpretation of the results. Several limitations are related to these laboratory approaches particularly in providing reliable data to transfer the results to the actual ongoing *in situ* processes.

Recently, ¹³C-labelled substrates in combination with ¹³C-enrichment of PLFA were used to characterise bacterial toluene degradation in soil, sediment or aquifer microcosms (Hanson et al., 1999; Pelz et al., 2001a; Pelz et al., 2001b) and to trace the assimilation of toluene along a food chain (Mauclaire et al., 2003) providing useful information on carbon fluxes and metabolites. At the field scale, microbial incorporation of ¹³C-labelled acetate into biomarker molecules like PLFA and DNA were successfully used to detect microbes which were suggested to be responsible for the reduction of uranium(VI) (Chang et al., 2005). To monitor the *in situ* dynamics of microbial communities in a BTEX polluted aquifer a mesocosm study was designed (Hendrickx et al., 2005). In this study, uncontaminated aquifer material was incubated either in the uncontaminated area or nearby a contaminated area of the aquifer and it was shown that the bacterial communities were not identical in these two zones using molecular biological techniques.

To overcome the limitations of *ex situ* laboratory test systems and to improve the monitoring of *in situ* biodegradation, an *in situ* microcosm system (BACTRAP[®]) was developed that can be directly incubated within the groundwater monitoring wells (Geyer et al., 2005). The system consists of pellets containing activated carbon that can be loaded with ¹³C-labelled contaminants. The pellets provide large amounts of interfacial area for colonisation or attached growth of degrading bacteria. If the indigenous bacteria colonise the BACTRAPS and consume the contaminants *in situ*, the ¹³C-labelled carbon will be transformed into the biomass and can be traced within biomarker molecules such as fatty acids or nucleic acids.

In a first approach, BACTRAPS containing activated carbon beads were loaded with ^{13}C -labelled benzene and toluene prior to incubation within the anaerobic zone of a contaminated aquifer (Geyer et al., 2005; Stelzer et al., 2006b). The pattern of fatty acids (FA) and the incorporation of ^{13}C into FA provided evidence for the *in situ* degradation of the test substrates and the metabolic transformation within a complex microbial community. In order to enable a more detailed characterisation of the bacterial community involved in biodegradation, we now analysed the microbial community colonising the *in situ* microcosms. For that reason the total DNA was directly extracted from the *in situ* microcosms and molecular tools were applied to study the microbial community structure. Partial sequences of the bacterial 16S rRNA genes, covering approximately 30 % (400 base pairs) of the complete gene, were PCR-amplified from total DNA and the diversity of the amplified product was visualised by genetic profiling, using the single-strand conformation polymorphism technique (SSCP) (Dohrmann and Tebbe, 2004; Schwieger and Tebbe, 1998).

Because the adsorbing material of the BACTRAPS can enrich degradation metabolites we intended to identify the metabolic process and pathway governing the *in situ* biodegradation of toluene in the aquifer.

In this paper we present our recent progress with BACTRAP systems. We compare *in situ* microcosm experiments with different contaminants at different field sites. This *in situ* approach was used to monitor the biodegradation potential within different geochemical zones of contaminated aquifers related to the vertical structure of a contamination plume. In addition, perspectives to use *in situ* microcosm experiments for microbial community structure analysis and for the identification of metabolic pathways are discussed.

3.3 Materials and Methods

3.3.1 Chemicals

The chemicals and solvents were obtained in p.A. quality from Merck unless stated. [$^{13}\text{C}_6$] benzene, [^{13}C]- α -toluene and benzylsuccinate were obtained from Sigma-Aldrich (St. Louis, USA). [$^{13}\text{C}_7$] toluene was purchased from Chemotrade Leipzig (Germany). All the isotopically labelled compounds had a chemical and an isotope purity higher than 99 %.

3.3.2 Field Sites

Zeitz (Saxony-Anhalt, Germany). The benzene, toluene, ethylbenzene and *o,m,p*- xylene (BTEX) contaminated aquifer is located in the area of a former hydrogenation plant close to the city of Zeitz, Germany (Fig. 3-1). A detailed description of the site hydrogeology and hydrochemical conditions of the contamination plume is given in previous studies (Fischer et al., 2004; Vieth et al., 2005).

The BACTRAP experiments were performed in a multilevel approach in the upper aquifer to investigate different geochemical zones of the vertical BTEX plume. The thickness of the upper aquifer varies between 4 to 6 m. In the source area total BTEX concentrations exceeded 900 mg L^{-1} (Fig. 3-1). Benzene and toluene were present in concentrations up to 850 mg L^{-1} and 50 mg L^{-1} , respectively. Ethylbenzene and xylenes were typically present in concentrations lower than 3 mg L^{-1} .

At the test site the predominant electron acceptor used for biodegradation was sulphate (Vieth et al., 2005). Active sulphate reduction in the course of the plume was evident by an enrichment of ^{34}S -sulphate (Fischer et al., 2004). In addition, methanogenesis had a minor impact on microbial BTEX-transformation in the BTEX-plume but a more significant impact in the source area (Fischer et al., 2005). Other electron acceptors like oxygen, nitrate and iron play a minor role for the overall biodegradation processes at the site (Dethlefsen et al., 2004; Fischer et al., 2004; Vieth et al., 2005).




Fig. 3-1: Concentration of BTEX [$\mu\text{g L}^{-1}$] in the upper aquifer in the area of the former hydrogenation plant in Zeitz (Saxony-Anhalt, Germany). The locations of the *in situ* microcosm experiments are indicated. At well 23/00 the multilevel experiment was carried out and well 32/00 was used for the *in situ* microcosm study by Geyer et al. (2005). The arrow depicts the general groundwater flow direction.

Gneisenau (Dortmund, Germany). The anaerobic tar oil and BTEX contaminated aquifer is located at the former coking plant Gneisenau in the vicinity of a coal mine in the Ruhr area. The hydrological situation is characterised by two aquifers separated by erratic marls and tills. The hydraulic conductivity of the upper porous quaternary aquifer is relatively low at

the basis (k_f of about 10^{-9} m s^{-1}) but is significantly higher in sandy layers ($k_f = 10^{-6}$ to 10^{-4} m s^{-1}). The water table is located 5 to 20 m below ground. The lower aquifer is a confined fractured cretaceous aquifer. Both aquifers are contaminated by BTEX, PAH and other contaminants typical for coking plants. The upper quaternary aquifer was mainly contaminated by BTEX (up to 1.5 mg L^{-1}) and naphthalene (up to $750 \mu\text{g L}^{-1}$), whereas in the deeper cretaceous aquifer mainly benzene (up to 45 mg L^{-1}) was found. Oxygen concentrations were always below 1 mg L^{-1} . The isotope enrichment of ^{34}S -sulphate in most contaminated areas provided evidence for sulphate reduction (Pfeifer, 2004). Sulphate, with concentrations up to 1660 mg L^{-1} , is likely the most important terminal electron acceptor at this site. Geochemical investigations suggested that methanogenesis as well as nitrate, iron and manganese reduction are probably less important for microbial processes in the subsurface (Pfeifer et al., 2003). The presence of benzylsuccinic acid, a typical metabolite of toluene, provided further evidence that anaerobic degradation processes are predominant at the site (Pfeifer, 2004).

3.3.3 Multi level packer systems used for depth discrete experiments

A multi level packer system (MLPS) was used for depth discrete sampling in Zeitz. The multi level packer system is a water-filled sock tube inserted into the well, which allows separating different sampling systems from each other within different depth. Small submersible pumps allowed depth-specific sampling without cross-currents. A detailed description of the multilevel sampling system is given by Schirmer et al. 1995 (Schirmer et al., 1995).

The depth discrete investigation of geochemical and isotope parameters in Zeitz at well 23/00 were previously described by Stelzer et al. (2006b). The position of BACTRAP systems, water table and geological information of well 23/00 are shown in Fig. 3-1 and Fig. 3-2.

3.3.4 Preparation and incubation of *in situ* microcosms (BACTRAP[®])

Bio-Sep[®] beads (K. Sublette, University of Tulsa, Tulsa, USA) were loaded with contaminants as carbon substrate for microorganisms. The spherical beads, 2 to 3 mm in diameter, consist of powdered activated carbon (PAC) incorporated within an aramid polymer matrix (Nomex[®]). The beads have an internal porosity of 75%, an internal surface area greater than $600 \text{ m}^2 \text{ g}^{-1}$, and outer pores of 1-10 microns (Peacock et al., 2004; White et al., 2003). The beads were heated at 300°C for 4 hours to remove organic residues and then 0.2 to 0.4 g of beads were filled in perforated Teflon[®] tubes. Glass wool was used as a plug to keep the beads inside the Teflon[®] tubes. The filled Teflon[®] tubes were autoclaved at 121°C for sterilization and rehydration of the beads. The microcosms were loaded with the substrate (benzene or toluene) via gas phase under reduced pressure as described previously (Geyer et al., 2005). The substrate and the microcosms were placed in small

glass containers which were evacuated at 60 mbar. To allow uniform adsorption of the contaminants the Bio-Sep® beads were incubated for at least 48 hours. Contrary to the previous method, the vacuum was then released by filling oxygen free water into the glass container and the BACTRAPs were stored under anoxic water until deployment in the monitoring wells. The oxygen free water was used to fill the free pore space of the beads due to the vacuum release to keep the *in situ* microcosm system anoxic. The BACTRAPs for the multi level experiment in Zeitz were loaded with [¹³C₆] benzene to a concentration of about 76 mg g⁻¹ and for the metabolite experiment with [¹³C]-α toluene to a concentration of about 50 mg g⁻¹. The BACTRAPs deployed in Gneisenau were loaded with uniformly labelled [¹³C₆] benzene and [¹³C₇] toluene to a concentration of 500 mg g⁻¹. In all experiments, BACTRAPs were loaded with non labelled benzene or toluene in similar concentration for control experiments. Material from these control experiments was used for molecular biological studies and to investigate the isotope composition of fatty acids at natural abundance.




Fig. 3-2: Left: Geological profile at monitoring well 23/00 (Zeitz, Germany). The location of the filter screen is illustrated. The groundwater table was located at 9.05 m below ground. *In situ* microcosms were installed at A, C, D, F, G, H and samples for geochemical analyses were taken from all depths except A. Right: The picture shows the multi level sampling system with stainless steel containers for *in situ* microcosms before subjection into the well.

The microcosms were deployed with the MLPS or in monitoring wells over a time period of 51 to 100 days. In the MLPS experiments the microcosms were placed in stainless steel cages as a spacer to ensure a permanent flow of groundwater through the *in situ*

microcosms and to avoid contamination with biological material by direct contact with the packer membrane or the filter screen of the well.

3.3.5 Extraction and Derivatisation of Fatty Acids

The Bio-Sep® beads were extracted using a dichloromethane-methanol-water mixture as solvent modified from Bligh and Dyer 1959 (Bligh and Dyer, 1959). After phase separation the dichloromethane phase containing the fatty acids (FA) was evaporated to dryness and derivatised using a trimethylchlorosilane (TMCS) methanol mixture (1:8; v:v) as reactant for 2 h at 70 °C to obtain fatty acid methyl esters (FAME) (Thiel et al., 2001). After evaporation to complete dryness the FAME fraction was dissolved in *n*-hexane for subsequent analysis by gas chromatography-mass spectrometry (GC-MS) and gas chromatography-combustion-isotope-ratio-monitoring-mass-spectrometer system (GC-C-IRMS).

3.3.6 Analysis

GC-MS. For identification and structural characterisation by GC-MS a Hewlett Packard 6890 gas chromatograph coupled to a 5973 quadrupole mass spectrometer (Agilent, Palo Alto, USA) was used. The carboxylic acid fraction was separated on a BPX-5 column (30 m*0.32 mm*0.25 µm) (SGE, Darmstadt, Germany) with a temperature program of 120 °C initial temperature for 4 min, heat at 4 °C min⁻¹ to 250 °C, heat at 20 °C min⁻¹ to 300 °C, and hold for 10 min. FAME were identified by co-injection of an authentic standard mix (bacterial acid methyl ester mix, Sigma-Aldrich, Germany). The fatty acids are designated in the form of A:B_ωC where A is the number of carbon atoms, B is the number of double bonds and C is the distance of the closest double bond from the aliphatic end of the molecule (unsaturation, $ω$ -nomenclature). The prefix *i* (*iso*) and *a* (*anteiso*) refer to methyl branching. The benzylsuccinic acid methyl ester was characterised by co-injection and comparison of mass spectra of the authentic reference compound which was derivatised as described above.

GC-C-IRMS. The carbon isotope composition of the carboxylic acids fractions was analysed using a GC-C-IRMS. The system consists of a gas chromatograph (6890 Series, Agilent Technology, Palo Alto, USA) coupled via a Conflow III interface (ThermoFinnigan, Bremen, Germany) to a MAT 252 mass spectrometer (ThermoFinnigan, Germany) as described previously (Richnow et al., 2003a). A BPX-5 column (50 m*0.32 mm*0.5) (SGE, Darmstadt, Germany) was used for chromatographic separation of fatty acid methyl esters with helium as carrier gas at a flow rate of 1.5 mL min⁻¹ and a temperature program with initial temperature of 60 °C for 2 min, heat at 20 °C min⁻¹ to 120 °C, heat at 2 °C min⁻¹ to 300 °C, and hold for 20 min (Miltner et al., 2004). The analysis of BTEX (Vieth et al., 2005) on the same system was described previously.

The carbon isotope ratio of fatty acids is reported in δ-notation (per mill) relative to the Vienna Pee Dee Belemnite standard (V-PDB) with known isotopic composition (Anonymus,

1995). The isotope composition of highly ^{13}C -enriched fatty acids could not be determined accurately. The instrument was calibrated to about 1000 δ units. Higher values are outside the calibration range of the instrument and the uncertainty increases with higher enrichments. Therefore, positive isotope compositions of fatty acids were only reported with the first 2 decimal places of the value.

3.3.7 Molecular biological analysis

For each sample, a total of 50 Bio-Sep® beads were transferred into 15 ml Falcon tubes containing 12 ml of TES buffer (50 mM NaCl₂, 10 mM Na₂EDTA, 50 mM Tris (hydroxymethyl) aminomethane hydrochloride [Tris HCl], pH 8.0) with 1% sodium dodecylsulphate (SDS). Five cycles of freeze-thaw lysis were performed, each with 5 min in liquid nitrogen (-196 °C) and 5 min at 65 °C in a shaking water bath, each cycle was interrupted by 10 s of vigorous vortexing. DNA was then extracted with phenol-chloroform, as described elsewhere (Schwieger and Tebbe, 1998; Schwieger and Tebbe, 2000). The isopropanol precipitated and purified DNA of each tube was resuspended in a total of 40 µl of 10 mM Tris in 10 µl aliquots. Aliquots were kept at 4 °C for further analysis or stored at -20 °C.

Partial 16S rRNA genes were amplified from total DNA by PCR, using primers Com1 and Com2ph (Schmalenberger et al., 2001). The PCR products were converted to single strands by lambda-exonuclease digestion, following the protocol of Schwieger and Tebbe (1998). Genetic profiles of each sample were generated by single strand conformation polymorphism (SSCP) on non-denaturing polyacrylamide gels, as described in detail elsewhere (Dohrmann and Tebbe, 2004; Tebbe et al., 2001). Single bands were selected for further DNA-sequencing after PCR-amplification and cloning in *Escherichia coli*. Sequencing is described in the same references (Dohrmann and Tebbe, 2004; Peters et al., 2000; Tebbe et al., 2001). Consensus sequences were compared to database sequences using the FASTA tool provided by the European Bioinformatics Institute (<http://www.ebi.ac.uk/fasta/>). DNA-sequences of this study have been deposited in the GenBank database under the accession numbers AJ123456 to AJ123478.

3.4 Results and Discussion

3.4.1 Zeitz

In order to test the BACTRAP concept, the BTEX contaminated aquifer in Zeitz (Germany) was chosen as a model site, because detailed information was available about the contamination plume and the geochemical zones from previous studies (Fischer et al., 2005; Fischer et al., 2004; Vieth et al., 2001; Vieth et al., 2005). At well 32/02 (Fig. 3-1), located downstream of the contamination source and characterised by moderate benzene concentrations of approximately 44 mg L⁻¹ in 2002, *in situ* microcosms supplemented with

¹³C-labelled toluene and benzene provided evidence for the *in situ* degradation of these substances, as described in detail elsewhere (Geyer et al., 2005).

For further BACTRAP studies we selected well 23/00 (Fig. 3-1, Fig. 3-2) because of generally higher concentrations of contaminants and clear indication of microbial sulphate reduction as the predominant terminal electron accepting process by sulphur isotope analysis (Fischer et al., 2004).

A depth specific MLPS approach was chosen to investigate different geochemical zones of the aquifer within only one monitoring well (Fig. 3-2). The benzene plume showed a vertical structure at this monitoring well (Fig. 3-3). The benzene concentration increased from 415 $\mu\text{mol L}^{-1}$ at 10.4 m below the surface, which was located near the water table at 9.05 m, to a maximum of more than 6000 $\mu\text{mol L}^{-1}$ in the depth interval between 13 to 14 m. Towards the aquitard, located at about 17 m, the benzene concentration decreased to about 1700 $\mu\text{mol L}^{-1}$ at 16.7 m depth.




Fig. 3-3: Left: Concentration of sulphate (\circ), nitrate (*), sulphide (\blacktriangle), ferrous iron (\blacksquare), ferric iron (\square) and manganese (\times) in $[\mu\text{mol L}^{-1}]$. Right: benzene (\blacksquare) and methane (\bullet) concentration in $[\mu\text{mol L}^{-1}]$ at Zeitz (Germany) monitoring well 23/00.

In uncontaminated zones of the aquifer upstream the plume, the concentrations of nitrate were relatively low (< 100 μmol) compared to sulphate concentrations (> 10 mmol) suggesting that nitrate was not a dominant electron acceptor in this aquifer. At the observation well manganese and nitrate were only present in the upper zone of the aquifer in elevated concentrations (Fig. 3-3). The low total manganese concentrations indicated that manganese mobilisation by Mn(IV) reduction was not an important electron accepting

process at the field site. Thus nitrate and manganese may act as potential electron acceptors at the water table close to the capillary fringe, where also some oxygen may enter from the gaseous phase of the soil into the aquifer system. Ferrous iron mobilisation suggested that ferric iron was a potential electron acceptor. However, the concentration of ferrous iron may not show the true extent of iron reduction, because ferrous iron can be precipitated in the presence of H_2S . However, as discussed elsewhere iron reduction is not an important terminal electron accepting process in this aquifer (Dethlefsen et al., 2004). Sulphate concentrations $> 10 \text{ mmol L}^{-1}$ are typical for uncontaminated parts of the aquifer (Fischer et al., 2004) and in the upper part of the vertical profile sulphate concentrations up to 5 mmol L^{-1} were found. The depletion of sulphate within the plume to concentrations of less than 0.2 mmol L^{-1} between 11.5 and 14.5 m suggested intensive microbial sulphate reduction although the development of sulphide was low. Sulphide may be precipitated by iron, which emphasises the difficulties to trace the degradation of contaminants by electron donor acceptor balances. Thus, sulphate was presumably the predominant electron acceptor.

Fatty acid composition. The fatty acid patterns extracted from the BACTRAPS were dominated by hexadecanoic (C16:0) and octadecanoic (C18:0) acids. Unsaturated hexadecenoic (C16:1) and octadecenoic (C18:1) acids were present in lower concentrations (Fig. 3-4). Other linear saturated fatty acids like tetra (C14:0), penta- (C15:0) and heptadecanoic (C17:0) acids as well as *iso* and *anteiso* branched fatty acids with 15 carbon atoms were present in much lower concentrations.




Fig. 3-4: The composition of the total fatty acid fraction (m/z 74) obtained from *in situ* microcosm experiment deployed at various depths (A to H) in well 23/00 at the field site Zeitz (Germany). The depths of the microcosm (A to H) and the position in the aquifer are shown in Fig. 3-2.

The *iso* and *anteiso* branched fatty acids are used as biomarkers for Gram-positive bacteria suggesting their involvement in biodegradation in some zones of the aquifer (Kaur et al., 2005). The fatty acid patterns of all depths were very similar although geochemical

conditions were changing, suggesting that the fatty acid pattern did not respond sensitively to the geochemical conditions. However, the composition of fatty acids found in benzene amended *in situ* microcosms at marginal part of the plume was more diverse suggesting a more diverse microbial community (Geyer et al., 2005). Further biomarker fatty acids specifically characterising some individual sulphate reducing bacterial strains were not observed (Kaur et al., 2005). In our studies we could not use the fatty acid pattern for a taxonomic interpretation.

Isotope signature of fatty acids. The isotope compositions of fatty acids ($\delta^{13}\text{C}_{\text{FA}}$) extracted from the BACTRAPS amended with [$^{13}\text{C}_6$] benzene were between -28 ‰ and 439 ‰ (PDB) (Tab. 3-1). The variation in isotope composition may indicate the presence of organisms feeding on various carbon sources. Fatty acids with an isotope signature lower than -25 ‰ show the typical natural abundance of $\delta^{13}\text{C}_{\text{FA}}$ found in soil and aquifer materials (Pelz et al., 2001a; Pelz et al., 2001b) and are typical for organisms using natural occurring non labelled carbon sources. In BACTRAP experiments with non labelled substrate or without any substrate supply via Bio-Sep® beads all fatty acids showed natural abundances (-35 to -20 ‰, data not shown). Fatty acids with a positive carbon isotope signature (>0 ‰) clearly indicated that microorganisms incorporated the labelled carbon from the provided [$^{13}\text{C}_6$] benzene (Tab. 3-1). The enrichment of ^{13}C in fatty acids can only stem from the labelled substrate and thus clearly provided evidence of *in situ* benzene degradation.

Tab. 3-1: Carbon isotope composition of fatty acids extracted from *in situ* microcosms incubated with $^{13}\text{C}_6$ labelled benzene at monitoring well 23/00 (Zeitz, Germany) for 6 weeks. A to H refers to incubation depth indicated in Fig. 3-2.

depth	$\delta^{13}\text{C}_{\text{FAME}} [\text{\textperthousand}]$							
	C14:0	i-C15:0	a-C15:0	C15:0	C16:1	C16:0	C18:1	C18:0
A	63	89	-22	-18	330	62	22	-21
C	-8	90	-19	-22	439	47	61	-15
D	-20	-17	-28	-23	35	-13	-21	-24
F	-17	1	-23	-23	86	15	-2	-25
G	-22	134	-23	-20	174	20	-6	-24
H	-22	-13	-22	-26	24	-3	-16	-26

BACTRAPS with ^{13}C -labelled benzene incubated in different geochemical zones showed variations in the ^{13}C -enrichment of fatty acids within the various depths although the composition of fatty acids did not change significantly. Hexadecenoic acid (C16:1) showed the highest incorporation of ^{13}C (439 ‰) next to i-C15 (134 ‰) and C16:0 (62 ‰). a-C15, C15:0, C14:0 and C18:0 displayed only low or even no incorporation of ^{13}C . The i-C15 generally showed a significantly higher labelling than a-C15 which was almost not labelled. The variation in $\delta^{13}\text{C}_{\text{FA}}$ clearly indicated that the organisms producing these FA grew on different carbon sources. Fatty acids displaying a higher incorporation of ^{13}C were very likely

produced by organism feeding on [$^{13}\text{C}_6$] benzene while fatty acids from bacteria with lower or no ^{13}C incorporation were likely not involved in the degradation of [$^{13}\text{C}_6$] benzene. Cross feeding by metabolites may channel labelled carbon into individual members of the microbial community.

In the upper zone of the aquifer the ^{13}C incorporation was higher, which may suggest a more intensive transformation of the [$^{13}\text{C}_6$] benzene. This might be a result of more attractive electron acceptors for degradation such as nitrate, manganese and iron as well as oxygen close to the water table. Nevertheless, also in the depth between 12 and 14 m indications for benzene degradation were found despite high benzene concentrations. Here sulphate reducing conditions were likely present.

On the other hand, labelling of biomass can be influenced by the availability of labelled benzene on the BACTRAPS relative to the unlabelled benzene from the groundwater itself. The concentration of benzene in the upper part of the plume was lower compared to deeper zones of the aquifer (Fig. 3-3). Therefore, benzene derived from the contamination might be available as a carbon source to a relatively higher amount in the depth of 12 and 14 m with benzene concentration up to 6 mmol L $^{-1}$ (Fig. 3-3). The relative intensity of ^{13}C incorporation into biomass changed more pronounced with depth than the composition of the fatty acid fraction and might therefore reflect the relative intensity of benzene metabolism and geochemical conditions. Compared to previous investigations with [$^{13}\text{C}_6$] benzene amended *in situ* microcosms at marginal parts of the plume where fatty acids were labelled up to 13000 % (Geyer et al., 2005), the ^{13}C incorporation found in the multi level investigation was significantly lower (max. 439 %) suggesting a less intensive metabolism at the different depth of this monitoring well.

3.4.2 Gneisenau: Upper quaternary aquifer

The compositions of the total fatty acid fractions obtained from *in situ* microcosms loaded with [$^{13}\text{C}_6$] benzene and [$^{13}\text{C}_7$] toluene were compared in order to investigate the microbial communities growing on these different substrates. The *in situ* microcosms were exposed to the upper quaternary aquifer Gneisenau, Germany (P10F) over 12 weeks and analysed for composition and isotope signatures of fatty acids. The patterns of fatty acids derived from toluene and benzene amended BACTRAPS incubated in the source zone of the Quaternary aquifer (P10F) were almost identical (Fig. 3-5). C18:0, C18:1 and C16:0 were dominating the fatty acid patterns. C18:1, iso and anteiso branched C15 compounds as well as C15:0 and C14:0 were present in minor concentrations. The similarity in fatty acid compositions of both the benzene and toluene amended BACTRAPS (Fig. 3-5) indicated that the substrate did not affect the pattern of fatty acids suggesting that the lipid pattern is not a sensitive indicator to determine changes in the microbial community.

However, the isotope composition of the fatty acids from the benzene amended microcosm incubated in well P10F showed drastic differences in comparison to the toluene loaded one (Tab. 3-2). Only the hexadecanoic acid (C16:0) of the [$^{13}\text{C}_6$] benzene amended BACTRAP were found to be labelled. Unsaturated C18:1 and C18:2 isomers were only slightly or almost not labelled. The concentrations of other fatty acids were too low for the determination of their isotope composition (Fig. 3-5, Tab. 3-2). In contrast, the [$^{13}\text{C}_7$] toluene amended BACTRAP of well P10F showed a very strong incorporation of ^{13}C into fatty acids with carbon numbers between C14 to C18 of which hexadecenoic acid (C16:1) was most intensively labelled (45000 ‰). Interestingly the fatty acids with 18 carbon atoms were labelled two orders of magnitude lower. The significant high labelling of fatty acids from the toluene amended BACTRAP demonstrated extensive transformation of toluene by some community members whereas benzene carbon was only assimilated to a limited extent. Obviously, other members of the community, producing the significantly lower labelled fatty acids, used other carbon sources in addition to the labelled contaminants.




Fig. 3-5: Comparison of the composition of the total fatty acid fraction (m/z 74) obtained from *in situ* microcosms loaded with [$^{13}\text{C}_6$] benzene (A) and [$^{13}\text{C}_7$] toluene (B). The *in situ* microcosms were both exposed in the upper quaternary aquifer (P10F) at the field site Gneisenau (Germany) over 12 weeks.

3.4.3 Gneisenau: Lower cretaceous aquifer

In situ microcosms were also applied to evaluate the biodegradation of benzene and toluene in the deeper fractured cretaceous aquifer. This aquifer was a hundred times higher contaminated with benzene (up to 45 mg L^{-1}) than the upper quaternary aquifer. BACTRAPS

In situ microcosms to evaluate natural attenuation potentials in contaminated aquifers

loaded with [$^{13}\text{C}_6$] benzene, [$^{13}\text{C}_7$] toluene and with non labelled substrates were exposed up stream the source zone (P8T) and in the centre of the plume (P11T). A higher enrichment of ^{13}C in fatty acids, e.g. in C16:1 (96000‰) and C16:0 (9700 ‰), was only observed in the toluene amended microcosms up stream the source area (P8T) in relatively uncontaminated zones of the deeper fractured aquifer (Tab. 3-2). The fatty acids of benzene or toluene amended BACTRAPs from the source zone (P11T) in the deeper aquifer were only slightly labelled in both cases (Tab. 3-2). The fatty acids occurring in relatively higher concentrations such as C16:0, unsaturated C18 isomers and C18:0 were almost not labelled in comparison to fatty acids commonly present in lower concentrations such as C14-, C15-species and C16:1 (Tab. 3-2). This suggests that only a minor part of the community was involved in the contaminant degradation.

The results show that the transformation of contaminant derived carbon into fatty acids was higher in toluene compared to benzene amended BACTRAPs, which is in accordance to earlier findings that toluene is more easily degradable under anaerobic conditions than benzene (Cunningham et al., 2001; Da Silva and Alvarez, 2004; Wiedemeier et al., 1999). In a direct comparison of parallel experiments with labelled toluene and benzene, the amount of ^{13}C transferred into fatty acids may be used as a direct indicator for the *in situ* degradability of the substrates. The highest enrichment of ^{13}C into fatty acids was observed in BACTRAPs loaded with labelled toluene indicating a higher degradation capacity for toluene compared to benzene. As the *in situ* microcosms were exposed in an almost uncontaminated zone of the deeper aquifer a significant potential to degrade the contaminants was present even in almost non-contaminated zones.

Tab. 3-2: Carbon isotope composition of fatty acids extracted from *in situ* microcosms incubated with $^{13}\text{C}_6$ labelled benzene and $^{13}\text{C}_7$ labelled toluene in the upper Quaternary monitoring well (P10F) and in the lower Cretaceous monitoring wells (P8T, P11T) at the field site Gneisenau (Germany) for 12 weeks.

well Fatty Acid	$\delta^{13}\text{C}_{\text{FAME}} [\text{\textperthousand}]$					
	P8T ^{13}C -Toluene	P10F ^{13}C -Benzene	P10F ^{13}C -Toluene	P11T ^{13}C -Benzene	P11T ^{13}C -Toluene	
14:0	n.d.	n.d.	2800	n.d.	390	
i-15:0	41000	n.d.	14000	n.d.	30	
a-15:0	n.d.	n.d.	1800	n.d.	-21	
15:0	n.d.	n.d.	580	172	n.d.	
i-16:0	n.d.	n.d.	7600	n.d.	-42	
16:1	96000	n.d.	46000	n.d.	140	
16:0	9700	9	3600	-9	-18	
18:2-18:1-Cluster	400	-17	540	-16	-19	
18:0	3	-27	290	-30	-26	

n.d. not determined due to low concentration

To sum up the results, Bio-Sep® beads spiked with ¹³C BTEX can be used to investigate the *in situ* biodegradation of BTEX compounds semi-quantitatively by the amount of incorporated ¹³C in lipid derived fatty acids. The intensity of ¹³C incorporation into biomass (FA) provides valuable information on microbial diversity and on the relative amount of metabolism. The quantification of biodegradation was not possible in these experiments, but with an absolute quantification of fatty acids and an estimation of the proportion of fatty acids in anaerobic bacteria it may be possible to quantify the incorporation into the biomass. Thus further development is needed to apply the BACTRAP system to quantify *in situ* biodegradation.

3.4.4 Molecular biological analysis of microbial communities on BACTRAPS

In order to characterise the contaminant degrading microbial community in the aquifer the biodiversity of the microbial biomass can be analysed by cultivation independent molecular biological approaches based on PCR-amplified partial 16S rRNA genes from total DNA extracted from groundwater samples (Scow and Hicks, 2005). In the present study, total DNA was extracted from the BACTRAPS and the PCR products amplified from the different samples (wells) were compared to each other by their SSCP profiles (Fig. 3-6). This technique is now widely used in microbial ecology to compare the structural diversity of microbial communities (Dohrmann and Tebbe, 2004). In addition to comparing the structural diversity of the bacterial community from different samples, bands of the SSCP-profiles can be characterised by DNA-sequencing in order to identify individual members of the bacterial community (Dohrmann and Tebbe, 2004).




Fig. 3-6: SSCP Gel of PCR amplified 16S rDNA isolated from BACTRAPS loaded with $^{13}\text{C}_7$ toluene or $^{13}\text{C}_6$ benzene after exposure (84 d) in a contaminated aquifer. The bands (numbers) are sequenced for identification (see Tab. 3-3).

Tab. 3-3: Characterization of rRNA genes isolated from BACTRAPS incubated in the lower (P11T) and upper (P10F) aquifer at the field site Greisenau (Germany).

Band	Clone	Length [bp]	Identity [%]	Closest relative in database (GenBank Accession No.)	Phylogenetic group	Source of sequence or isolate
Upper aquifer (P10F)						
1	P10F-01c	371	99.7	uncultured bacterium (AF534262)	Dehaloproteobacteria	marine and freshwater sediments
1	P10F-01a	368	99.5	uncultured bacterium (DQ404728)	Candidate division OP3	contaminated sediment
2	P10F-02b	371	98.4	uncultured bacterium (AF050590)	Firmicutes	hydrocarbon- and chlorinated-solvent-contaminated aquifer
2	P10F-02b	374	99.5	uncultured bacterium (AY050586)	Dehaloproteobacteria	monochlorobenzene contaminated groundwater
3	P10F-03c	371	97.6	unidentified bacterium (AF056010)	Dehaloproteobacteria	PCB-dechlorinating enrichment culture
4	P10F-04c	369	98.9	uncultured bacterium (AY475201)	Alphaproteobacteria	metal-rich and acidic river water
6	P10F-06b	369	100	<i>Ralstonia picketii</i> (AY741342)	Beta proteobacteria	clinical origin
7	P10F-07a	367	99.7	uncultured bacterium (AJ306737)	Bacteroidetes	1,2-Dichloropropane dechlorinating consortium
8	P10F-08b	369	98.9	uncultured bacterium (AB205680)	Alphaproteobacteria	denitrifying activated sludge
Lower aquifer (P11T)						
1	P11T-01a	368	97.3	uncultured bacterium (DQ404728)	Candidate division OP3	contaminated sediment
1	P11T-01b	371	99.4	<i>Geobacter pelophilus</i> (JG6918)	Dehaloproteobacteria	dissimilatory Fe(II)-reducing bacteria
2	P11T-02a	371	98.7	uncultured bacterium (AY945913)	Beta proteobacteria	quinoline-removal bioreactor
3	P11T-03a	371	99.2	uncultured bacterium (AF104274)	Dehaloproteobacteria	benzene-contaminated aquifer sediments
5	P11T-05b	369	99.2	uncultured bacterium (AB240520)	Beta proteobacteria	rhizosphere biofilm of <i>Phragmites</i> at river
5	P11T-05a	369	100	<i>Ralstonia picketii</i> (AY741342)	Beta proteobacteria	clinical origin
6	P11T-06d	371	98.4	uncultured bacterium (AY250093)	Beta proteobacteria	naphthalene-contaminated sediment
6	P11T-06c	367	97.3	uncultured bacterium (AB240238)	Bacteroidetes	reed bed reactor
7	P11T-07c	369	99.2	uncultured bacterium (AB205680)	Alphaproteobacteria	denitrifying activated sludge
9	P11T-16b	366	92.3	uncultured bacterium (AB205753)	Firmicutes	denitrifying activated sludge
10	P11T-10b	367	99.4	uncultured bacterium (AJ519404)	Bacteroidetes	uranium mining waste pile
13	P11T-14d	371	88.9	uncultured bacterium (UEU81707)	Chlamydiae	anaerobic digester

Most of the SSCP profiles generated in this study were relatively similar to each other, indicating a reproducible colonisation pattern of the bead material independently of the geochemical conditions in the investigated groundwater wells (Fig. 3-6). For most samples analysed, the bacterial community compositions on the benzene and toluene loaded microcosms were only slightly altered in comparison to the non-amended ones (blank). However, the fact that highly labelled microbial fatty acids were extracted from the ¹³C experiments indicated a substrate degradation and incorporation into biomass to a different extent (Tab. 3-2). This may show a coordinated degradation by certain members of the microbial community colonising the *in situ* microcosms and a channelling of the ¹³C carbon into the microbial food web. However, even though some fatty acids were highly labelled, the isotope composition showed that these organisms used also other carbon sources. In the highly contaminated part of the aquifer, benzene and toluene were presumably the major carbon sources but other BTEX compounds and minor contaminants as well as humic substances were certainly present and may be used as carbon sources as well. In addition, some fatty acids in ¹³C experiments were not labelled and indicated that microbes not involved in pollutant degradation were also colonising the bead material.

The comparison of the non-amended (blank) with the loaded BACTRAPs showed that the SSCP- and FA patterns were only slightly different (Fig. 3-5, Fig. 3-6). This reflects the fact that the bead material (activated carbon) of the blanks extracted a certain amount of the contaminant i. e. benzene from the contaminated aquifer leading to similar conditions in the non loaded microcosms compared to the benzene loaded ones. The SSCP- and FA patterns from the toluene loaded BACTRAPs were also only slightly different from the non-amended (blank) and benzene BACTRAPs showing that toluene degraders were members of the colonising community.

Detailed analyses showed that the profiles from the different wells clustered together mostly independent from the loaded substrates (data not shown). This indicated a dominant effect of the environmental conditions of each well on the communities developing on the BACTRAPs and a minor influence of the supplied substrate. Therefore, no signals with respect to new emerging bands showed the colonisation of specific contaminant degraders on the loaded *in situ* microcosms. The intense bands of the patterns from the low contaminated area (P8T) indicated a good growth on the microcosms whereas the generally smaller bands from the wells of the contaminated area (P10F and P11T) showed lower biomass accumulation and may point to a certain toxicity of the benzene in that area (Fig. 3-6). The identified DNA-sequences of the SSCP-profiles were mostly related to sequences from yet uncultivated bacterial species (Tab. 3-3, Fig. 3-7). Phylogenetic analyses revealed a community dominated by iron and sulphate reducing bacteria particularly of the *Geobacter* and *Desulfuromonas* group (Lovley and Phillips, 1988; Zhang et al., 2003). These groups of

organisms are known to be involved in contaminant degradation in anaerobic aquifers. This shows that iron and sulphate reducing microbial communities are likely to be involved in the contaminant degradation and that iron and sulphate were used as terminal electron acceptors at this field site.




Fig. 3-7: Phylogenetic analysis (maximum likelihood method) of PCR-amplified partial 16S rRNA genes from the group Deltaproteobacteria, retrieved from BACTRAPS exposed at the field site Gneisenau. Clones are indicated by their source of isolation (P10F for upper and P11T for lower aquifer) followed by their clone number. * indicates sequences which were found at two subsequent sampling dates in the same well.

The molecular biological data supported the results of the hydrological and geochemical analyses and provided valuable information about the biodiversity of the degrading microbial community. In addition, the BACTRAP approach with labelled contaminants provided an excellent tool for assessment and proving the microbial *in situ* activity in contaminated aquifers. However, the key organisms related to contaminant degradation could not be identified. In future, growth experiments to investigate the kinetic of the colonisation traced by quantitative PCR may elucidate the development of communities and may provide

indications for organisms supported by the test substrate. Provided that the total biomass on BACTRAPS is sufficient, stable isotope probing (SIP) may also be applicable to identify organisms which used the ^{13}C labelled contaminants primarily as a carbon substrate (Lueders et al., 2004; Manefield et al., 2004).

3.4.5 Investigation of metabolic pathways

The extracted carboxylic acid fraction from *in situ* microcosm experiments with [^{13}C]- α -toluene contained ^{13}C labelled metabolites which could be easily identified by GC-C-IRMS analysis, due to the very strong signal caused by the ^{13}C labelled carbon although their absolute concentration was very low (Fig. 3-8). An examination of the labelled peak by GC-MS gave a spectrum indicative for benzylsuccinic acid dimethyl ester (Fig. 3-9).




Fig. 3-8: GC-C-IRMS-chromatogram of the carboxylic acid fraction extracted from a [^{13}C]- α -toluene amended BACTRAP incubated at the field site Zeitz (Germany). The metabolites gave very strong signals in the 45/44 ratio plot (A) although their absolute concentration displayed by the amount of CO_2 (m/z 44) was relatively low (B).

The non labelled analogue was found in the carboxylic acids fraction of the BACTRAPS amended with non labelled toluene. The presence of benzylsuccinate was further confirmed by comparison of the mass spectra as well as the retention time of the derivatized authentic standard. The formation of benzylsuccinate is indicative for the anaerobic toluene degradation pathway which is employed by sulphate and iron reducing microorganisms able to degrade toluene (Heider et al., 1999; Spormann and Widdel, 2000; Widdel and Rabus, 2001). Benzylsuccinate has been used as an indicator for anaerobic toluene degradation in

contaminated aquifers before (Beller, 2000; Griebler et al., 2004b). Therefore, the benzylsuccinic acid extracted from the *in situ* microcosm is a good indicator for the biogeochemical conditions governing the toluene degradation in this particular aquifer.

This clearly shows that BACTRAPS can also be useful to identify degradation pathways in the aquifer by analysing the potential metabolites. Reusser and Field (2002) applied push and pull experiments with deuterium labelled toluene to verify *in situ* toluene degradation in a contaminated aquifer. Although deuterium labelled contaminants may be useful tracer compounds to provide evidence for *in situ* degradation, much higher amounts of labelled substances are needed during push and pull experiments in comparison to the BACTRAP approach. As described here, the application of *in situ* microcosms can significantly reduce the amount of labelled material and gives in principle similar information as obtained in push and pull experiments. When using ^{13}C labelled compounds, the transformation into biomass can be traced by analysing the isotope composition of fatty acids and perhaps may also provide kinetic information in future.




Fig. 3-9: Mass spectra of benzylsuccinate extracted from *in situ* microcosms incubated 92d within an anaerobic aquifer at the field site Zeitz (Germany). (A) BACTRAP loaded with non labelled toluene, (B) BACTRAP loaded with $[^{13}\text{C}]\text{-}\alpha\text{-toluene}$. The fragmentation pattern of ^{13}C labelled benzylsuccinate shows that the ^{13}C atom was found at the position of the former methyl group (C).

3.5 Conclusions

In situ microcosms with ^{13}C labelled substrate can be used to demonstrate *in situ* biodegradation of hardly degradable contaminants such as benzene and toluene under anoxic conditions by analysing the isotope composition of fatty acids extracted from the biomass grown on the BACTRAP. The isotope composition in combination with the pattern of

fatty acids indicated that a complex microbial community was colonising the BACTRAPs, but apparently not all community members were involved in the contaminant degradation. The fatty acid patterns were not of taxonomic value and no sensitive indicator to determine changes in the microbial community within various environmental conditions but isotope composition was very useful to provide evidence for *in situ* degradation. Molecular biological analyses showed that mainly uncultured iron and sulphate reducing communities were colonising the BACTRAPs which reflected the biogeochemical conditions and terminal electron accepting processes in the aquifer. By analysing the mass spectra of metabolites we are able to identify the *in situ* degradation pathways. Benzylsuccinate found on toluene amended BACTRAPs indicated that toluene was degraded via the benzylsuccinate pathway which is typical for sulfidogenic and iron reducing conditions.

In summary, BACTRAPs give qualitative information on *in situ* biodegradation of contaminants. Further more with additional molecular biological methods we were able to identify the structure of the microbial community growing on the BACTRAPs which may reflect the *in situ* microbial community to some extent. Compared to laboratory microcosm studies, the advantage of the BACTRAPs is that the incubation takes place under environmental conditions directly in the aquifer at the field site. At present, mass balances were not possible in BACTRAP experiments which is certainly a limitation compared to closed laboratory systems.

In terms of the assessment of *in situ* biodegradation in contaminated aquifers *in situ* microcosms with labelled substances are relatively simple, low-cost assays to gain information on the potential of the microbial community able to degrade contaminants within a reasonable timeframe under *in situ* conditions.

3.6 Acknowledgements

Dr. Kerry Sublette University of Tulsa, USA provided the Bio-Sep® beads for this work. Nicole Stelzer is supported by the Deutsche Bundesstiftung Umwelt (grant 20004/751). The Department of Groundwater Remediation and the SAFIRA Project of the UFZ is acknowledged to assistance in the field work and for providing well characterised test sites. We are grateful to our colleagues Dr. Matthias Gehre and Ursula Günther for support in the isotope laboratory and to Dr. Hermann Heipieper for critical discussion and valuable suggestions.

4 Assessment of Microbial *In Situ* Activity in Contaminated Aquifers

Matthias Kästner, Anko Fischer, Ivonne Nijenhuis, Roland Geyer, Nicole Stelzer, Petra Bombach, Christoph C. Tebbe, Hans-H. Richnow

(This chapter was published in Engineering Life Science 6 (3), p. 234-251)

4.1 Summary

Microbial ecologists and environmental engineers share the interest in identifying the key microorganisms responsible for compound turnover in the environment and in estimating the respective transformation rates. For successful application of Natural Attenuation processes, a reliable assessment of the *in situ* turnover of a contaminant in an aquifer is essential. Here, we review and present new details of two recently developed approaches concerning the assessment of *in situ* biodegradation: a) determination of biodegradation in a contamination plume by stable isotope fractionation analysis (SIFA) caused by microbial metabolism and b) determination of the actual degradation under the respective environmental conditions in the aquifer by using *in situ* microcosms (BACTRAP[®]) amended with ¹³C-labeled substrates as tracer compounds. Based on stable isotope fractionation analysis, the degradation occurring under anoxic biogeochemical conditions at a respective site can be calculated for the entire plume. This has been shown for benzene and toluene at the Zeitz site and partly for chlorobenzene at the Bitterfeld site. By use of the *in situ* microcosm approach with ¹³C-labeled compounds, the microbial *in situ* degradation under strictly anaerobic conditions could be proven for benzene and toluene in Zeitz and for chlorobenzene in Bitterfeld. The transformation of ¹³C-carbon of the labeled substrate into microbial fatty acids proved the assimilation of the pollutant resulting in the formation of biomass. In addition, metabolites such as benzylsuccinic acid were found in the toluene- amended microcosms indicating anaerobic degradation of toluene. This result corresponds to the geochemical conditions found at the field site and therefore, the microcosm approach with ¹³C-labeled compounds can be used to assign the predominant *in situ* degradation pathways in a contaminated aquifer. Since fatty acids profiles alone are often too unspecific for a community analysis at species level, the composition of the microbial communities were analyzed by genetic profiling and sequencing of partial 16S rRNA genes PCR-amplified from total DNA extracted directly from the microcosms. Sequences retrieved from the microcosms indicated a dominance of not yet cultivated bacteria. Several sequences were phylogenetically closely related to sequences of bacteria known to be iron and sulfate reducers, typically found at sites polluted with BTEX and/or mineral oil. The results show that the current methods for

monitoring microbial *in situ* activity at present stage are valuable tools for improving environmental control of compound turnover and will speed up engineering approaches.

4.2 Introduction

Since Winogradski's column experiments, microbiologists and biochemists have long tried to link the transformation of compounds in the environment with the key microorganisms involved in the respective transformation. This linking is a crucial prerequisite for shedding light into the black box of real world microbial processes. From a technical perspective, it is also necessary to have a detailed knowledge about the key processes for purposes of control and successful engineering. In the past decade, methodological advances in linking the structure of microbial communities with the actual activity and function of their members generated dramatic progress in the knowledge of microbial mediated environmental processes and led to the development of new innovative approaches such as waste water treatment using the Anammox process (Schmidt et al., 2003).

The fate of pollutants in contaminated aquifers is mainly governed by microbial degradation. Abiotic processes such as dispersion, dilution, sorption, and volatilization may contribute to a decrease in concentration but do not lead to a significant reduction of the mass of contaminants. Therefore, the evaluation of *in situ* biodegradation is essential for the implementation of Natural Attenuation (NA) concepts in ground water management strategies (US-EPA, 1999). Plate counting methods were used in the past for assessment of the number of microbes present; however, these methods only provide information about the cultivable species. Advanced conventional methods apply the laborious quantification of electron balances of typically complex electron donor - terminal electron acceptor interactions. Calculations then relate the decreasing contaminant concentrations to the consumption of electron acceptors in order to assess biodegradation processes (Clement et al., 2002). When tracing the fate of single compounds in contaminant mixtures, this approach is complicated particularly on the field scale in the very heterogeneous natural environment where several electron acceptor-donor interactions compete. Laboratory microcosm studies are often used to obtain information on *in situ* biodegradation. However, this approach is not reliable because the majority of microorganisms have not yet been cultivated (Ringelberg et al., 1997; Vestal and White, 1989). In particular, the reproducible cultivation of anaerobic bacteria that degrade typical contaminants such as BTEX and PAHs can be difficult and may lead to misinterpretation of the intrinsic biodegradation potentials. Laboratory microcosm studies or percolation column experiments, simulating the contaminant degradation under controlled water flow conditions, are time-consuming; furthermore, the conditions in the laboratory are mostly different to the natural environmental conditions resulting in certain limitations in the reliability of the data acquired.

Recently, the ^{13}C -enrichment of phospholipid-derived fatty acids (PLFA) related to the consumption of ^{13}C -labeled substrates was used to characterize bacterial toluene degradation in soil, sediment or aquifer microcosms (Hanson et al., 1999; Pelz et al., 2001a; Pelz et al., 2001b) and to trace the assimilation of toluene along a food chain (Mauclaire et al., 2003; Pombo et al., 2002). The ^{13}C -tracer concept can provide useful information on carbon fluxes, metabolites and microbial food webs. At the field scale, microbial incorporation of ^{13}C -labeled acetate into biomarker molecules such as PLFA and DNA was successfully assessed to indicate microbes which were suggested to be responsible for the reduction of uranium(VI) (Chang et al., 2005). A mesocosm was designed to monitor the *in situ* dynamics of the microbial community in a BTEX polluted aquifer (Hendrickx et al., 2005). In this study, uncontaminated aquifer material was incubated either in the uncontaminated area or in a contaminated area of the aquifer located nearby and the developed bacterial communities were analyzed by molecular biology techniques. Other authors applied the electron donors glucose and ethanol to acidic uranium and nitrate contaminated groundwater. Fortunately, they were able to show the increase of 16S rRNA gene sequences related to previously cultured metal reducing δ -proteobacteria by 1-2 orders of magnitude using quantitative PCR techniques and time resolved sampling at the site (North et al., 2004). These approaches provide valuable information on the functional biodiversity in soils and sediments. However, from a perspective of the contaminant history they do not provide reliable information on the compound turnover actually occurring under *in situ* conditions in an aquifer. In addition, it is difficult to provide any predictive data on the future fate of the residual contaminants under varying environmental conditions. These approaches may even fail, if organisms with necessary physiological properties were not cultured previously, or if the physiological capacities of the respective organisms are not generally expressed even though the organisms are present in the contaminated aquifer. Such effects are known from *Dehalococcoides* sp. (Holscher et al., 2004; Kube et al., 2005).

The objective of the present work is to review the recently developed approaches focusing on the assessment of the *in situ* biodegradation in contaminated aquifers. Two approaches will be described dealing with a) determining the cumulated biodegradation in a contamination plume by use of fractionation of stable isotopes at natural abundance during microbial metabolism and b) the assessment of the actual degradation under the respective environmental conditions in the aquifer. The latter approach employs *in situ* microcosms (BACTRAP[®]) amended with ^{13}C -labeled substrates such as toluene, benzene, and monochlorobenzene and enables the reliable quantification of the actual degradation. It will also enable the identification of the microbial key players in the near future. We present new results from applying the two approaches at field sites.

4.3 Materials and methods

4.3.1 Chemicals

The chemicals and solvents were obtained in p.A. quality from Merck unless otherwise stated. [$^{13}\text{C}_6$] benzene, [^{13}C]- α -toluene and benzylsuccinic acid were obtained from Sigma-Aldrich (St. Louis, USA). [$^{13}\text{C}_6$] monochlorobenzene and [$^{13}\text{C}_7$] toluene were purchased from Chemotrade Leipzig (Germany). All the isotopically labeled compounds had a chemical purity higher than 99 %.

4.3.2 Field Sites (Zeitz, Bitterfeld)

Zeitz (Saxony-Anhalt, Germany). The benzene, toluene, ethylbenzene and o,m,p- xylene (BTEX) contaminated aquifer is located in the area of a former hydrogenation plant close to the town of Zeitz in Saxony-Anhalt, Germany. A detailed description of site hydrogeology and hydrochemical conditions of the contamination plume at the SAFIRA (Remediation research in regional contaminated aquifers) site was given in previous studies (Fischer et al., 2004; Vieth et al., 2005). The thickness of the upper aquifer varies between 4 to 6 m. Depth discrete investigation of geochemical and isotope parameters in Zeitz have been previously described (Stelzer et al., 2006a). In the source area BTEX exceeded concentrations of 900 mg L⁻¹ and benzene and toluene were present in concentrations up to 850 mg L⁻¹ and 50 mg L⁻¹, respectively. Ethylbenzene and xylenes are typically present in concentrations lower than 3 mg L⁻¹. The groundwater was sampled from the wells indicated in Fig. 4-2. At the test site the predominant electron acceptor used for biodegradation was sulfate (Vieth et al., 2005) and the contribution of methanogenic processes in the microbial transformation in the BTEX-plume is of minor importance (Fischer et al., 2004). Other electron acceptors such as oxygen, nitrate and iron play a minor role for the overall biodegradation processes at the site.

Bitterfeld (Saxony-Anhalt, Germany). The monochlorobenzene (MCB) contaminated aquifer is located near the town of Bitterfeld in Saxony-Anhalt, Germany. Concentrations up to 30 mg L⁻¹ of MCB and lower concentrations of chlorinated aliphatic compounds (< 5 mg L⁻¹), benzene (< 0.77 mg L⁻¹), and dichlorobenzenes (< 3 mg L⁻¹) were found in the aquifer (Weiss et al., 2002). Due to the 150-year history of the chemical industry at the site, the time of spillage is not exactly known but chlorobenzenes have been produced at the industrial site since the first half of the last century. The hydrogeological situation is characterized by a shallow Quaternary aquifer system consisting of glaciofluvial sand and gravel and a deeper Tertiary aquifer system of micaceous sand overlying glauconite sand. A lignite layer overlaid by a Miocene clay layer formerly separated the Quaternary and Tertiary aquifer systems, which has been partially removed by open-cast mining. Today the two aquifer systems are

hydraulically connected at several locations. A Rupelian clay layer is the major regional aquitard of the Tertiary aquifer system (Heidrich et al., 2004a; Heidrich et al., 2004b). The MCB plume stretches down-gradient at the bottom of a Quaternary hydrogeological channel structure in a south-easterly direction (Kaschl et al., 2005). This Quaternary channel is located above the lignite layer, the local aquitard, separating the Quaternary from the underlying Tertiary aquifer. Oxygen concentrations in the aquifer were generally below 1 mg L⁻¹. Significant nitrate concentrations were only found in the upper layers of the aquifer indicating that nitrate reduction was not a relevant terminal electron acceptor (TEA). Some iron and manganese reduction was indicated by low Fe(II) and Mn(II) concentrations. Elevated sulfide concentrations associated with sulfate depletion indicated sulfate reduction and sulfate concentrations of up to 1.48 g L⁻¹ suggested that sulfate was the principle TEA in this aquifer (Kaschl et al., 2005). Methane was not found in the aquifer indicating that methanogenesis was a very unlikely electron acceptor process. The groundwater in Bitterfeld was sampled in a cross section A-B between 17 and 21 m below ground (Left fringe: wells SAF11, SAF12, SAF27, SAF28, and right fringe: SAF7, SAF30, SAF31, SAF23, see Fig. 4-7 and Fig. 4-8) of the lower aquifer in monitoring wells which are perpendicular to the contamination plume as described in details elsewhere (Kaschl et al., 2005). In addition, in well SAF 4/97 the upper, Quaternary aquifer was investigated during a depth discrete multilevel sampling campaign.

4.3.3 Isotope fractionation and calculations

The carbon isotope composition of contaminants and fatty acids is reported in δ-notation (‰) relative to the Vienna-PeeDee-Belemnite-Standard (V-PDB) with $^{13}\text{C}/^{12}\text{C} = (11237.2 \pm 2.9) \times 10^{-6}$ (Craig, 1957) (Eq. 4-1)

$$\delta^{13}\text{C} [\text{\textperthousand}] = \left(\frac{(^{13}\text{C}/^{12}\text{C})_{\text{Probe}}}{(^{13}\text{C}/^{12}\text{C})_{\text{Standard}}} - 1 \right) \times 1000$$

(Eq. 4-1)

The determination of the fractionation of stable isotopes caused by primary enzymatic transformation reaction of a compound during microbial degradation can be described with the Rayleigh equation (Hoefs, 1997; Mariotti et al., 1981). The Rayleigh equation (Eq. 4-2) can be used to relate the change in concentration (C_t/C_0) to the instantaneous change in isotope composition (R_t/R_0) by the isotope fractionation factor (α).

$$\frac{R_t}{R_0} = \left(\frac{C_t}{C_0} \right)^{\left(\frac{1-\alpha}{\alpha} \right)}$$

(Eq. 4-2)

R_0 and R_t are the isotope ratios at the beginning of the transformation reaction ($t = 0$) and after a given time (t), respectively. C_0 and C_t are the corresponding concentrations at the beginning of the reaction ($t = 0$) and after a distinct time (t), respectively. The isotope fractionation factor α quantifies the extent of isotope fractionation. The isotope fractionation is often reported as the enrichment factor $\varepsilon [\text{‰}] = (\alpha - 1) \times 1000$.

Isotope fractionation factors are determined in laboratory reference experiments with microcosms, column experiments or pure cultures where microbial degradation is the only sink of contaminants. In this case the change in isotope ratio (R_t/R_0) can be related unequivocally to the change in concentration (C_t/C_0). In a double logarithmic plot of C_t/C_0 and R_t/R_0 , the isotope fractionation factor can be determined from the slope of the linear regression curve. Isotope fractionation factors (α_D , α_C , α_{Cl}) published until 2004 can be obtained from recent reviews (Meckenstock et al., 2004a; Schmidt et al., 2004) or can be obtained from a frequently updated data base (www.isodetect.de).

The isotope fractionation concept for quantifying the microbial *in situ* degradation is based on the fact that *in situ* biodegradation is the major process altering the isotope composition of priority contaminants such as BTEX, chlorobenzenes and chlorinated ethenes in ground water to a significant extent. Other processes such as sorption and volatilization may affect the isotope composition but do not alter the isotope composition to a significant extent in comparison to biodegradation. Therefore, the isotope composition of contaminants in ground water provides information about the cumulated extent of intrinsic transformation. The isotope fractionation caused by biodegradation leads to an enrichment of heavy isotopomeres in the residual fraction of the contaminant. A modified Rayleigh equation is used to quantify the microbial *in situ* degradation degree (Biodegradation; $B[\%]$). $B[\%]$ represents the concentration decrease expected along a theoretical streamline plug flow without mixing and a single degradation process with a constant isotope fractionation factor. The change in concentration (C_t/C_0) down gradient from the source of contaminants is calculated employing the changes in the isotope ratio between the source (R_0) and a monitoring well (R_t) and an appropriate isotope fractionation factor (α) (Eq. 4-3).

$$(Eq. 4-2) \quad B[\%] = \left(1 - \frac{C_t}{C_0} \right) * 100 = \left[1 - \left(\frac{R_t}{R_0} \right)^{\left(\frac{1}{\alpha-1} \right)} \right] * 100$$

4.3.4 Preparation and incubation of *in situ* microcosms

The *in situ* microcosm experiments were performed using the BACTRAP system described elsewhere (Büning et al., 2005; Geyer et al., 2005). Briefly, Bio-Sep® beads (provided by K. Sublette, University of Tulsa, Tulsa, USA) were loaded with contaminants as carbon source

for microorganisms. The spherical beads, 2 to 3 mm in diameter, consist of powdered activated carbon (PAC) incorporated within an aramid polymer matrix (Nomex®). The beads have a porosity of 75%, an internal surface area greater than $600 \text{ m}^2 \text{ g}^{-1}$ and outer pores of 1-10 microns (Peacock et al., 2004; White et al., 2003). The beads were heated at 300 °C for 4 hours to remove organic residues and then 0.5 g of beads per trap were filled into perforated Teflon® tubes and fixed with glass wool plugs at both ends. The filled microcosms were autoclaved at 121°C for sterilization and hydration of the bead material and then air-dried for 24 hours. The microcosms were loaded with the substrates (benzene, toluene or chlorobenzene) via gas phase under reduced pressure. The substrate and the microcosms were placed in small glass containers and evacuated for at least 48 hours to 60 mbar in order to allow uniform adsorption of the contaminants on the beads. In order to keep the beads anoxic, the vacuum was released by filling oxygen free water into the glass container and the beads were stored under anoxic conditions until their deployment in the monitoring wells.

The BACTRAPs for the microcosm experiments in Zeitz were loaded with [$^{13}\text{C}_6$] benzene to a concentration of about 88 mg g $^{-1}$ bead material and with [^{13}C]- α -toluene to 112 mg g $^{-1}$. The microcosms were deployed in the well Saf Zz 32/02 at a depth of 14 - 15 m (4 m below the groundwater table). The *in situ* microcosms for the experiments in Bitterfeld were amended with [$^{13}\text{C}_6$] chlorobenzene to a concentration of 38 mg g $^{-1}$. For control experiments, the BACTRAPs were loaded with non-labeled benzene, toluene or chlorobenzene in similar concentration. Material from these control microcosms was used for molecular biological studies and to investigate the isotope composition of fatty acids at natural abundance.

A multi level packer system (MLPS) was used for depth discrete sampling from well SAF 4/97 in Bitterfeld (Kaschl et al., 2005; Schirmer et al., 1995) which was equipped with a filter screen from 4.5 to 22.3 m depth below ground. The multi level packer system is a water-filled sock tube inserted into the well, which allows separating various sampling systems to be separated from each other at different depths. Small submersible pumps allowed depth-specific sampling without cross-currents. A detailed description of the multi-level sampling system is given by (Schirmer et al., 1995). The microcosms were deployed in this well for 49 d at a depth of 10.7 m below the surface. The well is located in the anoxic central part of the plume and by the conventional sampling 0.3 mg MCB L $^{-1}$ was found. In the depth discrete sampling campaign, concentrations below 1.0 mg L $^{-1}$ were found up to the depth of 18 m and the concentration increase to more than 13 mg L $^{-1}$ in the deeper part of the well.

4.3.5 Groundwater sampling and analyses

Groundwater sampling for the determination of concentrations and isotope signatures was performed according to good sampling practice. No specific efforts are needed and the samples can be taken in parallel during conventional groundwater monitoring campaigns.

Sampling techniques are described in the literature for example according to the German Industrial Norm (DIN, 1985; DIN, 1996). The biological activity in the samples can be stabilized by adding HgCl_2 , sodium azide or other stabilizers depending on the target compound. Water samples with target compounds resistant to hydrolysis such as hydrocarbons can be stabilized by rising or lowering the pH. For the sampling of BTEX and monochlorobenzene (MCB) a 1 L bottle was completely filled with groundwater. NaOH pellets were added in order to raise the pH to at least 10 and the bottle was closed with a Teflon lined screw cap.

The sensitivity of gas chromatography-combustion-isotope ratio mass spectrometry (GC-C-IRMS) systems is lower than conventional gas chromatography-mass spectrometry (GC-MS) techniques and therefore GC-C-IRMS requires more analyte in order to obtain reliable data. For reproducible carbon isotope analysis about 12 ng carbon is necessary. For hydrogen isotope analysis 5 to 10 times more material is needed. Therefore, the efficiency of the extraction technique to enrich the target compound is critical for the lowest concentration that can be analyzed. Depending on the target, solvent extraction, solid phase micro extraction and purge & trap techniques have been shown to be appropriate for isotope analysis (Meckenstock et al., 2004a; Schmidt et al., 2004).

The solvent extraction procedure which is described in more detail elsewhere (Kaschl et al., 2005; Richnow et al., 2003a; Vieth et al., 2005) was employed for the isotope analysis of BTEX and MCB in this work. Briefly, depending on the estimated concentrations, 1-2 ml n-pentane were used for extraction of 1 L of groundwater. Although this extraction is not quantitative, no isotope effects were associated with this extraction method (Dempster et al., 1997) and MCB and BTEX compounds can be analysed to concentration as low as some $\mu\text{g L}^{-1}$.

The beads of the microcosms were extracted using a dichloromethane-methanol-water mixture as solvent modified according to Bligh and Dyer (1959). After phase separation the dichloromethane phase containing the fatty acids (FA) was evaporated to dryness and derivatized using a trimethylchlorosilane (TMCS) methanol mixture (1:8; v:v) as reactant for 2 h at 70 °C to obtain fatty acid methyl esters (FAME) (Thiel et al., 2001). After evaporation to complete dryness the FAME fraction was dissolved in n-hexane for subsequent analysis by GC-MS and GC-C-IRMS.

For the determination of concentrations and the identification or structural characterization of FAME and benzylsuccinic acid methyl ester by GC-MS, a Hewlett Packard 6890 gas chromatograph coupled to a 5973 quadrupole mass spectrometer (Agilent Technology, Palo Alto, USA) was used. The carboxylic acid fraction was separated on a BPX-5 column (30 m \times 0.32 mm \times 0.25 μm ; SGE, Darmstadt, Germany) with a temperature program of 120°C initial temperature for 4 min, heat at 4°C min^{-1} to 250°C, heat at 20°C min^{-1} to 300°C, and

hold for 10 min. FAME were identified by co-injection of an authentic standard mix (bacterial acid methyl ester mix, Supelco International) and quantified relative to an external standard mix or an internal standard (Henicosanoic acid, C21:0). The fatty acids are designated in the form of A:B ω C where A is the number of carbon atoms, B is the number of double bonds and C is the distance of the closest double bond from the aliphatic end of the molecule (unsaturation, ω -nomenclature). The prefix i (iso) and a (anteiso) refer to methyl branching. The benzylsuccinic acid methyl ester was characterized by co-injection and comparison with mass spectra obtained for the authentic reference compound which was derivated as described below.

The carbon isotope composition of the carboxylic acids fractions was analyzed using a GC-C-IRMS system. It consists of a gas chromatograph (6890 Series, Agilent Technology, Palo Alto, USA) coupled via a Conflow III interface (ThermoFinnigan, Bremen, Germany) to a MAT 252 mass spectrometer (ThermoFinnigan, Bremen, Germany). The performance of the measurement has been described previously (Richnow et al., 2003a). A BPX-5 column (50 m*0.32 mm*0.5; SGE, Darmstadt, Germany) was used for chromatographic separation with helium as carrier gas at a flow rate of 1.5 mL min⁻¹ and a temperature program with initial temperature of 60 °C for 2 min, heat at 20 °C min⁻¹ to 120 °C, heat at 2 °C min⁻¹ to 300 °C, and hold for 20 min (Miltner et al., 2004). The carbon isotope composition analysis of BTEX and MCB on the same system was described previously (Kaschl et al., 2005; Vieth et al., 2005).

The methylation of carboxylic acids for gas-chromatographic analyses introduces an additional carbon into the structure of the analyte which affects its isotopic composition. Therefore the isotope signature of fatty acids ($\delta^{13}\text{C}_{\text{FA}}$) was corrected for the isotope effect due to derivatization FAME with methanol as described previously (Abraham et al., 1998; Abrajano et al., 1994; Goodmann and Brenna, 1992).

Molecular biological analyses of the beads from the exposed microcosms were performed according to the following procedure: For each sample, a total of 50 beads were transferred into 15 ml Falcon tubes containing 12 ml of TES buffer (50 mM NaCl₂, 10 mM Na₂EDTA, 50 mM Tris (hydroxymethyl) aminomethane hydrochloride [Tris HCl], pH 8.0) with 1% sodium dodecylsulfate (SDS). Five cycles of freeze-thaw lysis were performed, each with 5 min in liquid nitrogen (-196 °C) and 5 min at 65 °C in a shaking water bath, each cycle was interrupted by 10 s of vigorous vortexing. DNA was then extracted with phenol-chloroform, as described elsewhere (Schwieger and Tebbe, 1998; Schwieger and Tebbe, 2000). The isopropanol precipitated and purified DNA of each tube was resuspended in a total of 40 µl of 10 mM Tris in 10 µl aliquots. Aliquots were kept at 4 °C for further analysis or stored at -20 °C.

Partial 16S rRNA genes, including two variable regions, were amplified from total DNA by PCR, using primers Com1 and Com2ph (Schmalenberger et al., 2001). The PCR products were converted into DNA-single strands by lambda-exonuclease digestion, following the protocol of Schwieger and Tebbe (1998). Genetic profiles of each sample were generated by single strand conformation polymorphism (SSCP) on non-denaturing polyacrylamide gels, as described elsewhere (Dohrmann and Tebbe, 2004; Tebbe et al., 2001). Single bands were selected for further DNA-sequencing after PCR-amplification and cloning in *Escherichia coli*. Sequencing is described in the same references. Consensus sequences were compared to database sequences using the Fasta Nucleotide Database Query provided by the European Bioinformatics Institute (<http://www.ebi.ac.uk/fasta>).

4.4 Results and Discussion

4.4.1 Benzene and Toluene degradation at the SAFIRA reference site Zeitz

Stable isotope fractionation analyses (cumulated biodegradation approach)

In order to obtain information about the degradation processes that have already been performed by the microorganisms at a contaminated site, the cumulated biodegradation approach was developed based on stable isotope fractionation analysis (SIFA). The overall workflow is summarized in Fig. 4-1. First, the concentrations of BTEX in the wells of the SAFIRA reference site Zeitz were analyzed where benzene is the predominant compound (Fig. 4-2A). Next, the natural carbon isotope composition of benzene in the course of the plume was determined (Fig. 4-2B). The centre of the plume with concentrations up to more than 950 mg L^{-1} was characterized by an average isotope composition of $-28.5 \pm 0.5\text{\textperthousand}$. Down gradient the concentration decreases to below 1 mg L^{-1} at the fringes of the plume and the benzene isotope composition was enriched to maximum values of $-24.4 \text{ \textperthousand}$ indicating biodegradation (Fischer et al., 2004).




Fig. 4-1: Approach I: Assessment of the cumulated biodegradation in a contaminated aquifer by use of stable isotope fractionation analysis (SIFA) at natural abundance; for details see text.

In order to quantify the extent of biodegradation, it is necessary to determine how a given microbial metabolic pathway fractionates $^{13}\text{C}/^{12}\text{C}$ during transformation. Isotope fractionation by microbial degradation is a result of the higher transformation rate of lighter isotopomers compared to heavier isotopomers by enzymatic reactions (Bisswanger, 2000; Galimov, 1985). This kinetic isotope fractionation enriches the amount of the ^{13}C isotopomer in the residual substrate. The isotope fractionation factor (α) describes the relation between the decrease in concentration and change in the isotope ratio during the degradation reaction and can be described by the Rayleigh equation (Eq. 4-2). The isotope fractionation factor can be obtained in reference degradation experiments with pure cultures or at least well defined cultures in which the degradation pathway or the redox conditions are known and in which no other elimination process may alter the concentrations except biodegradation (Meckenstock et al., 2004a). The extent of isotope fractionation depends on the initial enzymatic degradation reaction of the respective biochemical pathway. Without the knowledge about terminal electron acceptor process and degradation pathway, isotope fractionation factors from microcosm studies are not very precise because they may vary upon experimental conditions. Finally, with an appropriate isotope fractionation factor, the percentage of biodegradation along a flow path in a contamination plume can be calculated by the data from the SIFA using equation 4-3.

The metabolic degradation pathway governs the extent of isotope fractionation and therefore, it is important to apply the appropriate isotope fractionation factor for calculation. For

example, from pure cultures able to degrade toluene it is known that the aerobic degradation and the resulting isotope fractionation can vary significantly depending on the initial oxidation reaction at the methyl group or at the aromatic ring system (Morasch et al., 2002). However, under anaerobic conditions the benzylsuccinate pathway attacking toluene at the methyl group shows quite similar isotope fractionation factors under nitrate-, iron, and sulfate reducing conditions (Heider et al., 1999; Meckenstock et al., 1999; Morasch et al., 2001; Spormann and Widdel, 2000; Widdel and Rabus, 2001). At a contaminated site, the microbial pathway relevant for the degradation can be deduced by the redox conditions in the aquifer or by assessment of the primary metabolites of the compound of interest (Griebler et al., 2004b). In case of toluene, benzylsuccinate was detected in Zeitz, which supports the assumption that sulfate reducing conditions govern the anaerobic degradation of toluene in the aquifer. Similarly, the fractionation of benzene also varies under these conditions ($\alpha = 1.0036$) but is lower under methanogenic conditions ($\alpha = 1.002$) (Mancini et al., 2003). Because sulfate was the major electron acceptor in Zeitz, here we used the fractionation factor for benzene degradation under sulfate reducing conditions for calculating the *in situ* degradation. To show the uncertainty, we also calculated the degradation with a fractionation factor characterizing methanogenic conditions ($\alpha = 1.002$). Using the larger fractionation factor ($\alpha = 1.0036$) characteristic for sulfate reducing conditions, the interpretation of the benzene isotope composition give a more conservative estimate (up to 69 % degradation) whereas the calculation with the lower factor ($\alpha = 1.002$) indicative for methanogenic conditions suggest a higher degradation (up to 88%). A field survey over the contamination plume enabled mapping of degradation of contaminants (Fig. 4-3). A previously published monitoring campaign in Zeitz show similar results (Vieth et al., 2005). Examinations of other wells or other BTEX contaminated sites are described elsewhere (Büning et al., 2005; Geyer et al., 2005; Stelzer et al., 2006a; Stelzer et al., 2006b).



A**B**

Fig. 4-2: A) Spatial distribution of the benzene contaminations in the wells of the Zeitz aquifer in 2002 and 2004 (X = location of sampling wells; Saf Zz 32/02 well of the *in situ* microcosm experiment). B) Spatial distribution of the carbon isotope composition of benzene in the wells of the Zeitz aquifer.

In addition, a number of field studies exist where the enrichment of heavy isotopes in the residual fraction of mostly BTEX or chlorinated ethenes was used to estimate the *in situ* degradation (Chartrand et al., 2005; Fischer et al., 2004; Griebler et al., 2004b; Lollar et al.,

2001; Meckenstock et al., 2002; Morrill et al., 2005; Peter et al., 2004; Richnow et al., 2003a; Richnow and Meckenstock, 1999b; Richnow et al., 2003b; Steinbach et al., 2004; Vieth et al., 2001; Vieth et al., 2005; Vieth et al., 2003). For other compounds such as naphthalenes, chlorinated benzenes, phenols, fuel oxygenates (MTBE, ETBE, (Zwank et al., 2005)) less experience exists showing the need of further field studies. The rapid increase of the references within the last five years shows a broad acceptance of the general approach and the reliability of the technique will be further improved with increasing data sets for isotope fractionation factors. At present a range of fractionation factors for several compounds under different environmental conditions is available by recent reviews (Meckenstock et al., 2004b; Schmidt et al., 2004) and data bases (www.isodetect.de). Particularly for the mono-aromatic BTEX contaminants, the applicability of the technique is highly reliable. However, at present the cumulated biodegradation or SIFA approach is of limited value for contaminants such as tert-butyl alcohol, tri-, dichloroethylene, and vinylchloride which can be both, substrate or metabolite. Aquifers contaminated by tetra- and trichloroethene may show a sequential transformation pattern with varying isotope compositions which are not easy to interpret (Martin and Ertl, 2005).




Fig. 4-3: Percentage of benzene degradation in the wells of the contamination plume calculated by use of equation 3 (isotopic composition of the source $[R_0] = -28.5\text{\textperthousand}$, $\alpha = 1.0020$ (methanogenic) – 1.0036 (sulfate reducing); modified from (Fischer et al., 2004).

***In situ* microcosms with stable isotope labeled substrates (BACTRAP approach)**

At many sites, however, there is only limited access to the contamination plume due to the small number of wells or the insufficient data base for the respective compounds. Therefore, a method to prove the microbial degradation of the contaminants is necessary which can be

applied directly in the groundwater monitoring wells. In order to provide this proof and to estimate the turnover and the relevant pathway, the *in situ* microcosm approach was developed (Geyer et al., 2005). The system consists of beads containing activated carbon in a granular shape that are loaded with ^{13}C -labeled contaminants as shown for benzene or toluene at the Zeitz site. The beads provide a large amount of internal surface area for colonization or attached growth of degrading bacteria (biofilm formation). If the indigenous bacteria of the well colonize the beads and productively consume the contaminants under *in situ* conditions, the ^{13}C -labeled carbon will be incorporated into the growing cells and can be traced within biomarker molecules such as fatty acids or nucleic acids. The overall workflow of the approach is presented in Fig. 4-4.




Fig. 4-4: Approach II (A): Assessment of biodegradation potentials in a contaminated aquifer by use of *in situ* microcosms (BACTRAPS) with tracer compounds labeled with stable isotopes; (B): microcosm containing Bio-Sep® beads. (Picture: Copyright (2003) from Environmental Forensics, 2003, 4(1) by White et al.; reproduced by permission of Taylor & Francis Group, LLC., <http://www.taylorandfrancis.com>)

In order to analyze the microbial activity, sterilized *in situ* microcosms were loaded with ^{13}C -labeled benzene or toluene and were exposed for 32 d in well Saf Zz 32/02 in Zeitz. Non-amended microcosms served as controls. After exposure and development of the biofilms on the beads, the residual substrates and the lipid biomarkers were extracted and analyzed by means of mass spectrometry. About 88 to 112 mg of benzene and toluene per g of bead

material in the microcosm was initially applied and around 16 mg g⁻¹ was recovered after exposure (Tab. 4-1). Isotopic analysis of the residual amounts of ¹³C-benzene or ¹³C-toluene in the recovered microcosms from Saf Zz 32/02 in September 2003 showed only a slight accumulation of the contaminants from the aquifer in the bead material in this well. In July-August 2004, however, an accumulation of up to 5.6 mg g⁻¹ from the aquifer was observed in the controls after 112 d of exposure as a result of the much higher contaminant concentrations of about 300 mg L⁻¹ at this time. The overall standard deviation of the recovered substrates from the BACTRAP systems amounted to < 10 %.

Tab. 4-1: Concentrations and ¹³C abundance of contaminants in the *in situ* microcosms (BACTRAP®s) before and after exposure in well Saf Zz 32/02 (modified from Geyer et al., 2005).

	Before exposure		After 32 d		
	¹³ C (atom%)	load (mg g ⁻¹ a))	¹³ C (atom%)	conc. (mg g ⁻¹ a))	loss (%)
benzene	0.1	88 ± 1.0	0.1	15.8 ± 0.2	82
toluene	0.1	112 ± 1.1	0.1	16.0 ± 1.4	85
[¹³ C ₆]-benzene	98.0	88 ± 0.2	98.0	15.8 ± 0.2	82
[¹³ C ₁]-α-toluene	14.0	112 ± 1.2	14.0	17.8 ± 1.4	84
control	-	0.0	0.1	0.46 (benzene) ^{b)}	-
(control)	-	0.0	0.1	5.6 (") ^{c)}	-

a) per g bead material

b) non-loaded control; accumulation from aquifer contaminant plume in September 2003

c) non-loaded control; accumulation from the aquifer after 112 d of exposure in July-August 2005 showing much higher benzene concentrations and lower biofilm formation

Total lipid fatty acid (TLFA) patterns of extracts and their isotopic compositions, as well as polar lipid fatty acids (PLFA) of the beads from the microcosms exposed in well Saf Zz 32/02 were analyzed and the TLFA patterns are shown in Fig. 4-5. TLFA from both, ¹³C-benzene and ¹³C-toluene amended microcosms showed ¹³C-enrichment of up to δ = 13360 ‰ in the fatty acids, providing clear evidence for benzene and toluene biodegradation with transformation of the ¹³C into the microbial biomass. The decrease in contaminant concentrations of the microcosms can thus be considered to be attributable to microbial activity. Extracts from ¹³C-toluene-amended beads showed incorporation of ¹³C into more or less the same fatty acids as found in the ¹³C-benzene microcosm.



A**B**

Fig. 4-5: GC-MS profiles and isotope enrichment of fatty acids in total lipid extracts obtained from the [$^{13}\text{C}_1$]- α -toluene (A) and [$^{13}\text{C}_6$] benzene (B) amended microcosms deployed in well Saf Zz 32/02 in the Zeitz aquifer for 32 days. Major fatty acids with retention time in brackets given in minutes were 14:0 (13.83), i15:0 (15.46), a15:0 (15.66), 15:0 (16.42), i16:0 (17.98), 16:1ω7c (18.36), 16:0 (18.91), 18:1ω9c (22.96), 18:1ω7c (23.10), 18:0 (23.59), br16:0 (19.92, 20.41, 20.60) with undetermined position of methyl-branched, * phthalate (21.31) from extraction process. (The $\delta^{13}\text{C}$ values for the isotopic composition of the respective methyl esters represent the isotopic shift compared to the PDB standard.)

Estimation of the ^{13}C recovery in the biomass. The incorporation of ^{13}C into the fatty acid fraction causing a shift of up to δ 13360 ‰ indicates that [^{13}C]- α -toluene (~14.5 atom %, see Tab. 4-1) was more intensively incorporated into microbial biomass than [$^{13}\text{C}_6$] benzene (98 atom %). This was expected because toluene is considered to be more easily degradable

than benzene. For 16:1 ω 7c, the isotope composition was 13360 ‰ (13.9 atom %), very similar to the toluene substrate, which indicates that at least one organism producing this fatty acid used the applied toluene nearly exclusively as a carbon source. The isotope signatures of C15:0, C16:0 and 18:1 ω 7c of 7946 ‰ (9.1 atom %); 6283 ‰ (7.6 atom %); and 9684 ‰ (10.7 atom %), respectively, demonstrate that other organisms in the consortium assimilated the labeled C from toluene to a major extent but not exclusively. Fatty acids with a lighter isotope composition such as for 18:0 with 92 ‰ (1.2 atom %) indicated that some organism colonized the *in situ* microcosms using other carbon sources than the labeled toluene, e. g. other contaminants or natural organic matter to a major extent.

The most intensively labeled fatty acid on the [¹³C₆] benzene amended microcosms was 16:1 ω 7c with an isotope composition of 13410 ‰ corresponding to 13.9 atom %. The isotope composition of [¹³C₆] benzene was about 98 atom % and therefore around 1/7th of the carbon was derived from the labeled benzene. Thus, the majority of carbon was derived from non labeled carbon sources which may comprise non-labeled benzene from the contamination, natural organic carbon sources or CO₂ used for synthesis of fatty acid. The C16:0 and C18:0 fatty acids were much lower labeled indicating that some organism use benzene only to a minor extent for biosynthesis. Almost non-labeled fatty acids (C12:0) suggest that parts of the microbial community colonizing the beads are not involved in biodegradation of benzene and use other carbon sources exclusively.

The concentrations and isotope compositions of individual fatty acids were used to calculate the amount of ¹³C converted from the contaminant according to the method described by Richnow et al. (1999a). Since the isotope composition of fatty acids from the non-amended *in situ* microcosms were typically between -35 and -25 ‰, it was considered that an isotope composition enriched to more than -20 ‰ was significantly affected by the transformation of the labeled contaminants. On the benzene amended trap about 3.1 nmol ¹³C g⁻¹ bead material corresponding to 43 ng ¹³C g⁻¹ labeled benzene was bound in the fatty acid fraction. The ¹³C- α -toluene derived carbon bound within fatty acids was an order of magnitude higher. The amount of 456 nmol toluene derived carbon g⁻¹ corresponds to the equivalent of 6062 ng labeled toluene. Therefore, the assimilation of C from toluene was more than 140 times higher than that of benzene. Considering that fatty acids account for up to 5 % of the total biomass carbon and that the carbon isotope signature in lipids is similar to that in other fractions of the biomass, the overall amount of labeled carbon bound within the biomass is 20 times higher than found in the fatty acids. If we expect furthermore, that around 95% of the carbon source is metabolized and 5 % is used for biosynthesis of biomass ($y_{xs} = 0.005$), the total productive metabolism can be estimated as 2400 µg toluene and 17.2 µg benzene during incubation in the aquifer. These values only account for 2.5 % of the toluene

loss and 0.024 % of the benzene loss. The ^{13}C -label of benzene, however, was highly diluted due to the background concentrations of benzene from the aquifer.

Phospholipid fatty acids (PLFA) are parameters for microbial membrane phosphoglycerolipid content and thus an indicator for viable cells (Guckert et al., 1985). The total PLFA amounts were highest in the ^{13}C -toluene microcosms with 8870 pmol and 8480 pmol in the ^{12}C -toluene microcosm (Geyer et al., 2005). The microcosms with ^{13}C - or ^{12}C -benzene and the control contained 5860 pmol, 2410 pmol and 860 pmol PLFA, respectively. Applying the conversion factor of $1.4 - 4.0 \times 10^4$ cells pmol $^{-1}$ of PLFA, these values correspond to 2.3×10^7 - 2.4×10^8 bacterial cells with the size of *Escherichia coli* (White et al., 1996) verifying a considerable microbial colonization in the bead material of the microcosms.

Even though no strictly anaerobic pure cultures able to degrade benzene under iron or sulfate reducing conditions are available yet, the *in situ* microcosm approach with ^{13}C -labeled compounds could prove the microbial *in situ* degradation of benzene under strict anaerobic conditions in Zeitz. At present, only nitrate or perchlorate reducing pure cultures responsible for the degradation of benzene have been described (Chakraborty and Coates, 2005). The transformation of ^{13}C -carbon from the labeled substrate into microbial fatty acids provided clear evidence for the assimilation of the pollutant resulting in the formation of biomass. This easy to handle and cost-effective microcosm approach can be used to provide the information required to implement Monitored Natural Attenuation (MNA) as requested by the OSWER directive (US-EPA, 1999) for approval of Natural Attenuation as a treatment measure for contaminated sites.

Metabolites. The activated carbon of the beads may enrich degradation metabolites that can be used to identify the involved degradation pathways. The formation of labeled benzylsuccinate upon toluene degradation was observed in the Zeitz aquifer (data not shown) and is indicative for the anaerobic degradation pathways of aromatic compounds in microorganisms using sulfate and iron as terminal electron acceptors (Heider et al., 1999; Spormann and Widdel, 2000; Widdel and Rabus, 2001). The non-labeled analogue was also found in the control microcosms indicating a degradation of the aquifer contaminants under the respective conditions. Reusser et al. (2002) applied push and pull experiments with deuterium labeled toluene and xylene to verify *in situ* degradation of the compounds in a contaminated aquifer. Although deuterium labeled contaminants may be useful tracer compounds for proving *in situ* degradation, much higher amounts of labeled substances are needed during push and pull experiments compared to the BACTRAP approach. The application of *in situ* microcosms as described here can significantly reduce the amount of labeled material and gives in principle, information similar to that obtained from push and pull experiments. However, by using ^{13}C -labeled compounds in the *in situ* microcosms the

transformation into biomass can be traced by analyzing the biomarker molecules providing the proof of principle for biodegradation.

Biodiversity. The biodiversity of the microbial biomass from *in situ* microcosms exposed in well Saf Zz 32/00 in summer 2004 over a period of 7 to 98 d was characterized by cultivation independent molecular biological approaches based on PCR-amplified partial 16S rRNA genes from total DNA (Scow and Hicks, 2005). Total DNA was directly extracted from the *in situ* microcosms and the PCR products amplified from the various samples were compared to each other on the basis of their SSCP profiles (Fig. 4-6). The community profiles were typically composed of approx. 20 bands with about 5 to 10 bands being more dominant. Most of the profiles generated from *in situ* microcosms taken after different incubation times were relatively similar to each other, indicating a highly reproducible colonization pattern of the bead material. For most samples analyzed, the bacterial community compositions on the benzene and toluene loaded microcosms were not severely altered in comparison to the non-amended controls although highly labeled microbial fatty acids extracted from the ^{13}C -experiments showed a substrate degradation and incorporation into biomass (Tab. 4-1). This indicates for a coordinated degradation by certain members of the indigenous microbial community colonizing the *in situ* microcosms and channeling of the ^{13}C carbon into the microbial food web. However, the isotope composition showed that these organisms also used other carbon sources. Benzene and toluene were the major carbon sources but other BTEX compounds, contaminants, or humic substances were certainly present and may be used as additional carbon sources.

The comparison of the profiles of the non-amended controls with the loaded microcosms showed that the DNA and FA patterns were only slightly different. This reflects the fact that the bead material (activated carbon) of the control (blank) extracted a certain amount of contaminants in particular benzene from the contaminated aquifer (Tab. 4-1) leading to similar conditions in the non-loaded and benzene-loaded microcosms. A comparison of the SSCP-profiles of the bead samples with the profiles of the groundwater samples from the same well showed that with the exception of two bands all bands found in the water were also present on the beads; conversely, only three bands found on the beads were not present in the groundwater samples (Fig. 4-6A). Hence, it can be concluded that the bead material provided a nearly non-selective surface for the growth of the indigenous microbial community from BTEX contaminated groundwater.

DNA-patterns from the toluene loaded microcosms were also highly similar indicating that toluene degraders were already members of the colonizing community and may have conducted the toluene degradation at the site. Cluster analysis of the profiles (Fig. 4-6B) revealed four major groups (Fig. 4-6B). Cluster I contained communities from all microcosms

collected after short time exposure (24 d) or without loaded substrates (T1, see also legend of Fig. 4-6).




Fig. 4-6: SSCP Gel (A) and digital image analyses of SSCP-profiles (B) of PCR amplified 16S rDNA isolated from groundwater and microcosms loaded with toluene or benzene and controls (blanks) after exposure for 24 d (T1), 49 d (T2), and 92 d (T3) in well Saf Zz 32/02. Selected bands of these profiles were sequenced for identification, numbers indicated in the gel image can be found in Tab. 4-2.

In addition, two toluene loaded microcosms from a later sampling (T3) were found in this group. Cluster II contained samples from microcosms loaded with benzene. Other replicates of this treatment and sampling were found in Cluster III which mainly contained samples from substrate loaded treatments, collected later during the kinetic study. As expected on the one hand, the community profiles generated from the surrounding groundwater was different from the biofilm material extracted from the beads (Cluster IV), but on the other hand, similarities

between both sample types were evident. This indicated that in fact the dominant bacteria from the surrounding groundwater contributed to the biofilm communities. Interestingly, the similarity of the groundwater samples was higher with Cluster III (later samplings) than that with Cluster I, which can be explained by the fact that several of the bacteria from the aquifer were only slowly growing on the bead material.

Due to the similarity of the profiles to each other it was in most cases not really possible to identify bands which were highly characteristic for a certain substrate or period of incubation. However, for toluene, a characteristic band seems to be generated during the incubation (Fig. 4-6, band 3 and 4). DNA-sequencing of selected bands revealed that most bacteria in this community were Proteobacteria from the Alpha- und Beta-subgroup (Tab. 4-2). Similarities to known sequences in the database for these partial rRNA genes ranged from 85 to 100 %. Complete similarity was found with an uncultured bacterial clone isolated from monochlorobenzene contaminated groundwater (Alfreider et al., 2003). Other sequences, which were derived from the toluene-specific bands, were found to originate from *Azoarcus* sp., one of them most closely related to an organism that has been shown to degrade toluene and *m*-xylene under denitrifying conditions (Barragan et al., 2004; Zhou et al., 1995). Despite the fact that 16S rRNA genes are generally not reliable indicators for bacterial functions, the so far characterized sequences give no rise to assume growth of sulfate-reducing or other strictly anaerobic bacteria but rather indicates the presence of bacteria capable of growing under denitrifying conditions. Other sequences indicated α -proteobacteria adapted to the exposition with metals, assuming physiological similarities with their closest relatives (Lopez-Archilla et al., 2004; Stein et al., 2002). However, it is known from other strictly anaerobic bacteria e. g. *Dehalococcoides* sp. that no PCR products were found with the universal COM1/COM2 primers.

Tab. 4-2: Characterization of rRNA genes isolated from groundwater and microcosms loaded with toluene or benzene and controls after exposure in well Saf Zz 32/02 (see also Fig. 4-6)

Band No. ¹⁾	Length [bp]	Similarity [%]	Closest relative or sequence (Accession No.)	Phylogenetic group	Source of isolation or habitat of closest relative or relevant metabolic activity of closest relative
1	370	100	uncultured bacterium (AY05094)	Betaproteobacteria	monochlorobenzene contaminated aquifer
2	205	85	<i>Acidovorax</i> sp. (AJ864847)	Betaproteobacteria	high mountain lake habitat
3	370	99	<i>Azoarcus</i> sp. (AF515816)	Betaproteobacteria	degradation of toluene and benzene under denitrifying conditions
4	360	95	<i>Azoarcus denitrificans</i> (L33690)	Betaproteobacteria	growth on toluene under denitrifying conditions
5	350	93	uncultured bacterium (AY475201)	Alphaproteobacteria	from metal rich, acidic river Tinto (Spain)
11	210	90	uncultured bacterium (AF418953)	Alphaproteobacteria	from metal rich freshwater reservoir

¹⁾ PCR-products obtained from bands indicated in figure 6A; no products were obtained from other bands.

4.4.2 Monochlorobenzene degradation at the SAFIRA reference site Bitterfeld

Stable isotope fractionation analysis (cumulated biodegradation approach)

Monochlorobenzene (MCB) is a xenobiotic compound which has been commonly used for chemical manufacturing processes over decades and is a common groundwater contaminant at many chemical production sites in Europe and the United States (Arneth et al., 1989; Schiedeck et al., 1997). In addition, higher chlorinated benzenes can be degraded via reductive dechlorination to MCB under anaerobic conditions (for review see Van Agteren et al., 1998) and may be also produced from the microbial transformation of hexachlorocyclohexane (HCH) in anoxic aquifers (Phillips et al., 2005; Van Eekert et al., 1998). MCB is the major contaminant of the pollution plume in the Quaternary anoxic aquifer in Bitterfeld. The aquifer belongs to the regional aquifer system, in which about 200 million m³ of groundwater are contaminated with predominantly chlorinated compounds (Heidrich et al., 2004a; Heidrich et al., 2004b). Field studies demonstrating the anaerobic microbial transformation of MCB are completely lacking. Thus, no pathways and mechanisms of anaerobic transformation of MCB are yet known. Taking into account that the plume formed in the aquifer several decades ago, a significant selection pressure for MCB degrading microorganisms can be expected. Therefore, assuming biodegradation we applied the SIFA approach (Fig. 4-1) to the complex contamination plumes in Bitterfeld (Kaschl et al., 2005).




Fig. 4-7: Monitoring wells and distribution of the monochlorobenzene plume in Bitterfeld modified from (Kaschl et al., 2005). Concentrations are given by the dotted lines in mg L⁻¹. The arrows indicate the groundwater flow in November 2002. The solid line (A to B) represents the position of the cross-section perpendicular to the monochlorobenzene plume at the location of the SAFIRA research station. (Saf 4 = monitoring well for the vertical analysis).

The MCB plume stretches downgradient in a SE-direction, following a Quaternary hydrogeological channel structure oriented in a NW/SE direction, which strongly affects

contaminant flow. Since the access to the groundwater by wells was limited, we decided to analyze a cross section of wells A-B perpendicular to the groundwater flow in the lower aquifer which covers the concentration gradients at both sides of the plume (Fig. 4-7). The concentrations of MCB increased up to 20 mg L^{-1} in the centre of the plume and the isotopic signatures increased from values of -26.7 ‰ at the centre to -23.0 ‰ at the fringes of the plume (Fig. 4-8 left side) showing a significant ^{13}C -enrichment in the residual MCB at the fringes. In addition, we analyzed the vertical distribution of MCB and its isotope signature in the upper aquifer by multilevel sampling in well SAF 4/97 showing the same trends with a clear vertical MCB stratification (Fig. 4-8 right side). The ^{13}C -enrichment found at the fringes of the plume relative to the source area of about 4 δ units clearly shows the occurrence of *in situ* biodegradation of MCB. This would result in a field isotope fractionation factor of $\alpha = 1.0005$. However, field fractionation factors are always lower than factors obtained from laboratory experiments in closed systems in which only biodegradation can diminish the concentrations. In the field, the concentrations may be affected by abiotic effects leading to much smaller fractionation factors suspending the calculation of the effective biodegradation according to equation 4-3.




Fig. 4-8: Monochlorobenzene concentrations and isotopic signatures of the cross-section A – B, depth of 17–21 m (left) and the vertical profile in Saf 4 (right); (modified from Kaschl et al., 2005); ● = MCB concentrations; □ and ○ = $\delta^{13}\text{C}$ of MCB

Several degradation pathways are conceivable under anaerobic conditions. MCB may be degraded similarly to other aromatic compounds by processes such as nitrate-, Fe(III)-, sulfate-reduction, or fermentation coupled to methanogenesis (Chakraborty and Coates, 2005; Heider et al., 1999; Spormann and Widdel, 2000; Ulrich et al., 2005; Widdel and Rabus, 2001). In addition, microbial reductive dechlorination of MCB requiring an electron donor may also be considered. Batch experiments using several aerobic pure cultures of bacteria showed that the known aerobic pathway initiated by dioxygenases did not result in a

significant carbon isotopic fractionation (Kaschl et al., 2005). Conversely, the field isotope fractionation factor for MCB obtained in the anoxic aquifer was higher, especially if dispersion and dilution effects are accounted for (Kaschl et al., 2005). Thus, it is evident that in this aquifer a novel anaerobic pathway resulting in an isotopic fractionation which is untypical for aerobic MCB transformation must be considered to be the predominant process of the *in situ* MCB degradation. However, the present data do not allow the anaerobic pathway of MCB degradation to be identified. The transformation of MCB appears to be very slow and most of the microbial activity appears to be located at the plume fringes.




Fig. 4-9: Rayleigh plot illustrating field-derived and laboratory-derived fractionation factors: aerobic degradation of MCB in batch experiments ($\alpha=1.0001-1.00028$), the field fractionation factor for the whole MCB plume ($\alpha=1.00043$), the (\blacktriangle) left (Saf11, Saf12, Saf27, Saf28) and (\bullet) right fringe (Saf7, Saf30, Saf31, Saf23) of the plume at the cross-section A - B ($\alpha=1.00083$ and 1.00106 , respectively) and, for comparison, batch experimental data for the reductive dechlorination of TCBs ($\alpha=1.0031-1.0037$ (Griebler et al., 2004a) and anaerobic degradation of benzene ($\alpha=1.0019-1.0036$ (Mancini et al., 2003)).

Fig. 4-9 illustrates the various fractionation processes likely for the MCB degradation in comparison to the field data. Since these field-derived fractionation factors for MCB are significantly higher than the laboratory-derived factors for aerobic degradation, it is evident that an anaerobic pathway governs the MCB fractionation in the Bitterfeld aquifer. All known aerobic microbial cultures investigated used the ring-dioxygenase pathway to degrade MCB (Vogt et al., 2004) but the dioxygenase pathway did not lead to a significant isotope fractionation of MCB in batch experiments. This result is in accordance with studies using

toluene and naphthalene as model compounds showing that the dioxygenase reaction does not lead to significant carbon isotope fractionation (Morasch et al., 2002). Aerobic degradation of TCB by pure cultures exhibiting the dioxygenase reaction also did not show any significant isotope fractionation (Griebler et al., 2004a). However, the field fractionation factors obtained are lower compared to laboratory experiments investigating the anaerobic degradation of benzene or 1,2,3-trichlorobenzene (TCB) (Griebler et al., 2004a; Mancini et al., 2003). This may be explained by dilution, dispersion and sorption processes in the aquifer, which affect concentrations (C_t) without altering the isotopic composition (R_t). Hence, when using the Rayleigh equation (Eq. 4-2) for field data, the resulting field isotope fractionation factor is always lower than determined in culture experiments. However, in a cross section perpendicular to the contaminant plume, the dispersion and sorption processes should have affected MCB concentrations in a similar way, since the distance from the original source of the contamination is comparable. Accordingly, the fractionation factor determined at the left and right fringe was higher than that of the whole plume and was significantly higher than the low isotopic fractionation expected from aerobic transformation. The anaerobic degradation of various aromatic hydrocarbons led to a significant isotope fractionation of about $\alpha=1.0017$ to 1.0036 (Mancini et al., 2003; Meckenstock et al., 1999; Morasch et al., 2001). In addition, a number of studies have demonstrated that the anaerobic reductive dehalogenation of halogenated ethenes and TCBs is always associated with higher isotope fractionation factors (Griebler et al., 2004a; Griebler et al., 2004b; Hunkeler et al., 1999a; Hunkeler et al., 2002). Analogously, it can be expected that a reductive dehalogenation of MCB should also be associated with a significant isotope fractionation. Alternatively, MCB may be degraded similarly to benzene under anaerobic conditions. Although the anaerobic degradation pathway for benzene has not yet been completely identified, the isotope fractionation of benzene yields fractionation factors of 1.0020 (methanogenic), 1.0036 (sulfate-reducing) and 1.0022-1.0024 (nitrate-reducing conditions) in enrichment cultures (Mancini et al., 2003). A degradation of MCB following the anaerobic benzene pathway should therefore also show a significant isotope fractionation.

***In situ* microcosms with stable isotope labeled substrates (BACTRAP approach)**

The *in situ* approach based upon microcosms amended with ^{13}C -labeled tracer compounds was also applied for MCB in the aquifer in Bitterfeld. The accumulation of MCB in anoxic aquifers initially led to the assumption, that anaerobic biodegradation of MCB is difficult or even impossible, because neither pure nor enrichment cultures and microcosm studies had demonstrated biodegradation of MCB under these conditions. However, the isotopic signatures of the contamination plumes clearly show biotransformation in the course of the MCB plume (Kaschl et al., 2005). BACTRAPS amended with MCB were incubated for 7 weeks in groundwater well Saf 4/97 equipped with a MLPS in order to avoid any oxygen

access changes in redox conditions. Geochemical analysis led to the conclusion that anaerobic conditions prevailed in the well where the BACTRAPS were tested.

After exposition for 49 d, more than 80% of ^{13}C -MCB had been lost from the microcosms and the extracted total lipid fatty acids were found to be significantly labeled (Fig. 4-10). Hexadecanoic acid (16:0) showed the highest incorporation of ^{13}C with $\delta^{13}\text{C}$ of 579 ‰ (Fig. 4-10). Octadecenoic acid (18:1) was also labeled significantly to 158 ‰. Octadecanoic acid (C18:0) and tetradecanoic acid (14:0) showed a lower incorporation of ^{13}C of 10 ‰ and 46 ‰, respectively. Unsaturated hexadecenoic acid (16:1) showed a strong ^{13}C signal indicating high label incorporation but the concentration was also too small for quantitative analysis. Similar to the *in situ* microcosm studies with benzene and toluene, the data provided evidence for the assumption of an unknown anaerobic MCB degradation process actually occurring in Bitterfeld. In addition, the incorporation of the label into marker molecules of the biomass also clearly indicates that this pathway supports growth and energy conservation from MCB for the respective bacteria under the given environmental conditions.




Fig. 4-10: Total ion current of the fatty acid extracts from a BACTRAP amended with $[^{13}\text{C}_6]$ monochlorobenzene exposed for 49 d in an anaerobic aquifer in well Saf 4/97 at the Bitterfeld field site. Identified fatty acids and their isotope compositions are indicated in the boxes.

4.5 Concluding remarks

Both approaches presented in this overview provide promising tools for the assessment of *in situ* microbial degradation activity in contaminated aquifers. They have certain implications and the evaluation of the practical benefits and limitations is currently in progress. The isotope fractionation approach interpreting the isotope signatures of contaminants provides a technique for assessing the cumulated biodegradation in an aquifer which has been occurred

on the groundwater flow path down gradient from a source of contaminants. The SIFA approach relies on the availability of a sufficient number of monitoring wells, the availability of distinct isotope fractionation factors for the respective degradation pathways and biogeochemical conditions, and a detailed knowledge about the hydrology of the aquifer. Even if no specific isotope fractionation factor is available, a qualitative estimation of the biodegradation can be made by comparison with other isotope fractionation factors of known pathways and may provide valuable information on the processes likely to occur in the investigated aquifer. However, if multiple sources of contaminants with different initial isotopic signatures contribute to a mixed plume, this approach may be too complex for reliable quantification of *in situ* biodegradation. In addition, the applicability of the approach is limited, if processes without significant isotope fractionation govern *in situ* biodegradation as shown for aerobic toluene degradation pathways catalysed by mono- or dioxygenase enzymes reacting with the aromatic ring system (Meckenstock et al., 1999).

The *in situ* microcosm approach (BACTRAP) with isotopic labeled contaminants and analysis of the isotope composition of lipid biomarkers provides an excellent tool for assessment by proving the microbial *in situ* activity in contaminated aquifers and for monitoring natural attenuation processes. This relatively simple, low-cost approach provides information in a reasonable time frame and the results are more reliable than *ex situ* approaches. Even if the details of the metabolism of a target compound are not yet known, valuable information about the presumable pathway can be obtained by this approach. In particular, if metabolites of the parent compound can be identified on the bead material of the microcosms, a clear indication for the relevant degradation pathways can be given by this approach. Future investigations will provide the basis for a quantification of the biodegradation potential by this approach. Therefore, it is necessary to investigate the sorption behavior of the bead material in the presence and absence of bacteria. The key organisms related to contaminant degradation could not yet be identified; however, this aspect of the approach is currently under investigation. Future experiments to investigate the kinetics of the colonization as traced by quantitative PCR may elucidate the development of communities and may provide indication for organisms supported by the test substrate. When the concentration of the label in the biomass is sufficiently high, stable isotope probing (SIP) can also be employed to identify organisms which use the ¹³C-labeled contaminants as a carbon substrate (Lueders et al., 2004; Manefield et al., 2004; Whiteley et al., 2006). This may open the "black box" of subsurface microbial ecology by linking structure and function.

4.6 Acknowledgements

Dr. Kerry Sublette University of Tulsa, USA provided the Bio-Sep® beads for this work. The fellowship of Nicole Stelzer is supported by the Deutsche Bundesstiftung Umwelt (grant

20004/751). The Department of Groundwater Remediation and the SAFIRA Project of the UFZ are acknowledged for assistance in the field work and for providing well-characterized test sites. We are grateful to Dr. Matthias Gehre and Ursula Günther for support in our isotope laboratory. The authors thank the BMBF (grant 02WT0022) and the EU for financial support within the 6th framework research project WELCOME (EVK1-2001-00132) and the Marie Curie Development Host Fellowship project BIOISOTOPE (EVK1-CT-2000256120).

5 Sensitive detection of anaerobic monochlorobenzene degradation using stable isotope tracers

Ivonne Nijenhuis, Nicole Stelzer, Matthias Kästner, Hans-H. Richnow

(This chapter was published in Environmental Science and Technology 41(11), p. 3836-3842)

5.1 Abstract

Microbial degradation of monochlorobenzene (MCB) under anaerobic conditions was investigated using a stable isotope tracer under *in-* and *ex situ* conditions. *In situ* microcosms were incubated directly in an anoxic aquifer and carbon derived from [$^{13}\text{C}_6$] MCB was found to be incorporated into the microbial biomass. In laboratory microcosms, amended with [$^{13}\text{C}_6$] MCB, anaerobic mineralization of MCB was indicated by the production of $^{13}\text{CO}_2$. Further, recovery of the ^{13}C -label in the fatty acids confirmed the assimilation of MCB derived carbon into microbial biomass. The described approach may be applied to various other organic groundwater contaminants of concern using carbon (^{13}C) as well as other stable isotope tracers, such as nitrogen (^{15}N), allowing direct and sensitive detection of biodegradation.

5.2 Introduction

Monochlorobenzene (MCB) is a widespread groundwater contaminant found at many former chlorine chemistry sites particularly when chlorinated pesticides have been produced [www.who.int/en/]. The Bitterfeld/Wolfen region in Germany, is a former production site for lindane (hexachlorocyclohexane) where today MCB is present as an ubiquitous contaminant throughout the entire aquifer with concentrations up to 30 mg L^{-1} (Heidrich et al., 2004a; Heidrich et al., 2004b; Popp et al., 2000). While MCB contamination may be the result of a spillage event, it can also be produced *in situ* by microbial lindane degradation or during dechlorination of higher chlorinated benzenes (Adrian et al., 2000a; Adrian et al., 2000b; Boyle et al., 1999; Fennell et al., 2004; Van Agteren et al., 1998; Van Doesburg et al., 2005). MCB is the most mobile chlorobenzene due to its relatively high solubility and lower sorption potential to the aquifer matrix and usually long plumes form indicating its high persistence in anaerobic aquifers. Drinking water limits for MCB in the USA are currently set at $<0.1 \text{ mg L}^{-1}$ [www.atsdr.cdc.gov/toxfaq], in Germany, the quality target for surface waters is set at $1\mu\text{g L}^{-1}$ conform with European Union standards [www.umweltbundesamt.de]. While aerobic degradation of MCB has been well studied (for a review see Van Agteren et al., 1998), complete anaerobic degradation and mineralization has, to our knowledge, not yet been reported. Only the complete anaerobic dehalogenation of the chlorinated benzenes to

benzene has been described but the microbiota involved in the process have not been identified (Nowak et al., 1996).

In recent years, stable isotope approaches have been developed to investigate natural attenuation of contaminants. Stable isotope fractionation analysis may be used to investigate *in situ* biodegradation of pollutants (Meckenstock et al., 2004a), additionally, stable isotopes may be applied as tracer compounds (Geyer et al., 2005; Kästner et al., 2006; Peacock et al., 2004; Stelzer et al., 2006a). Previously, evidence for the biodegradation of MCB in an anoxic aquifer in Bitterfeld was provided applying isotope fractionation techniques (Kaschl et al., 2005).

Recently, *in situ* microcosms (BACTRAP[®]), applying stable isotope tracers, were developed and used to investigate degradation of BTEX compounds in the field (Geyer et al., 2005; Peacock et al., 2004; Stelzer et al., 2006a). These *in situ* microcosms consist of a sorbent material serving as a surface for microbial growth and as a source for the isotopically labeled and unlabeled test substrates. During incubation of these microcosms in an aquifer for several months, microorganisms capable of degradation of the contaminant will incorporate carbon derived from the substrate into their biomass (e.g. fatty acids) and the biodegradation of the contaminant can be proven if the ¹³C-label is recovered from the biomass. A preliminary investigation applying these *in situ* microcosms loaded with [¹³C₆] MCB supported the stable isotope fractionation analysis providing evidence for the biodegradation of MCB under field conditions (Kästner et al., 2006).

In this study, stable carbon (¹³C) tracer methods were applied to investigate the biodegradation of MCB under anoxic conditions. *In situ* microcosm systems were incubated directly in the groundwater at the field site in parallel to a laboratory set up. Production of ¹³C-labeled degradation products and incorporation of the ¹³C into biomass were used to demonstrate anaerobic biodegradation of MCB.

5.3 Materials and Methods


5.3.1 Chemicals

The [¹³C₆] chlorobenzene was purchased from Chemotrade Leipzig (Germany) with chemical and isotopic purity of >99 %. The [¹³C₆] chlorobenzene preparation further contained an impurity of 0.02% [¹³C₆] benzene. All other chemicals were obtained in p.a. quality or higher.

5.3.2 Site description of the aquifer in Bitterfeld (Saxony-Anhalt, Germany)

- The chlorobenzene (MCB) contaminated aquifer is located near the city of Bitterfeld, Germany. Concentrations up to 30 mg L⁻¹ of MCB and lower concentrations of chlorinated aliphatic compounds (up to 5 mg L⁻¹), benzene (up to 0.77 mg L⁻¹), and dichlorobenzenes (up to 3 mg L⁻¹) were found in the Quaternary aquifer (Kaschl et al., 2005). The time of spillage is

not known but chlorobenzenes have been produced at the industrial site since the first decade of the last century. The hydrogeology and geochemistry of the aquifer system was described previously (Kaschl et al., 2005). Oxygen concentrations in the aquifer were analyzed using a flow-through chamber and an oxygen electrode (CellOx 325, Wissenschaftlich-Technische Werksttten, Weilheim, Germany) and were generally below 0.3 mg L^{-1} , except for the highest level, close to the groundwater table where oxygen concentrations were around 1 mg L^{-1} . Using this method, oxygen concentrations below 0.5 mg L^{-1} could not be resolved accurately. Significant nitrate concentrations were only found in the upper layers of the aquifer indicating that nitrate reduction was not a relevant terminal electron acceptor (TEA) in the deeper layers of the investigated aquifer. Some iron and manganese reduction was indicated by low Fe(II) and Mn(II) concentrations. Elevated sulfide concentration associated with sulfate depletion indicated sulfate reduction and sulfate concentrations up to 1.48 g L^{-1} suggested that sulfate was available as the principle TEA in this aquifer (Kaschl et al., 2005). Elevated methane concentrations were not found in the aquifer so methanogenesis apparently was not an important process. Overall, the geochemical analysis points at prevalently anoxic conditions in the aquifer.






Fig. 5-1: A). Multilevel sampling system (MLPS), preventing groundwater circulation and penetration of oxygen from the surface, used for the *in situ* microcosm experiment in well SAF4. The black rectangles indicate the location of the *in situ* microcosms (m1-m6) in general between the filter screen of the well and the membrane of the MLPS, whereas the small rectangles (I-IV) show the detailed experimental set up with [¹²C] MCB (I), Blank (II), [¹³C₆] MCB (III) and Blank (IV) *in situ* microcosms as it was performed in every depth. The black arrows show the groundwater flow direction, gs: ground surface and wt: water table. B). Carbon isotope composition of fatty acid methyl esters (FAME) extracted from [¹³C₆] MCB amended *in situ* microcosms incubated for 7 weeks at different depth (m1-m6) at the multilevel monitoring well SAF4 in Bitterfeld. The grey shaded area indicates the natural abundance of $\delta^{13}\text{C}_{\text{FAME}}$. C) Relative abundance of total lipid fatty acids [FAME] extracted from *in situ* microcosm loaded with ¹³C₆-labelled MCB incubated in different depths (m1-m6) at the monitoring well SAF4 in Bitterfeld.

5.3.3 Preparation and incubation of *in situ* microcosms

The *in situ* microcosms were prepared as described before by Stelzer et al. (2006a). Briefly, Bio-Sep® beads (University of Tulsa, Tulsa, USA) were loaded with MCB as carbon substrate for microorganisms via gas phase under reduced pressure. The *in situ* microcosms for the experiments in Bitterfeld were amended with [¹³C₆] MCB or natural abundance MCB to a concentration of 76 mg g⁻¹ sorbent material. In parallel, unamended *in situ* microcosms (blank) were deployed. Material from these control experiments was used to investigate the isotope composition of fatty acids at natural abundance. The *in situ* microcosms were

incubated for 7 weeks in groundwater well SAF4 equipped with a multilevel sampling device (MLPS) to avoid any oxygen penetration and groundwater circulation between depths (Fig. 5-1; Fig. S1 Supporting Information (Anhang E)) (Schirmer et al., 1995). In the MLPS experiments the microcosms were placed in stainless steel cages as a spacer to ensure a permanent flow of ground water through the *in situ* microcosms and to avoid contamination with biological material by direct contact with the packer membrane or the filter screen of the well. The set of *in situ* microcosms used as an inoculum for the enrichment cultures were incubated directly in the groundwater in the wells without a MLPS or stainless steal cages.

5.3.4 Extraction and derivatization of fatty acids (FA)

After removal of the *in situ* microcosms from the groundwater well, the Bio-Sep® beads were extracted using a dichloromethane-methanol-water mixture as solvent modified after Blight et al. (1959). For investigation of the laboratory microcosms and enrichment cultures, 15 or 3 ml of liquid, respectively, was removed from the microcosms for extraction with the dichloromethane-methanol mixture. After phase separation the dichloromethane phase containing the fatty acids (FA) was evaporated to dryness and derivatized using a trimethylchlorosilane (TMCS) methanol mixture (1:8; v:v) as reactant for 2 h at 70 °C to obtain total lipid fatty acid methyl esters (FAME) (Thiel et al., 2001). After evaporation to complete dryness the FAME fraction was dissolved in n-hexane for subsequent analysis using a gas chromatograph coupled to a mass spectrometer (GC-MS) and a gas chromatography-combustion-isotope-ratio-monitoring-mass-spectrometer system (GC-C-IRMS) (See Supporting Information, Anhang E). FAME were identified by co-injection of an authentic standard mix (bacterial acid methyl ester mix, Supelco, Munich, Germany) and concentrations of FA were quantified relatively to the internal standard (Henicosanoic acid, C21:0). The fatty acids are designated in the form of A:B where A is the number of carbon atoms, B is the number of double bonds.

5.3.5 Laboratory microcosms

Microcosms

Groundwater was taken from 2 wells (SAF11 and SAF30) at the MCB contaminated field site in Bitterfeld. The filter screens of the wells are at 19 to 20 and 19 to 21 meters below ground, respectively. After transport to the laboratory and transfer into an anaerobic glovebox (98-96% N₂/ 2-4% H₂), the groundwater was amended with 1 mg L⁻¹ of resazurin as a redox indicator (Tratnyek et al., 2001). The microcosms were prepared in 120 mL vials to which 100 mL of the groundwater mixture was added. The vials were closed with Teflon coated grey butyl rubber stoppers and crimped with aluminum crimps. For each well, 11 microcosms were prepared. Of each set, 2 microcosms were autoclaved for 40 minutes on three consecutive days to prepare killed controls. Then, 5 of the live microcosms and the 2 killed

microcosms were amended with natural abundance MCB and the other 4 with [$^{13}\text{C}_6$] MCB to a concentration of approx. 100 $\mu\text{mol L}^{-1}$. MCB concentrations were followed by GC-FID analysis. CO_2 concentrations were estimated by GC-C-IRMS vs. an external calibration curve. SAF11 and SAF30 contained 6.7 and 3.7 mmol L^{-1} CO_2 respectively. For details of the analysis see Supporting Information (Anhang E). Sulfide concentrations were determined according to the method of Cline (1969). Iron(II) concentrations were determined by the Department of Analytics at the UFZ-Centre for Environmental Research using the Spectroquant® Iron test (Merck, Darmstadt, Germany).

***In situ* microcosm enrichment**

In situ microcosms loaded with natural abundance MCB from a preliminary experiment incubated for approx. 8 weeks in the groundwater wells SAF11 and SAF30 were used as an inoculum for the enrichment cultures. Therefore, the *in situ* microcosms were recovered after incubation and the sorbent material was added to 27 mL anaerobic culturing tubes filled with groundwater from the monitoring wells. The tubes were closed with Teflon coated grey butyl rubber stoppers and crimped. In the laboratory, the tubes were amended as follows with anoxic sterile stock solutions to a final concentration of: 1 mg L^{-1} resazurin, 2 mg L^{-1} yeast extract, 2 mg L^{-1} vitamin B12, 1 mM Na_2S and either, for the sulfidogenic enrichments 5 mM FeSO_4 or, for the methanogenic enrichments, 2 mM FeCl_2 . Approx. 20 μmol MCB was added to each tube. Due to sorption of MCB to the Bio-Sep® beads, it was not possible to quantify MCB in these enrichment cultures. Qualitatively, MCB appeared to decrease over time and, apparently, microbial conversion occurred (data not shown). After incubation of approx 4 month, sub cultures were made from each *in situ* microcosm enrichment. Liquid from each microcosm was diluted (10%) into fresh basal medium according to the description by Zinder (Zinder, 1998) amended with 0.1 % NaHCO_3 , vitamin solution, 1 mM Na_2S and, as indicated 5 mM FeSO_4 or 2 mM FeCl_2 . Two parallels were prepared for each transfer set: one culture was amended with natural abundance MCB while the 2nd culture was amended with [$^{13}\text{C}_6$] MCB to a concentration of 500 $\mu\text{mol L}^{-1}$.

5.4 Results and Discussion

5.4.1 *In situ* microcosms

The recently developed *in situ* microcosms were directly incubated in the anoxic aquifer in Bitterfeld to investigate the MCB degradation (Geyer et al., 2005; Peacock et al., 2004; Stelzer et al., 2006a). Previous studies applying isotope fractionation techniques and a preliminary *in situ* microcosms study provided first evidence that MCB was degraded within this aquifer (Kaschl et al., 2005; Kästner et al., 2006). The *in situ* microcosms were installed inside a monitoring well in a multilevel fashion (microcosm set m1-m6; see Fig. 5-1A). The

use of a multilevel sampling device allowed depth discreet sampling and should prevent circulation of groundwater between the different levels and entry of oxygen to the microcosms during exposure (Schirmer et al., 1995). However, water may circulate between different levels through the filter material of the well screens or the surrounding sediments. After incubation in the groundwater for 7 weeks, the *in situ* microcosms were removed from the monitoring well and used for further analysis. Direct extraction of the sorbent material from the *in situ* microcosm showed that in addition to the parent [$^{13}\text{C}_6$] MCB, ^{13}C -labeled benzene was present at several of the investigated depths (m2, m3, m5, m6) with an isotope signature of $\delta^{13}\text{C} > 1000 \text{ ‰}$ (Tab. 5-1).

Tab. 5-1: Carbon isotope composition of benzene and MCB extracted from blank, unloaded, and [$^{13}\text{C}_6$] MCB loaded *in situ* microcosms incubated for 7 weeks at different depths at the multilevel monitoring well SAF4 in Bitterfeld. n.a.= not analysed. The standard deviation for the analysis of the carbon stable isotope composition was <1‰ with $\delta^{13}\text{C} < 500 \text{ ‰}$ and between 5-10 ‰ with $\delta^{13}\text{C} > 500 \text{ ‰}$.

Depth [m]	Blank	Blank	Labeled
	$\delta^{13}\text{C}_{\text{Benzene}} [\text{‰}]$	$\delta^{13}\text{C}_{\text{MCB}} [\text{‰}]$	$\delta^{13}\text{C}_{\text{benzene}} [\text{‰}]$
m1 (8.1)	3590	190	n.a.
m2 (14.5)	1000	265	1310
m3 (16.5)	485	245	3940
m4 (18.5)	1080	50	n.a.
m5 (20.5)	700	60	8520
m6 (21.9)	560	-25.1	3780

^{13}C labeled benzene was also observed in blank, non-amended microcosms. At depth m6, ^{13}C -labelled benzene ($\delta^{13}\text{C} = 560 \text{ ‰}$) and MCB with an isotope composition typical for this aquifer section ($\delta^{13}\text{C} = -25 \text{ ‰}$) was observed in the blank, non-amended microcosms. Both compounds must have been adsorbed during incubation on the non amended microcosm from the surrounding groundwater and were present in concentrations in a similar order of magnitude, benzene concentrations were approx. one third that of MCB. (See Fig. S2, Supporting Information (Anhang E)). Although [$^{13}\text{C}_6$] benzene was already present in trace concentrations (0.02 %) in the commercial [$^{13}\text{C}_6$] MCB, it is unlikely that the labeled benzene was derived from the impurity of the chemical. Since benzene and MCB have similar physico-chemical properties, it would be not probable that a separation of both chemicals would occur upon diffusion or transport with the groundwater resulting in sorption of labeled benzene without the co-transport of labeled MCB. The results suggest that benzene is

formed upon dehalogenation of MCB in the aquifer. Additionally, at the field site, benzene could be detected in the aquifer in low concentrations (up to $20 \mu\text{g L}^{-1}$).




Fig. 5-2: Carbon isotope signature of CO₂ (♦) and methane (○) in A) a microcosm prepared with groundwater from SAF11 and B) the enrichment culture. Based on the CO₂ content and isotope signature, the amount of [¹³C₆] MCB degraded was calculated in μmol per microcosm (◊) and % remaining of the total added [¹³C₆] MCB (■) for C) the microcosm and D) the enrichment culture. The shaded area in panel A indicates the observed isotope signature of CO₂ in the control microcosms.

To investigate assimilation of MCB derived carbon into bacterial biomass, the isotope composition of fatty acids (FA) was determined. FA extracted from the *in situ* microcosm systems loaded with [¹³C₆] MCB were significantly labeled providing evidence for the complete degradation of MCB via central metabolism and assimilation of MCB derived carbon for synthesis of biomass (Fig. 5-1B). The relative abundance of the various identified FA was similar comparing the microcosms incubated at the different depths m1 to m6 (Fig. 5-1B). C16:0 and C18:0 were the most abundant FA, followed by lower amounts of the mono-unsaturated C16:1 and C18:1. Only minor amounts of C12:0, C14:0 and C15:0 were detected (Fig. 5-1C). Close to the groundwater table (m1) all identified FA were labelled. Possibly, at the transition to the capillary fringe oxygen (1 mg L⁻¹ at m1) is available as electron acceptor for degradation in this zone (Kaschl et al., 2005); therefore MCB may be degraded aerobically at the upper level. At deeper levels (m2-m6), conditions were strictly anaerobic and significant enrichment in ¹³C was observed in the hexadecanoic acid (C16:0; up to δ¹³C = 352 ‰ at depth m6); hexadecenoic acid (C16:1; up to δ¹³C = 503 ‰ at depth m5); and the octadecenoic acid (C18:1; δ¹³C = 94 ‰ at depth m6) (Fig. 5-1A, B). Other FA were not labelled significantly. The incorporation of ¹³C derived from [¹³C₆]-MCB into FA

provided clear evidence for *in situ* biodegradation of MCB under the environmental conditions present in the aquifer.

5.4.2 Laboratory microcosms and enrichments

Laboratory microcosms were used in parallel to investigate anaerobic MCB degradation in a closed system and to attempt cultivation of MCB degrading microorganisms. A first set of laboratory microcosms was prepared from a mixture of groundwater derived from the same MCB plume as described above. A decrease in MCB concentration was observed in several microcosms amended with electron donors (e.g. lactate, benzoate) or acceptors (sulfate), but no intermediates or degradation products were found and the activity could not be sustained (data not shown). For this reason in a second set up of microcosm systems, we supplied isotopically labeled MCB to enable a more sensitive detection of the degradation processes. Two separate sets of microcosms were prepared with groundwater from wells SAF11 and SAF30 and ^{13}C -ring labeled MCB was used as a tracer substance. No MCB decrease relative to the killed control could be observed (data not shown). The concentrations and isotope signatures of benzene, methane and CO_2 were monitored over time. In Fig. 5-2A an example is given of the observed isotope signatures for methane and CO_2 in the investigated time frame. The concentrations of CO_2 did not change significantly over the investigated period (data not shown), but the CO_2 became clearly enriched in ^{13}C (Fig. 5-2A). Over time, the isotope signature of CO_2 increased from $\delta = -24.4$ to $+305\text{\textperthousand}$ in the microcosms prepared from groundwater of well SAF11 (Fig. 5-2A). The isotope signature of CO_2 in the microcosms prepared with groundwater of well SAF30 changed similarly ($\delta = -25.3$ to $+443\text{\textperthousand}$). All replicates of microcosms in both sets (SAF11, SAF30) showed enrichment of the ^{13}C -label in the CO_2 fraction although the lag period for the degradation was highly variable (data not shown). After 700 days of incubation, all microcosms amended with [$^{13}\text{C}_6$] MCB had significantly produced $^{13}\text{CO}_2$. Production of $^{13}\text{CO}_2$ was not observed in any of the killed controls amended with [$^{13}\text{C}_6$] MCB. The enrichment of ^{13}C in CO_2 clearly showed the mineralization of MCB in these microcosms. Additionally, the isotope signature of methane remained stable around $\delta = -40$ to $-45\text{\textperthousand}$ and $\delta = -45$ to $-50\text{\textperthousand}$ for microcosms from well SAF11 and SAF30 respectively and no methane production was observed. Apparently, methanogenesis was not an important process involved in MCB degradation in our laboratory microcosms. Resazurin was added to the microcosms allowing visual monitoring of anoxic conditions during the incubation time. All microcosms remained clear throughout the incubation period indicating a reduction potential of $< -0.051 \text{ mV}$ (Tratnyek et al., 2001). At the end of the study period, soluble Fe(II) and total sulfide was measured to confirm the presence of anoxic conditions and redox buffers in the system (See Tab. S1, Supporting Information (Anhang E)). Sulfide concentrations were in the same order of magnitude,

generally a bit higher, compared to the original groundwater while soluble Fe (II) concentrations were lower confirming anoxic conditions. It should be considered that Fe(II) concentrations could have been underestimated due to precipitation with sulfide. In these microcosms, early in the incubation period (100-300 days), low levels of ¹³C-labeled benzene could be identified by the means of retention time using headspace analysis, potentially derived from the impurity in the labeled MCB preparation. Benzene did not accumulate and disappeared over time suggesting its conversion in the microcosms.




Fig. 5-3: Total ion current of fatty acid methyl esters derived from microcosms prepared with aquifer water of well SAF30. Shown are samples from a killed control, natural abundance MCB amended and [¹³C₆] MCB amended microcosms. Identified peaks are labeled and underlined (_). FAME were identified via GC-MS and the isotope signature in the various samples are shown in ‰ [PDB].

Using the concentration of CO₂ (6.7 or 3.7 mmol L⁻¹ for microcosms from SAF11 or SAF30 respectively) and corresponding isotope signatures in the *ex situ* microcosms, we estimated the amount of MCB mineralized as shown in Fig. 5-2C for the microcosms from SAF11. After 700 days of incubation, 3 out of 4 microcosms from each set incubated with the labeled MCB were estimated to have converted 3 to 4% of the added MCB. Interestingly, in each set, one microcosm had converted up to 11 and 30% of MCB based on ¹³CO₂ evolution for microcosms derived from groundwater from well SAF30 and SAF11, respectively.

Using the analysis of the stable isotope signature of CO₂, degradation could be determined extremely sensitively. A significant enrichment in ¹³C (e.g. δ = +50‰) would, in these microcosms from SAF11, represent an estimated degradation of about 100 nmol MCB corresponding to about 1% of the total added MCB. This order of sensitivity to measure biodegradation would be impossible to obtain using other methods except for methods using radioactive carbon tracers.

To prove incorporation of MCB-derived carbon into microbial biomass, fatty acids were extracted from the laboratory microcosms, derivatized and FA isotope signatures were analyzed. In Fig. 5-3 the total ion currents of 3 selected samples from the set SAF30 are shown. While the amount of biomass was highly similar in the different samples, [$^{13}\text{C}_6$] MCB amended microcosms showed FA which were significantly enriched in ^{13}C (16:1; 16:0; 18:0 and 18:1 with $\delta = +5060$; $+1025$; $+50$ and $+61 \text{ ‰}$, respectively) indicating incorporation of MCB derived carbon into biomass. Comparing the 3 samples, a shift in FA composition could be observed as octadecanoic acid (18:0) was present in the killed control microcosm but nearly absent in the live microcosms. Since the fatty acids found are common for most bacteria and have relatively low specific taxonomic value, no conclusions can be drawn about the microbial community involved in the biodegradation (Kaur et al., 2005).

According to the literature, no anaerobic MCB mineralizing microbial cultures have been described yet. In an attempt to improve the cultivability of anaerobic MCB degrading microorganisms, the sorbent from *in situ* microcosms was used as an inoculum. Therefore, after pre-enrichment of organisms during approximately 8 weeks exposure in wells SAF11 or SAF30, the *in situ* microcosms were removed and the sorbent material was used as an inoculum for enrichments. In these initial enrichments, disappearance of MCB was observed, but quantification of MCB was impossible due to the adsorption of MCB on the sorbent material resulting in only very low dissolved concentrations in the enrichment cultures. After approx. 4 months, 10% liquid transfers were prepared from these enrichments into fresh medium and [$^{13}\text{C}_6$] MCB was added as a tracer. Methane, CO_2 and benzene concentrations as well as their isotope compositions were monitored in these transfer cultures (Fig. 5-2C). The appearance of $^{13}\text{CO}_2$ was observed but, again, no labeled methane was formed. An analysis of the incorporation of carbon into biomass was also attempted. FAs were analyzed but concentrations were too low for a quantitative assessment. On the GC-C-IRMS, labeled products could be observed in the isotope trace at the same retention time as commonly found FA (C16:1; C16:0; C18:1; C18:0) implying the presence of labeled biomass (data not shown). Apparently, only small quantities of biomass were formed in our enrichment cultures which corresponded with the absence of obvious biomass formation and turbidity.

Two major hypothetical degradation pathways would come into consideration for the anaerobic biodegradation MCB: 1) initial dechlorination, followed by the degradation of benzene or 2) direct attack on the benzyl ring with the formation of a substituted chlorinated aromatic substance (e.g. chlorobenzoate, chlorophenol). The first pathway would suggest a similar pattern as observed previously by Nowak et al. (1996). Benzene could then be further degraded via benzoate or phenol as intermediates (Chakraborty and Coates, 2005; Edwards and Grbic-Galic, 1992; Lovley, 2000; Phelps et al., 2001; Ulrich et al., 2005). Alternatively, a chlorinated phenol, benzoate or toluene could be formed which then is dechlorinated via

reductive dechlorination as described for chlorinated benzoate to benzoate by e.g. *Desulfomonile tiedje* or for chlorinated phenols to phenol by *Desulfitobacterium* spp. (Löffler et al., 2003; Mohn and Tiedje, 1990; Van de Pas et al., 1999). Although the observed traces of benzene on blank *in situ* microcosms suggest degradation via dehalogenation, because of the impurity of benzene in the labeled MCB preparation the 1st pathway could not be verified in the laboratory experiments. However, the *in situ* experiments indicated that reductive dehalogenation may occur in the aquifer. Further detailed investigations of the pathway and development of a stable mixed culture and further isolation of microorganisms involved in the anaerobic MCB degradation are required.

To summarize, for the first time we could demonstrate the sensitive detection of anaerobic biodegradation of MCB, which was thus far considered persistent, *in situ* as well as in *ex situ* laboratory studies. A previous investigation supported MCB degradation in an anaerobic aquifer via stable isotope fractionation analysis but no definite proof for anaerobic degradation was presented thus far (Kaschl et al., 2005). By means of stable isotope tracers (¹³C) we provided evidence for anaerobic MCB degradation although only very small portions of the contaminant (nmol) were mineralized. Our concept based on the use of ¹³C labeled tracers in combination with *in situ* and laboratory microcosms was found very valuable to analyze the slow degradation of recalcitrant organic chemicals. The application of modern analytical techniques for isotope analysis enables stable isotope probing of cell components (FA), allowing the sensitive elucidation of carbon fluxes on a molecular level. Although, stable isotope probing of DNA or RNA could give valuable information about the microbial community involved in the degradation of MCB, application of these techniques in our field and laboratory systems is currently not possible yet due to the lack of biomass formed.

The described approach can be applied to detect the biodegradation and to investigate the persistence of a variety of recalcitrant environmental pollutants using ¹³C or ¹⁵N stable isotope tracers which are needed for the synthesis of cell components. Furthermore, we can use the *in situ* microcosms as pre-enrichments to improve cultivation efficiency of the microbial community involved in the degradation processes. In the future, our concept may be applied to investigate the environmental fate of recalcitrant organic chemicals providing valuable support for evaluation of chemicals for regulatory framework such as Registration, Evaluation and Authorisation of Chemicals (REACH) recently developed in Europe.

5.4.3 Acknowledgements

The UFZ and the SAFIRA Project, in particular Ralf Trabitzsch and Holger Weiss, are acknowledged for providing infrastructure and assistance in the field work. We are thankful to Matthias Gehre, Ursula Günther for their technical support in our stable isotope laboratory, Hermann Heipieper and colleagues of the departments for Isotope Biogeochemistry and Bioremediation for discussion and Sylvia Uhlig for performing parts of the practical work. We

Sensitive detection of anaerobic monochlorobenzene degradation using stable isotope tracers

further would like to thank Max Hägglom and Stephen Zinder and three anonymous reviewers for critical reading. Kerry Sublette University of Tulsa, USA provided the Bio-Sep® beads for this work. We thank the European Union Marie Curie Development Host Fellowship (Bioisotope, EVK1-CT-2000256120) for financial support of Ivonne Nijenhuis and the Deutsche Bundesstiftung Umwelt (grant 20004/751) for support of Nicole Stelzer.

Supporting Information Available

Additional information is available on the detailed methods for analysis using GC-MS and GC-C-IRMS, the analysis of the *in situ* and laboratory microcosms (Anhang E).

6 Integrative approach to delineate Natural Attenuation of chlorinated benzenes in anoxic aquifers

Nicole Stelzer, Gwenaël Imfeld, Martin Thullner, Jürgen Lehmann, Alexander Poser, Hans-H. Richnow, Ivonne Nijenhuis

(This chapter is submitted to Environmental Science and Technology)

6.1 Abstract

Assessment of intrinsic microbial activity under anoxic conditions at a chlorobenzene contaminated field site was addressed by hydrogeochemical analysis, stable isotope tools and multivariate statistics to elucidate the fate of chlorinated benzenes with particular emphasize on most persistent monochlorobenzene (MCB). *In situ* microcosm analysis provided evidence of microbial assimilation of MCB-derived carbon and laboratory investigations asserted mineralization of MCB at very low degradation rates. To further investigate *in situ* biodegradation of chlorinated benzenes (CBs), compound-specific stable isotope analysis (CSIA) was applied. During sequential degradation of CBs the isotope signature of a particular chlorobenzene species might be affected simultaneously by depletion and enrichment of ^{13}C , complicating the application of the Rayleigh approach for quantification of *in situ* degradation. Therefore, an isotope balance concept was applied. The enrichment of the cumulative isotope composition of all CBs indicated *in situ* biodegradation. Additionally, the linkage between geochemistry, contaminant concentration and the observed degradation activity was investigated by principal component analysis (PCA) underlining hydrogeochemical heterogeneity at the study site. The application of an integrative approach relying on multiple lines of evidence to document that capability of the intrinsic microbial community to anaerobically degrade MCB and DCB was found useful to characterize the natural attenuation potential of the site.

6.2 Synopsis

Multiple lines of evidence demonstrated *in situ* degradation of chlorinated benzenes in an anoxic aquifer.

6.3 Introduction

Chlorinated aromatic compounds are worldwide intensively used to synthesize pesticides and other chemicals leading to ubiquitous distribution in the environment (Field and Sierra-Alvarez, 2008). Chlorinated benzenes (CBs) can also be formed during anaerobic microbial transformation of hexachlorocyclohexane, of which the γ -isomer (Lindane) is worldwide used as

pesticide (Phillips et al., 2005; Van Agteren et al., 1998). Due to their toxicity, persistence and accumulation in the food chain CBs are of great environmental concern.

CBs may be subjected to both aerobic and anaerobic microbial degradation (Adrian and Gorisch, 2002; Field and Sierra-Alvarez, 2008; Van Agteren et al., 1998). With increasing chlorination CBs may undergo reductive dehalogenation (RDH) either as cometabolic reaction or energy yielding halorespiration (Adrian and Gorisch, 2002; Van Agteren et al., 1998). The occurrence of RDH under methanogenic and sulfate reducing conditions has been observed in a variety of anaerobic mixed cultures (Adrian and Gorisch, 2002; Van Agteren et al., 1998), but only one bacterial strain (*Dehalococcoides* strain CBDB1) capable to couple energy conservation with RDH of CBs (≥ 3 chlorine substituents) has been isolated so far (Adrian et al., 2000b; Jayachandran et al., 2003). RDH generally results in the transformation of higher chlorinated CBs but may lead to an accumulation of lower chlorinated CBs such as monochlorobenzene (MCB) under strongly reducing conditions. Preliminary indications for RDH of MCB have been presented (Nowak et al., 1996). Kaschl et al. suggested that anaerobic MCB degradation may lead to significant stable carbon isotope fractionation and applied compound-specific stable isotope analysis (CSIA) to further indicate *in situ* biodegradation of MCB in an anoxic aquifer (Kaschl et al., 2005). Recently, complete mineralization of MCB has been proven using stable isotope tracer techniques which enabled very sensitive detection of low rate degradation processes (Nijenhuis et al., 2007).

CSIA has been established for monitoring the biodegradation of pollutants at contaminated field sites (Meckenstock et al., 2004a). This method was successfully applied for BTEX (Fischer et al., 2007; Griebler et al., 2004b; Mancini et al., 2002; Vieth et al., 2005), fuel oxygenates (Kuder et al., 2005; Zwank et al., 2005) and chlorinated ethenes (Hunkeler et al., 1999a; Imfeld et al., 2008; Nijenhuis et al., 2005; Sherwood Lollar and Slater, 2001) but, to our knowledge, beside the work of Kaschl et al. (2005) no further field studies exist for CBs. Moreover, most studies were conducted to characterize degradation of single contaminants instead of looking at complex scenarios involving sequential degradation mechanisms, multiple contaminants and/or various degradation pathways.

Microbial degradation of organic contaminants is associated with isotope fractionation, leading to an enrichment of heavier isotopes in the residual non degraded fraction. The magnitude of isotope fractionation mainly depends on the reaction mechanism. As shown for RDH of chlorinated ethenes (Cichocka et al., 2008; Nijenhuis et al., 2005) and trichlorobenzenes (TCB) (Griebler et al., 2004a), degradation of CBs should also be associated with significant isotope fractionation, if the reduction of a Cl-C-bond is involved. Similar to anaerobic BTEX degradation, the anaerobic oxidation of the benzene ring for which cleavage of a C-H-bond is expected should lead to isotope fractionation (Fischer et al., 2008; Mancini et al., 2003). However, if the fate of a contaminant is simultaneously controlled by its production (associated

with depletion in ^{13}C) and further degradation (leading to enrichment in ^{13}C), as expected from sequential RDH, the isotope enrichment might be masked due to opposite isotope fractionation limiting the use of the Rayleigh approach for contaminants (Hunkeler et al., 1999a; Nijenhuis et al., 2005).

At the study site, release of CBs resulted in a severe contamination of the groundwater. It is hypothesized that RDH and/or anaerobic oxidation govern the removal of CBs under anoxic conditions in the aquifer. Since active remediation technologies were recognized as technically and economically not feasible (Thullner and Schäfer, 1999), this study aimed at evaluating the efficacy of natural attenuation (NA) as remedial option.

For this purpose, in accordance with guidelines provided by the US-EPA (1999), we developed an integrative approach including (i) assessment and monitoring of CB degradation and plume stability by means of CSIA and tracer studies under field and laboratory conditions as well as (ii) characterization of hydrogeochemical conditions governing *in situ* biodegradation applying multivariate statistics. Special attention was put in interpreting MCB degradation since its accumulation seems to be the limiting factor to apply NA as remediation strategy. To overcome complications associated with the Rayleigh concept for interpretation of the isotope data, an isotope balance was calculated to assess biodegradation of CBs.

6.4 Material and Methods

6.4.1 Study Site and sampling

The field site is located at a former chemical plant where the production of mostly pesticides caused an intensive release of chlorinated organic contaminants to the subsurface. The direct source zone characterized by high loads of contaminants and dense nonaqueous phase liquid (DNAPL) was completely encapsulated and hydraulically disconnected from the surrounding natural groundwater system to prevent further release of contaminants. The initial contaminant pattern dominated by higher chlorinated benzenes is preserved inside the containment whereas a plume mainly consisting of MCB and DCB with total concentrations up $2840 \mu\text{g L}^{-1}$ has developed. A detailed site description is given in supporting information (SI) (section 1.1, Anhang F).

Groundwater samples for hydrogeochemical and isotope analyses were taken in 2005 and 2006 from 22 monitoring wells (Fig. 6-1). The sampling, extraction procedure and analytical methods are described in SI (Anhang F). A table summarizing the geochemical data collected during sampling in 2005 is given in SI (Tab. S2, Anhang F).

6.4.2 Chemicals

The [$^{13}\text{C}_6$] MCB was purchased from Chemotrade Leipzig (Germany) with chemical and isotopic purity of > 99%. All other chemicals used were obtained in p.a. quality or higher.

6.4.3 Field study with *in situ* microcosms

In February 2005 a field experiment was performed using an *in situ* microcosm test system (BACTRAP) as described elsewhere (Geyer et al., 2005; Kästner et al., 2006; Nijenhuis et al., 2007; Peacock et al., 2004; Stelzer et al., 2006a). Briefly, Bio-Sep® beads were loaded with [¹³C₆] MCB to a concentration of about 100 mg g⁻¹ beads. The *in situ* microcosms were deployed in 6 different wells at the low (H, L), medium (D, E) and high contaminated (A, B) area of the plume in the lower strictly anoxic part of the aquifer (Fig. 6-1 and Fig. S1, Anhang F). After 72 days *in situ* microcosms were removed from the aquifer and the carbon isotope signature of total lipid fatty acids (TFLA) was analyzed. Detailed information about the extraction procedure and analytical methods can be found in SI (section 1.4, Anhang F).

6.4.4 Laboratory study with enrichment cultures

Material of *in situ* microcosms amended with natural abundance MCB as described above were used as inoculum for the laboratory enrichment cultures. From each selected well (A, B, D, E, H, L) four enrichment cultures were prepared in 38 ml vials. Each vial finally contained approx. 27 ml of groundwater and 4-6 Bio-Sep® beads. MCB was added to the vials: always two cultures were amended with 1 µl natural abundant and another two with [¹³C₆] MCB. Cultures were incubated stationary at 20°C in the dark and were sampled at regular time intervals to determine the carbon isotope signature of mineralization products (CO₂, CH₄). A more detailed description of the methods is given in SI (section 1.5, Anhang F).

6.4.5 Determination of δ¹³C values for the total chlorinated benzenes

The isotope signatures of individual CBs ($\delta^{13}\text{C}_i$) were analyzed by GC-C-IRMS and are given in δ-notation (per mill) (for details see SI, section 1.6, Anhang F).

The isotope signature (per mill) of the total CBs ($\delta^{13}\text{C}_{\text{CB}}$) was calculated by multiplying the molar concentration of each compound (C_i) with its respective carbon isotope signature ($\delta^{13}\text{C}_i$), adding all contributions and dividing by the total molar concentration of all chlorinated benzenes (C_{CB}) (Eq. 6-1).

$$\delta^{13}\text{C}_{\text{CB}}[\text{\%}] = \frac{\sum (C_i * \delta^{13}\text{C}_i)}{C_{\text{CB}}}$$

Equation (6-1)

In this paper we refer to the term isotope balance to clearly distinguish between the calculated isotope signature of the total CBs ($\delta^{13}\text{C}_{\text{CB}}$, isotope balance) and the directly measured isotope signatures ($\delta^{13}\text{C}_i$) of the single species of CBs, such as MCB and DCB isomers.

The uncertainty associated with the isotope balance was calculated based on the error propagation using the standard deviation of each chlorobenzene species (Eq. 6-2).

$$\Delta_{tot} \delta_{stdv} = \frac{\sqrt{\sum (C_i * \Delta\delta^{13}C_i)^2}}{C_{CB}}$$

Equation (6-2)

For all samples the standard deviation for the values of the isotope balance was $\leq 0.4 \text{ ‰}$.

6.4.6 Principal Component Analysis

Principal component analysis (PCA) was used to condense the hydrogeochemical and isotope data into a reduced number of orthogonal linear combinations and to gain insight into the relationship between variables. The numerical data matrices were converted and the correlation analyses carried out using the program R (R: Copyright 2005, The R Foundation for Statistical Computing Version 2.1.1). PCA was performed on the correlation matrix applying the logarithmic transformation to meet normality requirements. Two data sets were subjected to PCA: (i) the hydrogeochemistry data (Tab. S2, Anhang F), isotope composition of MCB and of total CBs corresponding to the year 2005 (Tab. S4, Anhang F), as well as (ii) concentrations of MCB and DCB isomers, isotope composition of MCB and the total CBs corresponding to years 2005 (Tab. S4, Anhang F) and 2006.

6.5 Results and Discussion

6.5.1 *In situ* microcosms

The patterns of TFLA extracted from *in situ* microcosms are provided for one representative well located at the low (H), medium (E) and high contaminated (A) area of the plume (Fig. S1, Anhang F). All samples showed significant amounts of saturated hexadecanoic (C16:0) and octadecanoic (C18:0) acids as well as the unsaturated hexadecenoic (C16:1) and octadecenoic (C18:1) acids. Further, an octadienoic acid (C18:2) was identified in all samples. Comparison of the three samples revealed only minor differences in the fatty acid (FA) compositions. While sample H was dominated by C18:0, the most abundant FA in the other two samples was a C18:1 isomer. In sample A, octadecanoic acid was one order of magnitude less abundant compared to E and H (Fig. S1, Anhang F). TLFA profiles showed low taxonomic value similar to previous findings (Geyer et al., 2005; Stelzer et al., 2006a). The total concentrations of TLFA varied between 4000-12000 pmol per microcosm (data not shown) suggesting a significant microbial colonization in the bead material similar to previously published results (Geyer et al., 2005).

PLFA may offer a sensitive measure to characterize viable bacterial community structures whereas TLFA fraction can additional comprise lipids of dead biomass (Green and Scow, 2000; Kaur et al., 2005; Lu et al., 2007). Basically, the microbial community trapped on *in situ* microcosms may consist of both viable and non-living organisms as well as degraders and non-

degraders. Therefore extraction of TLFA was favoured over PLFA to investigate more sensitively the incorporation of ^{13}C into total biomass as an indicator for *in situ* activity of the bacterial community.

Compared to unlabeled controls ($\delta^{13}\text{C}_{\text{FAME}} = -30 \pm 5 \text{ ‰}$, data not shown), the TLFA methyl ester fraction of labeled samples showed a clear enrichment of ^{13}C up to 4500 ‰ in individual fatty acids (Tab. 6-1). Fatty acids with 16 carbons generally represented the highest enrichment in ^{13}C and unsaturated FA were higher labeled than the saturated ones as similarly observed by Nijenhuis et al. (2007). FA with odd chain length (C17) showed the highest $\delta^{13}\text{C}$ value of 4500 ‰ which corresponded to 6 atom % incorporation, but was only found in sample A in very low quantity. For some FA (C18:2, C18:1) no label was found (-31 to -23 ‰) indicating part of the microbial community were not involved in MCB degradation.

However, the transformation of the labeled carbon from the [$^{13}\text{C}_6$] MCB into bacterial fatty acids provided evidence of microbial degradation of MCB under ambient aquifer conditions although an enrichment of ≤ 6 atom percent indicated that microorganisms colonizing the *in situ* microcosms mainly used other carbon sources than the labeled MCB. In general, the labeling was comparable in all three samples but slightly higher at A suggesting that the presence of a more MCB adapted microbial community correlated with high MCB concentrations.

Tab. 6-1: Carbon isotope signature of TLFA methyl esters $\delta^{13}\text{C}_{\text{FAME}}$ [‰] extracted from *in situ* microcosms amended with [$^{13}\text{C}_6$] MCB and incubated in the contaminated aquifer at the study site.

Well [C_{MCB} in $\mu\text{g L}^{-1}$]	A [1400]	E [430]	H [120]
C16:1	2600	2000	2000
C16:1	nd	nd	2900
C16:0	1600	70	10
iC17:0	4500	nd	nd
C18:2	-28	-31	-31
C18:1	-28	-31	-31
C18:1	360	43	1
C18:0	50	-25	-31
C21:0	-23	-26	-30

nd - not detectable

6.5.2 Laboratory enrichment cultures

To prove the potential for mineralization of MCB by the microflora colonizing the *in situ* microcosms, cultivation-dependent methods were used in the laboratory. Surfaces are very

often required for successful cultivation and pre-incubation in the field may improve the later cultivation in the lab (Herrmann et al., 2008). Therefore, material of *in situ* microcosms, which were directly incubated at the field site, served as pre enriched-inoculum. Application of ^{13}C -labeled substrates in enrichment cultures is one of the few approaches suitable to sensitively detect complete mineralization of a single contaminant in a complex mixture determining the evolution of labeled CO_2 (Morasch et al., 2007). Addition of [$^{13}\text{C}_6$] MCB allowed exclusive detection of anaerobic MCB degradation, a supposedly slow process (Nijenhuis et al., 2007). After 197 days highest enrichment of ^{13}C in CO_2 was observed for the samples from well L with values of $629 \pm 85\text{‰}$ and lowest for wells H and D with values of $91 \pm 59\text{‰}$ and $130 \pm 40\text{‰}$ respectively. Analyses of enrichment cultures from A, B and E showed comparable results with $\delta^{13}\text{C}_{\text{CO}_2}$ of $250 \pm 91\text{‰}$, $222 \pm 19\text{‰}$ and $296 \pm 122\text{‰}$, respectively (Tab. S3, Anhang F). Labeled methane could not be detected indicating that MCB degradation was not coupled to methanogenesis. All unlabeled controls showed no ^{13}C enrichment in CO_2 during the course of the experiment.

The extent of MCB mineralization was calculated according to the method of Morasch et al. (2007) based on the $^{13}\text{C}-\text{CO}_2$ production during the course of the experiment in respect to the total CO_2 initially analyzed from the groundwater of the respective wells (Tab. S3, Anhang F). Mineralization rates varied between 0.1 to 1.1 nmol per day indicating very slow MCB mineralization underlining the difficulties associated with cultivation of such organisms. The results are in good agreement with data obtained from Nijenhuis et al. (2007) for a similar experiment at another field site. An electron balance to link MCB degradation to iron or sulfate reduction was not feasible due to too high background concentrations of potential electron donors (Tab. S2, Anhang F). In all samples MCB mineralization was further maintained over time up to 1000 days (data not shown). However, the underlying pathway could not be elucidated.

In summary, both the *in situ* microcosm and laboratory investigations confirmed anoxic degradation of MCB, providing proof of assimilation of MCB-derived carbon and mineralization of MCB. *In situ* microcosms provide a very promising tool to directly test *in situ* biodegradation of recalcitrant contaminants within reasonable time and this technique opens prospects for detailed analysis of the microbial key-players in future.

6.5.3 Isotope balance computation

CSIA of MCB and DCB was performed to gain information on *in situ* biodegradation within the plume and to investigate the relevance of degradation processes at the field site. Presence of these contaminants in the plume is presumably related to RDH of higher chlorinated benzenes and HCH which were initially spilled and which are still present inside the containment (SI, sections 1.1, 2.4; Anhang F).

The isotope signatures of MCB and DCB dissolved in the groundwater inside the containment were in the range of -27.7 to -29.7 ‰ and -24.0 to -27.5 ‰, respectively. In the plume DCBs showed a significant enrichment of ^{13}C compared to the containment and the isotope signatures were in the range of -25.5 to -15.6 ‰, -28.1 to -22.3 ‰ and -28.1 to -22.0 ‰ for 1,2-, 1,3- and 1,4-DCB, respectively (Tab. S4, Fig. S5; Anhang F). Assuming that due to their relatively low concentrations, RDH of TCB is not a relevant process within the plume and other sources for DCB production are lacking (SI, section 2.4; Anhang F), this high variability is supposedly related to microbial DCB degradation although no systematic correlation between concentrations and isotope signatures was observed. With increasing distance from the containment an enrichment of ^{13}C in the residual DCB fraction up to 4-5 ‰ compared to the source area was observed. Nevertheless, the highest isotopic shift (10 ‰ for 1,2-DCB and 4 ‰ for 1,3-DCB) was recorded in the area defined as contaminant source of the plume close to wells A, B, D suggesting that this part of the aquifer is probably favorable for RDH of DCB possibly associated with an accumulation of MCB. Further, our data supported the hypothesis of a preferential degradation of 1,2- over 1,3- and 1,4-DCB presumably due to higher yields in Gibbs free energy (Dolfing and Harrison, 1993) since lowest concentrations were determined for 1,2-DCB accompanied by the highest enrichment in ^{13}C and accordingly 1,4-DCB showed highest concentration and lowest isotope fractionation.

Under anoxic conditions RDH is the known degradation pathway for DCB leading to the formation of MCB, which consequently should be depleted in ^{13}C . Indeed in most of the wells the isotope signature of MCB was isotopically lighter compared to DCB. Theoretically, also a direct mineralization of DCB to CO_2 is feasible, but so far this has only been described for aerobic biodegradation (Van Agteren et al., 1998). The high abundance of MCB in the plume further indicated that apparently RDH of DCB and probably TCB led to the formation of MCB. Since no significant amounts of higher chlorinated benzenes were detectable in the plume, we suppose that the isotope signatures of DCB were only affected by its degradation.

The isotope signatures of MCB in the plume ranged between -29.6 ‰ and -25.3 ‰. Highest enrichments were found at the N and SE fringe of the plume (Fig. S5; Anhang F), where MCB concentrations are rather low. The ^{13}C -enrichment in combination with low MCB concentrations indicated MCB biodegradation in this aquifer zone. In the central and western part, presumably representing the former centerline of the plume, $\delta^{13}\text{C}$ values of MCB were more depleted indicating microbial production of MCB (Tab. S4, Fig. S5; Anhang F).

As previously suggested, two main mechanisms for MCB degradation should be considered: (i) RDH of MCB to benzene and its subsequent degradation or (ii) anaerobic oxidation of MCB which in this case would serve as electron donor (Braeckeveldt et al., 2007a; Nijenhuis et al., 2007; Nowak et al., 1996). Similar to RDH of TCB or anaerobic degradation of benzene, it is expected that microbial degradation of MCB under anoxic conditions would also lead to

significant isotope fractionation (Fischer et al., 2008; Griebler et al., 2004a; Mancini et al., 2003). At the anoxic aquifer of the study site the situation is more complex: MCB could be the product of DCB dehalogenation and simultaneously be subjected to microbial degradation. Both processes are associated with an opposite isotope fractionation. The RDH of DCB may result in isotopically lighter MCB and the MCB degradation would lead to an enrichment of ^{13}C in the residual MCB fraction. For this reason the isotope signature of MCB may reflect both processes, production of MCB from DCB as well as MCB biodegradation and the Rayleigh concept is not applicable straight forward to quantify biodegradation of MCB.

In order to investigate whether or not degradation of MCB occurred, an isotope mass balance including all CBs detected was performed based on two main assumptions: transformation of DCB to MCB via RDH which is most probably associated with isotope fractionation would not influence the cumulative isotope composition of all CB ($\delta^{13}\text{C}_{\text{CB}}$) and consequently only further MCB degradation would affect the isotope balance. Therefore significant enrichment of $\delta^{13}\text{C}_{\text{CB}}$ can be considered as indicator for MCB biodegradation. Regardless of the extent of isotope fractionation of individual contaminants upon formation and degradation, the cumulative isotope composition becomes enriched upon the degradation process. This assumption is valid even if a particular degradation reaction in the reductive sequence is not associated with significant isotope fractionation. However, at least one reaction step of the reduction sequence should be linked to isotope fractionation to document biodegradation. The isotope mass balance approach is corrupted, if alternative degradation pathways, for example mineralization of a DCB isomer without formation of MBC, occur in parallel and affect concentration or isotope signature of one member of the hypothetical sequence. In this case DCBs and MCB will not form a complete reaction sequence and the enrichment of individual species reflects degradation. However, isotope enrichment of DCBs will suggest *in situ* biodegradation in any case. In addition, the isotope enrichment of MCB at the plume fringes compared to the source areas will clearly indicate *in situ* degradation. Both, the mass balance approach as well as the individual evaluation of MCB may provide evidence that NA processes governing MCB concentration. In addition, different contaminants may vary significantly concerning their physico-chemical properties which could cause a preferential transport of contaminant species between source and down gradient the plume. For DCB and MCB no considerable differences concerning their retardation are expected, since adsorption coefficients for both are almost identical (Thullner and Schäfer, 1999).

The spatial distribution of concentration and respective carbon isotope signatures for the total CBs ($\delta^{13}\text{C}_{\text{CB}}$) is presented in Fig. 6-1. The $\delta^{13}\text{C}_{\text{CB}}$ values ranged between -27.2 ‰ up to -24.8 ‰. The lowest $\delta^{13}\text{C}_{\text{CB}}$ value was found for well A, located nearby the containment and defined as the centre of the plume due to highest contaminant concentrations. The decrease in total CB concentration was associated with significant enrichment of ^{13}C by ≥ 1 ‰ (Fig. 6-2)

indicating *in situ* biodegradation. Highest ^{13}C -enrichment of total CBs was mainly observed in the west (well O and N) and north (well J) of the plume suggesting removal of CBs from the system due to biodegradation (Fig. 6-1, Fig. 6-2). Overall, with the exception of well B and I, the data obtained from the isotope balance computation indicated a destructive removal of total CBs from the aquifer and demonstrated that chlorobenzene degradation is a relevant process in most parts of the plume. Abiotic processes such as dilution were generally not expected to cause considerable isotope effects, but they can influence the contaminant concentration (Fischer et al., 2006; Kopinke et al., 2005) (Fig. 6-2).




Fig. 6-1: Concentrations [$\mu\text{g L}^{-1}$] and carbon isotope signatures [%] of total chlorinated benzenes (numbers) at the study site for sampling in 2005. The letters indicate the name of the wells and the rectangle in diagonal crosses surrounds the area of the containment, which is hydraulically disconnected from the plume. Additionally the dashed (- - -) black line shows the area where the ratio between MCB and DCB is < 0.3 and the solid (—) line indicates where the ratio between MCB and DCB is > 0.9 . The symbols indicate the samples which belong to a common cluster as obtained by PCA: ▲-cluster I (A, B, C, D); ■-cluster II (G, H, N, O); ●-cluster III (E, F, I, J); ◆-cluster IV (J, M) (see Fig. 6-3). Wells which were not considered for PCA are indicated by ○.

The source zone (A, D, B) and the western fringe of the plume, where DCB is still present in higher amounts, were characterized by more enriched $\delta^{13}\text{C}_{\text{DCB}}$ and $\delta^{13}\text{C}_{\text{CB}}$ values suggesting microbial degradation of DCB leading to the formation of MCB more depleted in ^{13}C . MCB isotope signatures did not reflect MCB degradation in the western part of the plume. Nevertheless, the isotope balance of total CBs showed ^{13}C enrichment suggesting that

chlorobenzenes are destructively removed from the aquifer. At the SE fringe, higher $\delta^{13}\text{C}_{\text{MCB}}$ values directly indicated MCB degradation. DCB is not very abundant at this part of the aquifer and consequently significant formation of MCB due to DCB degradation is unlikely.




Fig. 6-2: Correlation between concentration [$\mu\text{g L}^{-1}$] and isotopic enrichment [%] of total chlorinated benzenes. The symbols indicate the sampling locations which belong to a common cluster as obtained by PCA: ▲-cluster I (A, B, C, D); ■-cluster II (G, H, N, O); ●-cluster III (E, F, I, J); ♦-cluster IV (J, M) (see Fig. 6-3).

6.5.4 Statistical analysis of the field data

Principal Component Analysis (PCA) was used to highlight relations between hydrogeochemical patterns of various sampling locations and to support delineation of prevailing biogeochemical processes at the contaminated groundwater system (Andrade et al., 2008; Lee et al., 2001).

PCA permitted to identify factors driving the separation between clusters and consequently highlighting geochemical variables associated with changes in the isotopic signatures. In the hydrogeochemical analysis of year 2005, the percent of variance within the data accounted for by the first (PC1) and second principal component (PC2) was high (55 %; Fig. 6-3). Scores of PC1 correlated positively to $\delta^{13}\text{C}_{\text{MCB}}$, total organic carbon (TOC), electric conductivity, ferrous iron, methane and $\delta^{13}\text{C}_{\text{CB}}$ (loadings > 0.3) and negatively to total CB and MCB concentrations. This emphasizes the significance of these variables in distinguishing clusters of sampling locations. Moreover, isotopic composition values of both MCB and CBs were negatively

correlated with their concentration values (Pearson's product-moment correlation, $P<0.05$) which supported the results of the isotope balance illustrating stable isotope fractionation as a function of microbial degradation. Sampling locations experiencing degradation activity (cluster II and IV) could be distinguished from apparently less active locations (cluster III) or locations where the activity could hardly be detected (cluster I). Overall, higher PC1 and PC2 scores indicated higher biodegradation activity in a well (Fig. 6-3).




Fig. 6-3: PCA ordination plot of geochemistry, contaminant concentrations, carbon isotope composition of MCB and total CBs in groundwater samples obtained during sampling in 2005. Values on the axes indicate % of total variation explanation by the corresponding axis (PC1, principal component axis 1; PC 2, principal component axis 2). The symbols indicate the samples which belong to a common cluster: ▲-cluster I (A, B, C, D); ■-cluster II (G, H, N, O); ●-cluster III (E, F, I, J); ♦-cluster IV (J, M). The figure in the upper right corner of the plot depicts the relative contribution of the descriptors in the reduced space. Description vectors correspond to: $\delta^{13}\text{C}_{\Sigma\text{CB}}$, isotope balance of total CBs; $\delta^{13}\text{C}_{\text{MCB}}$, carbon isotope signature of MCB; Fe^{2+} , ferrous iron; EC, electric conductivity; TOC, total organic carbon; CH_4 , methane; Cl^- , chloride; MCB, monochlorobenzene; ΣCB , total chlorobenzenes; NH_4^+ , ammonium; Eh, redox potential; SO_4^{2-} , sulphate; ΣMn , total manganese.

Interestingly, four clusters could be distinguished by PCA corresponding to geographical zonation: (i) heavily contaminated area (wells A, B, C, D), (ii) SW fringe (G, H, N, O), (iii) SE

fringe (wells E, F, I, L) and (iv) northern fringe of the plume (wells J, M) (Fig. 6-3, Fig. 6-1 for location of the wells). Cluster I representing the high contaminated region of the plume is consequently correlated with high contaminant concentrations. Cluster II grouped positively along PC2, which is associated with manganese, sulfate and $\delta^{13}\text{C}_{\text{CB}}$ emphasizing relatively higher values for these variables at the SW fringe. Although enrichment in ^{13}C indicating biodegradation has been observed in this region, a linkage between anaerobic oxidation and presence of sulfate or manganese could not be ruled out. Samples from the SE fringe of the plume (cluster III) grouped close to the origin and thus show intermediate hydrogeochemical patterns with no extreme values in the variables taken into account for PCA. Alternatively, sampling locations of the northern fringe (cluster IV) clustered positively along PC1 and displayed relatively higher values of ferrous iron, methane, TOC and electric conductivity. At this area, MCB is present > 60% of total CB and isotope enrichment of CB and MCB is likely to mirror MCB degradation. Because $\delta^{13}\text{C}_{\text{MCB}}$ values were positively correlated with ferrous iron, TOC and electric conductivity ($P<0.05$), a linkage between these geochemical indicators and degradation activity may exist. Indeed, TOC may serve as a source of electron donors to support RDH of CBs. Moreover, positive correlation of ^{13}C enriched MCB with increased ferrous iron concentrations may suggest anoxic oxidation of MCB during ferric iron reduction. Caution should be taken classifying sampling locations, as PCA directly reflects the limits of the methodology on which each variable integrated in the analysis has been derived. For instance, samples belonging to cluster I were collected in a highly contaminated part of the plume and showed no shift in $\delta^{13}\text{C}$ values of MCB or of total CBs. This could be used as indicator for recalcitrance to biodegradation. However, CSIA suggested DCB degradation due to relatively high isotopic enrichment accompanied by MCB formation (Tab. S4, Fig. S5, Anhang F). The significant DCB enrichment was masked in the isotope balance possibly due to the accumulation of relatively ^{13}C depleted MCB which may point to RDH of DCB. Moreover, the results of the *in situ* and laboratory tracer studies showed that *in situ* biodegradation of MCB is feasible under the ambient aquifer conditions, but could not be determined by means of CSIA. This emphasizes the interest of combining several tools of various sensitivity levels to gain evidence of *in situ* degradation.

Overall, the PCA results underlined the hydrogeochemical heterogeneity of the investigated field site. Delineated clusters of sampling locations showed various hydrogeochemical trends and corresponded to defined zones of the contamination plume, which putatively differ with respects to their potential and development of CB degradation activity. However, causality or mechanistic interpretation could often hardly be proved based on statistical analysis of the field data alone. Together with other lines of evidence, this delineation may be relevant in terms of risk assessment when coupling information about dominant contaminant flow patterns with zonation of potential biogeochemical characteristics at the plume scale.

To further confirm that monitored natural attenuation might be suitable as remediation method at the study site stability of the plume as well as of NA processes needs to be demonstrated (US-EPA, 1999). The comparative analysis of contaminant and isotope signature patterns of sampling locations in year 2005 and 2006 by PCA did not reveal substantial changes (Fig. S6, Anhang F). Additionally, the temporal evolution of contaminant concentration (2000-2007) and corresponding isotope signature (2005-2007) were analyzed in more detail for wells H, F, L at the southern fringe of the plume (Fig S7-9, Anhang F). In all three wells, concentrations varied significantly while the isotope signatures remained relatively stable over time. Although minor fluctuations within the data occurred, the constant enrichment of heavy isotopes compared to central parts of the plume indicated that the biodegradation potential was maintained over time demonstrating plume stability presumably controlled by microbial activity (see also SI, section 2.5; Anhang F).

6.5.5 Implications for environmental studies and site management in terms of natural attenuation

This study aimed at evaluating the efficacy of NA applying an integrated approach. Multivariate statistics permitted to delineate various geochemical zones within the plume associated with microbial degradation suggesting variability of processes involved in contaminant removal underlining the complexity and heterogeneity of the field site. PCA supported the results obtained by the isotope balance and enabled a combined analysis of geochemical and isotope data to evaluate the *in situ* biodegradation potential. To explore significant spatial and temporal changes of the hydrogeochemical patterns statistical methods are helpful and CSIA can reveal if biodegradation controls plume stability. Such methods can be easily implemented into a monitoring campaign. At the investigate area plume stability was demonstrated over time. Major changes in environmental conditions that may reduce the efficacy of the NA processes especially that of *in situ* biodegradation, are not likely to occur at the field site.

As basically required by authorities and scientists we presented an integrative concept encompassing multiple lines of evidence for NA of recalcitrant contaminants such as CBs at contaminated complex subsurface environments (McKelvie et al., 2007; US-EPA, 1999). At many field sites numerous contaminants are present and different mechanisms may govern the fate of pollutants *in situ*. Especially at such complex systems characterized by multiple contaminants, sequential degradation and/or different potential pathways one single method can not elucidate all relevant biogeochemical processes. Therefore, coupling of several techniques may lead to a more robust assessment and allows verifying plausibility of lines of evidence for *in situ* biodegradation processes. This further results in complex sets of data which can be treated more easily by statistical methods to interpret the relationship between certain parameters and to analyze the influence as well as temporal changes of

hydrogeochemical factors on *in situ* biodegradation without substantial additional monitoring costs (Andrade et al., 2008; Lee et al., 2001).

Even for recalcitrant contaminants (e.g. MCB) with unknown degradation pathways, an appropriate methodology may provide evidence of *in situ* biodegradation. An integrative approach consisting of hydrogeochemical analysis combined with novel stable isotope tools and multivariate statistics may substantially support identification of mechanisms and main biogeochemical factors associated with contaminant removal. At the complex chlorobenzene contaminated field site evaluation of NA potential showed that CBs are subjected to microbial degradation leading to sustainable removal of pollutants from the groundwater.

6.5.6 Acknowledgment

We thank the "Deutsche Bundesstiftung Umwelt" (grant 20004/751) for financial support of N. Stelzer. G. Imfeld is supported by a European Union Marie Curie Early Stage Training Fellowship (AXIOM, contract N° MEST-CT-2004-8332). We like to acknowledge A. Fischer and anonymous reviewers for critical reading of the manuscript. Further we are grateful to M. Gehre and U. Günther for technical assistance in the isotope laboratory. We also thank K. Sublette who provided the Bio-Sep® beads. The help of the Bauerumweltgruppe at the field site is acknowledged and we especially want to thank the site owner and the responsible environmental agency for good cooperation.

6.5.7 Supporting Information Available

Detailed description of the field site, sampling, experimental set up, analytical methods, hydrogeochemical data, chromatograms of *in situ* microcosm analysis, $\delta^{13}\text{C}_{\text{CO}_2}$ of laboratory enrichment cultures, isotope analysis of chlorinated benzenes, plume stability is available free of charge via the Internet at <http://pubs.acs.org> (Anhang F).

7 Assessment of *in situ* biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland

Mareike Braeckeveld, Hemal Rokadia, Gwenaël Imfeld, Nicole Stelzer, Heidrun Paschke, Peter Kuschk, Matthias Kästner, Hans-H. Richnow, Stefanie Weber

(This chapter was published in Environmental Pollution 148(2), p. 428-437)

7.1 Abstract

The degradation of monochlorobenzene (MCB) was assessed in a constructed wetland treating MCB contaminated groundwater using a detailed geochemical characterisation, stable isotope composition analysis and *in situ* microcosm experiments. A correlation between ferrous iron mobilisation, decreasing MCB concentration and enrichment in carbon isotope composition was visible at increasing distance from the inflow point, indicating biodegradation of MCB in the wetland. Additionally, *in situ* microcosm systems loaded with ¹³C-labelled MCB were deployed for the first time in sediments to investigate the biotransformation of MCB. Incorporation of ¹³C-labelled carbon derived from the MCB into bacterial fatty acids substantiated *in situ* degradation of MCB. The detection of ¹³C-labelled benzene indicated reductive dehalogenation of MCB. This integrated approach indicated the natural attenuation of the MCB in a wetland system. Further investigations are required to document and optimise the *in situ* biodegradation of MCB in constructed and natural wetland systems treating contaminated groundwater.

7.2 Capsule

An integrated approach including isotope composition analysis and *in situ* microcosm experiments provided evidences for *in situ* biodegradation of MCB in a wetland system.

7.3 Introduction

Monochlorobenzene (MCB) is encountered worldwide as a groundwater pollutant, and persists in the essentially anaerobic aquifer at the large-scale contaminated site in Bitterfeld, Germany (Heidrich et al., 2004b; Wycisk et al., 2004). In recent years, interest has grown in using phytoremediation processes for the elimination of recalcitrant organic substances from waste- and groundwater (Macek et al., 1998; Schnoor et al., 1995; Shimp et al., 1993; Trapp, 2000) including chloroaromatics (Gilbert and Crowley, 1997). Wetland systems represent an effective and inexpensive option to treat groundwater polluted with organic compounds by taking advantage of the geochemical and biological processes (e.g. Baker, 1998; Dunbabin and Bowmer, 1992; Gumbrecht, 1993). Indeed, rapid degradation of chlorinated organics has been observed in the rhizosphere (Anderson and Walton, 1995; Jordahl et al., 1997; Lorah and Olsen, 1999; Pardue et al., 1996).

While aerobic degradation of MCB has been well studied (Van Agteren et al., 1998), only some evidence for MCB transformation under anoxic conditions has been presented yet and the degradation pathway is unknown (Kaschl et al., 2005; Liang and Gribic-Galic, 1990; Nowak et al., 1996). Moreover, only very few studies focus on the anaerobic microbial transformation of MCB under field conditions. Recently, indications of anaerobic MCB degradation taking place in the Bitterfeld contaminated aquifer were provided on the basis of isotope fractionation patterns (Kaschl et al., 2005). Kinetic isotope fractionation processes have been employed to demonstrate the biological transformation of various contaminants (Lollar et al., 2001; Richnow et al., 2003a; Richnow et al., 2003b; Song et al., 2002). A substantial enrichment of ^{13}C in the non degraded fraction in the course of a contaminant plume indicates microbial degradation, as dilution and sorption do not affect the isotope composition of contaminants significantly (Harrington et al., 1999; Schüth et al., 2003; Slater et al., 2000). Combining stable isotope composition analysis with information obtained in simple *in situ* microcosm experiments (BACTRAPS) using isotope labelled substrate may provide a suitable approach to qualitatively support *in situ* biotransformation and to monitor spatial and temporal natural attenuation processes. Previously, BACTRAPS were exclusively installed in groundwater monitoring wells (Geyer et al., 2005; Kästner et al., 2006; Stelzer et al., 2006a) and were deployed in sediment for the first time in the framework of this study.

For the assessment of *in situ* biodegradation in constructed wetlands and wetlands treating contaminated groundwater, it may be necessary to use several methods providing more than one line of evidence. A combined approach may be of additional benefit in particular when systems are complex, possess several compartments and convincing evidence is required. Moreover, a better understanding of the controlling geochemical processes in wetland systems is necessary to reliably predict the retention and transformation of contaminant. In this study, we evaluated the natural attenuation of MCB in a constructed wetland treating MCB contaminated groundwater using a detailed geochemical characterisation, stable isotope composition analysis and *in situ* microcosm experiments. The spatial variations of geochemical parameter were studied with the help of multivariate statistics to investigate the main processes controlling the wetland system. The concentration and carbon stable isotope composition of MCB was analysed to monitor the *in situ* contaminant degradation and *in situ* microcosms were used to provide qualitative evidences of *in situ* biotransformation of MCB.

7.4 Materials and methods

7.4.1 Design and characteristics of the wetland

The pilot constructed wetland at the experimental site in Bitterfeld was set up in December 2002. The horizontal subsurface flow wetland consisted of a stainless steel tank divided into

two segments. Each segment was 6 m x 1 m and was filled to an average depth of 0.5 m with autochthonous quaternary aquifer material consisting predominantly of Bitterfeld mica sand (25%) and gravel (67%), which was embedded in lignite (10%) with an effective porosity of 28% (Vogt et al., 2002). The hydrogeochemical characteristics of the study site and the filling material originating from the local aquifer are described in previous studies (e.g. Vogt et al., 2002; Weiss et al., 2001). One of the segments was planted with common reed (*Phragmites australis*, Cav.), whereas the other side was left unplanted. In both segments, a 1 m long open water pond at the outflow side allows direct contact between the atmosphere and the water surface. The water level was maintained at approximately 10 cm below the surface of the wetland. The groundwater was collected from the MCB contaminated aquifer and conveyed from 16 to 22 m depth directly to the wetland. Both segments were operated in a flow-through mode at a flow rate of 4.7 L h⁻¹, corresponding to a retention time of 6 days.

7.4.2 Sampling

In the period from April to September 2005, pore water samples were collected five times (d0; d53; d66; d143; d172) in order to investigate the geochemical processes and the contaminant behaviour in the wetland system. The pore water was collected in both segments along a transect from the inflow up to the outflow of the wetland, at respectively 0 (inflow valves), 0.5, 1, 2, 3 and 4 m using a stainless steel lance. At each sampling point, three depths, 30, 40 and 50 cm were systematically investigated. Water samples were also collected at the ponds (6 m). In addition, to assess the *in situ* biodegradation using isotope composition analysis, pore water samples from both segments were collected at day 0 and day 53, at 0, 1 and 3.5 m along the wetland at 0.5 m depth, as well as in the ponds.

7.4.3 Physico-chemical and geochemical parameters of the pore water samples

The redox potential was measured on-line in the field using a SenTix ORP electrode (PT 1000, PreSens, Regensburg, Germany). The temperature was determined by a temperature sensor (PT 1000, PreSens, Regensburg, Germany). Samples for the pH analysis and the quantitative ions were filtered through a 5 µm syringe filter (Ministart NML, Sartorius) for particle removal. The pH value was measured with a SenTix41 electrode with pH 537 Microprocessor (WTW, Weilheim, Germany). Oxygen measurement was carried out using an optical oxygen trace sensor system (oxygen meter Fibox-3-trace and flow-through cell type sensor FTC-TOS7) with automatic temperature compensation (temperature sensor PT 1000) (PreSens, Regensburg, Germany). For the analysis of Mn(II), total iron and Fe(II), hydrochloric acid was added and samples were diluted with deionised water (1:10, v:v). Total iron and Mn(II) concentrations were analysed by atomic emission spectrometry with ICP excitation and CCD detection (Spectro Ciros Vision CCD, Spectro Analytical Instruments, Kleve, Germany). Photometric

analysis of ferric iron was carried out at 562 nm after derivatisation with ferrocin using a Cadas 100 photometer (Hach Lange, Düsseldorf, Germany). Chloride and sulphate concentrations were determined by ion chromatography (DX 500) with conductivity detection (CD 20) and a IonPacAG11 (4x250 mm) column (Dionex Corporation, Sunnyvale, USA). For the analysis of sulphide concentrations samples were spiked with sulphide anti-oxidant buffer (200 ml L⁻¹ 10 M NaOH, 35 g L⁻¹ ascorbic acid, 67 g L⁻¹ EDTA) (1:1, v:v) and measured with an ion selective Ag/S 500 electrode and reference electrode R 503 (WTW, Weilheim, Germany).

7.4.4 Analysis of benzene, MCB and metabolites

Pore water samples for the analysis of benzene and MCB concentrations were collected in 20 mL glass flasks (Supelco, Bellefonte, USA), and sealed with Teflon-lined septa. Sodium azide solution was added to the samples to inhibit microbial activity. Benzene and MCB concentrations were quantified by automatic headspace gas chromatography using an HP 6890 gas chromatograph with flame ionisation detector (Agilent technologies, Palo Alto, USA). For headspace analysis a volume of 1000 µl was injected at an injection temperature of 250°C with split 1:5 (measurements in duplicates). The chromatographic separation was achieved on a HP-1 capillary column (Agilent technologies, Palo Alto, USA) (30 m x 0.32 mm x 5 µm) with the following oven temperature program: 45°C (1 min), 20°C min⁻¹ to 200°C (2.5 min), 65°C min⁻¹ to 250°C (1 min) and a detector temperature of 280°C.

For the determination of the carbon isotope composition of MCB, 1 L glass bottles (Schott, Mainz, Germany) containing NaOH pellets to prevent microbial growth were filled completely with groundwater, stored at 4°C and extracted within 24 h using 2 mL n-pentane as described previously (Richnow et al., 2003b). The analysis of volatile metabolites obtained in the *in situ* microcosm experiments was carried out using a HP 6890 gas chromatograph with HP 5973 mass spectrometer (Agilent Technologies, Palo Alto, USA). Aliquots of 1 µl liquid samples were injected at a temperature of 280°C with split 1:40 and separated on a Zebron BPX-5 column (30 m x 0.32 mm x 0.25 µm) (Phenomenex, Torrance, USA). The following oven temperature program was applied: 40°C (2.5 min), 10°C min⁻¹ to 70°C (0 min), 60°C min⁻¹ to 280°C (4 min).

7.4.5 In situ microcosms (BACTRAPS)

Preparation of microcosms and derivatisation of fatty acids

The *in situ* microcosms were prepared as described previously (Stelzer et al., 2006a). Different sets of *in situ* microcosm experiments were prepared. One set was loaded with [¹³C₆]-labelled MCB (Cambridge Isotope Laboratories, Andover, USA) and another one with natural abundance MCB. A third set was kept unloaded to observe the background effects. The loading was done via gas phase under reduced pressure with approximately 40 mg MCB per g Bio-Sep®. The microcosms were deployed at 1.5, 2.5 and 4.5 m from the inflow in both planted

and unplanted segments at 50 cm depth. The microcosms were collected after 6 weeks and fatty acid extraction was carried out according to Bligh and Dyer (1959). The derivatisation to obtain fatty acid methyl esters (FAME) was done according to Thiel et al. (2001). After evaporation to complete dryness and addition of heneicosanoic acid methyl ester (C21:0) as an internal standard the FAME fraction was dissolved in *n*-hexane for further identification, structural characterisation and carbon isotope composition analysis.

GC-MS analysis

For identification and structural characterisation of FAME a HP 6890 gas chromatograph coupled with a HP 5973 quadrupole mass spectrometer (Agilent Technologies, Palo Alto, USA) was used. The FAMEs were separated on a Zebron BPX-5 column (30 m x 0.32 mm x 0.25 µm) (Phenomenex, Torrance, USA) with the following temperature program: 70°C (1 min), 20°C min⁻¹ to 130°C, 2°C min⁻¹ to 150°C (5 min), 2°C min⁻¹ to 165°C (5 min), 2°C min⁻¹ to 230°C, 20°C min⁻¹ to 300°C (5 min). FAME were identified by comparing with the retention time and mass spectra of an authentic standard mix (bacterial acid methyl esters mix, Sigma-Aldrich, Germany) and quantified relatively to the internal standard.

Isotopic composition analysis

The carbon isotope composition of MCB and the FAMEs was measured with a gas chromatography-combustion-isotope ratio mass spectrometry system (GC-C-IRMS) consisting of a GC unit (HP 6890, Agilent technologies, Palo Alto, USA), a combustion device (Finnigan MAT GC III, ThermoFinnigan Bremen Germany) with water-removal assembly (Nafion® membrane, 50 cm long, T = 0°C) and a mass spectrometer (Finnigan MAT 252; ThermoFinnigan, Bremen, Germany), as previously described (Richnow et al., 2003a). Helium was used as carrier gas at a flow rate of 1.5 ml min⁻¹.

Stable isotope samples were measured in triplicates and the analyses were carried out immediately after each sampling. Aliquots of 1 µl liquid samples were injected at 250°C with split 1:10 to the GC-C-IRMS and separated on a capillary column (Zebtron ZB-1, 60 m x 0.32 m x 1 µm; Phenomenex, Torrance, USA). The following chromatographic conditions were applied: injector temperature 250°C, oven temperature program: 40°C (1 min), 4°C min⁻¹ to 150°C, 20°C min⁻¹ to 250°C (2 min). The carbon isotope composition is reported in the delta notation as δ¹³C values [‰] relative to Vienna Pee Dee Belemnite standard (V-PDB, IAEA-Vienna) (Eq. 7-1) (Hoefs, 1997).

$$(Equation\ 7-1)\qquad \delta^{13}\text{C}[\text{\%}] = \left(\frac{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{Sample}}}{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{Standard}}} - 1 \right) * 1000 = \left(\frac{R_{\text{Sample}}}{R_{\text{Standard}}} - 1 \right) * 1000$$

Eq.(7-2) is applied to calculate the remaining substrate fraction (f_t) using the isotope fractionation factor (α_C).

$$(Equation \ 7-2) \quad f_t = \left(\frac{R_t}{R_0} \right)^{\left(\frac{1}{\alpha_C - 1} \right)}$$

The R_t and R_0 give the isotope composition of MCB at time t and zero. Eq. (7-3) is applied to calculate the percentage of biodegradation of the residual substrate fraction (B_t).

$$(Equation \ 7-3) \quad B[\%] = (1 - f_t) * 100$$

For the separation of the FAME fractions, a Restek RTX 5 column (60 m x 0.32 mm x 0.1 µm; Restek, Bellefonte, USA) was used with the same temperature program applied for the GC-MS-analysis of FAME. Aliquots of 1 µl were injected with split 1:5. The methylation of fatty acids for gas chromatographic analysis introduces an additional carbon atom into the structure of the fatty acid molecules which affects its isotopic composition. Therefore the isotope signature of fatty acids ($\delta^{13}\text{C}_{\text{FA}}$) was corrected for the isotope effect upon derivatisation to FAME with methanol as described previously (Abraham et al., 1998; Abrajano et al., 1994; Goodmann and Brenna, 1992). The methanol used for the derivatisation had an isotope composition of -38.2‰.

7.4.6 Statistical analysis

Statistical analyses were carried out using the R software (R, Version 2.1.1, 2005). Statistical significance of the difference in geochemical parameters as well as concentrations and isotopic compositions of MCB between the planted and unplanted segment was determined with the unpaired Wilcoxon or the Kruskall-Wallis rank sum tests. Correlation analyses were carried out using the Spearman rank sum coefficient. Principal Component Analysis (PCA) was used to analyse the relationship between the different samples with reference to their respective pore water parameters. The sampling location corresponds to the object and the chemical parameters to the descriptors (represented by the vectors) of the multivariate analysis. The PCA were scaled as correlation biplots.

7.5 Results

7.5.1 Characterisation of pore water chemistry

Distribution of MCB and benzene

The MCB concentration was measured as a function of the distance from the inflow point in both the planted and unplanted segment (Tab. 7-1). The average amount of MCB ranged from

Assessment of in situ biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland

14.4 mg L⁻¹ to 17.7 mg L⁻¹ at the inflow down to 2.0 mg L⁻¹ to 2.2 mg L⁻¹ in the ponds for the planted and the unplanted segment, respectively. No significant difference in MCB concentration between the three depths over the study period were generally observed ($p<0.05$). Benzene was found in low concentration in both segments (<26 µg L⁻¹), with generally higher concentration values in the unplanted segment (Tab. 7-1).

Tab. 7-1: Geochemical characterisation, MCB and benzene concentrations of samples collected from (a) unplanted plot and (b) planted plot. Values represent the depth profiles average in the soil compartments (0.5 - 4 m) and average (0 m and pond) over the study period (13.04.05-29.09.05). Standard deviation is indicated in parentheses.

Sampling point [m from inflow]	pH	Temp. [°C]	O ₂ [mg L ⁻¹]	Eh [mV]	Fe(II) [mg L ⁻¹]	Sulphate [mg L ⁻¹]	Cl ⁻ [mg L ⁻¹]	Benzene [µg L ⁻¹]	MCB [mg L ⁻¹]
(a)									
0	7.1 (3.6) ^d	14.8 (8.5) ^d	0.04 (0.02) ^e	38 (37) ^d	0.18 (0.23) ^c	969 (402) ^c	280 (116) ^c	25.6 (3.6) ^c	17.7 (8.0) ^c
0.5	7.0 (2.2) ^b	16.3 (5.4) ^a	0.05 (0.04) ^b	61 (26) ^b	2.43 (1.83) ^a	969 (253) ^a	273 (69) ^a	21.2 (6.1) ^a	14.6 (5.0) ^a
1	6.7 (2.1) ^a	17.1 (5.4) ^a	0.04 (0.02) ^a	84 (38) ^a	9.58 (5.80) ^a	967 (253) ^a	269 (68) ^a	23.3 (3.9) ^a	14.5 (5.1) ^a
2	6.9 (2.0) ^a	17.8 (5.5) ^a	0.04 (0.04) ^a	70 (41) ^a	12.40 (6.74) ^a	973 (254) ^a	267 (67) ^a	22.2 (5.4) ^a	14.5 (5.2) ^a
3	6.8 (1.9) ^a	18.4 (5.5) ^a	0.04 (0.03) ^a	57 (34) ^a	21.12 (7.45) ^a	974 (254) ^a	261 (71) ^a	22.5 (5.5) ^a	14.9 (5.6) ^a
4	6.8 (1.9) ^a	18.9 (5.6) ^a	0.04 (0.02) ^a	57 (37) ^a	30.23 (7.08) ^a	973 (256) ^a	267 (72) ^a	21.1 (6.3) ^a	14.4 (5.5) ^a
Pond	6.8 (3.0) ^d	20.9 (8.4) ^d	n.a.	250 (44) ^d	3.06 (8.81) ^c	986 (403) ^c	287 (118) ^c	1.8 (3.0) ^c	2.0 (7.9) ^c
(b)									
0	6.8 (3.1) ^d	14.8 (8.5) ^d	0.04 (0.02) ^e	35 (40) ^d	0.20 (0.21) ^c	977 (400) ^c	262 (109) ^c	24.0 (6.9) ^c	14.4 (6.4) ^c
0.5	6.8 (2.2) ^b	15.6 (5.5) ^a	0.13 (0.05) ^b	47 (42) ^b	5.73 (3.62) ^a	994 (251) ^a	257 (68) ^a	11.8 (5.5) ^a	8.7 (7.7) ^a
1	6.8 (2.1) ^a	16.6 (5.4) ^a	0.12 (0.09) ^a	54 (40) ^a	13.34 (7.48) ^a	980 (248) ^a	256 (68) ^a	15.7 (6.6) ^a	9.8 (7.2) ^a
2	6.7 (2.0) ^a	17.6 (5.4) ^a	0.05 (0.02) ^a	46 (36) ^a	25.48 (7.60) ^a	1006 (248) ^a	266 (68) ^a	11.9 (8.8) ^a	8.2 (6.6) ^a
3	6.7 (1.9) ^a	18.0 (5.4) ^a	0.05 (0.03) ^a	51 (34) ^a	36.57 (7.33) ^a	1000 (248) ^a	271 (69) ^a	11.3 (8.9) ^a	7.9 (6.1) ^a
4	6.5 (1.9) ^a	18.8 (5.5) ^a	0.03 (0.02) ^a	57 (34) ^a	37.40 (8.94) ^a	998 (259) ^a	266 (74) ^a	10.2 (8.0) ^a	7.7 (5.9) ^a
Pond	6.8 (3.1) ^d	20.5 (8.1) ^d	4.85 (1.89) ^e	204 (43) ^d	2.42 (9.28) ^c	1068 (397) ^c	275 (111) ^c	0.2 (0.4) ^c	2.2 (7.8) ^c

^aProfile average of five sampling dates (n=15)

^b Profile average of four sampling dates (n=12)

^c Average of five sampling dates (n=5)

^d Average of four sampling dates (n=4)

^e Average of three sampling dates (n=3)

n.a. not assessed

Pore water geochemistry

The evolution of oxygen, redox potential, manganese, sulphide, sulphate, ferric iron and total iron were monitored along the flow path at three depths in order to characterise the geochemical conditions prevailing in the wetland system. The average values of the three depths investigated at each sampling point (0, 0.5, 1, 2, 3, 4 m from the inflow) were computed (Tab. 7-1). To investigate the differences between the sampling locations and to explore existing gradients, the data sets were analysed by principle component analysis, separately for the unplanted and the planted segments (Fig. 7-1). In both cases, the vectors representing the temperature, Fe(II), and total Fe parameters were orientated in the same direction (positive correlation), in the opposite direction with regards to vectors representing manganese and

ammonium (negative correlation), and are perpendicular to the MCB vector (absence of correlation). The orthogonal projection of an object on a descriptor allows approximating the correlation between that object and the descriptor. For both segments, samples from the inflow part of the wetland (0.5 to 1 m) were associated with MCB, manganese and ammonium, whereas samples from the outflow (3 and 4 m) were generally associated with total Fe and Fe(II). Indeed, the Fe(II) concentration at the inflow were below 0.5 mg L⁻¹ and systematically increased along the flow path in both planted and unplanted segment, indicating ferrous iron mobilisation. A clear shift in the parameters characterising predominantly the trends of variation of the samples along the flow path is operating on the first principal component. Indeed, the axis I corresponds to a spatial gradient from the inflow to the outflow of the system, and separates sampling sites accordingly (from the right to the left). The variations of geochemistry along the flow path contributed more in characterising the samples than the variations occurring along the vertical profile.

The maximum concentration of Fe(II) reached an average value of 37.4 mg L⁻¹ at 4 m from the inflow of the planted segment. The levels of total dissolved iron detected throughout the experimental period ranged from 0.8 to 50.6 mg L⁻¹. Interestingly, Fe(II) mobilisation was more important in the planted than in the unplanted segment. Correlation analysis revealed, however, very similar Fe(II) mobilisation patterns in both segments over the study period ($p>0.9$; $p<0.01$). Iron reduction was apparently a relevant process at the three depths. Due to the high background concentration of sulphate in the supplied groundwater (700-1100 mg L⁻¹), a sensitive analysis of the microbial sulphate reduction on the basis of sulphate concentration was not possible. However, trace concentrations of sulphide (<3 µg L⁻¹) were detected along the horizontal transect, indicating sulphate reduction. Mn(II) concentrations systematically ranged below 0.2 mg L⁻¹, suggesting a low relevance of Mn as electron acceptor in the wetland (data not shown). The redox potential displayed averages of values ranging from 35 mV at the inflow to 250 mV in the ponds, and maximal concentration of oxygen in the soil compartments was 0.12 mg L⁻¹, indicating the prevalence of anoxic conditions in the soil compartments and oxic conditions in the ponds, respectively. The dissolved organic and total carbon concentrations ranged from 0.24 to 0.45 mM L⁻¹ and from 2.27 to 2.41 mM L⁻¹ in the unplanted and planted segments, respectively (data not shown). Nitrate concentrations were assessed in previous studies and ranged systematically under the detection threshold in the groundwater supplied to the wetland (Vogt et al., 2002), and nitrate may therefore not represent a relevant electron acceptor. The chloride concentrations did not show any significant variation along the transect.




Fig. 7-1: Ordination plot generated by principal component analysis representing the relationship between the sampling locations and the average geochemical parameters measured in both the planted (a) and the unplanted (b) experimental wetlands over the study period. Description vectors correspond to: Fell = ferric iron; Tot.Fe = Total iron ($Fe(II)+Fe(III)$); MCB = Monochlorobenzene; Mn^{II} = Manganese(II); NH_4^+ = Ammonium; T = Temperature. Objects correspond to: 0.5 to 4 = distance [m] from the inflow point; u = upper depth (30cm); m = medium depth (40cm); l = lower depth (50cm). Values on the axes indicate % of total variation explained by the axes. PC 1 = principal component axis 1; PC 2 = principal component axis 2.

7.5.2 Carbon isotopic composition of MCB

The concentration and isotopic composition ($\delta^{13}C$) of MCB were plotted as a function of the distance from the inflow point (Fig. 7-2). A decrease in MCB concentrations over the flow path was

systematically associated in both segments with a significant enrichment in $\delta^{13}\text{C}$. In the planted segment, MCB showed a maximal isotope shift of 0.6‰, whereas it reached 0.9‰ in the unplanted segment. This indicates that MCB is subjected to *in situ* biodegradation in both segments. Assuming an isotopically homogenous source of MCB, these values are slightly higher than the typically defined analytical error of 0.5‰ associated with compound specific isotope analysis (Dempster et al., 1997; Mancini et al., 2002).




Fig. 7-2: Concentration and isotopic composition of MCB (diamond = concentration; squares = isotopic composition) in both planted (a) and unplanted (b) segments and for both sampling dates (line = 14.04.2005, dashed = 09.05.2005). Error bars show the standard deviation.

A quantitative assessment of *in situ* MCB biodegradation requires a fractionation factor representing the *in situ* conditions. Fractionation factors for aerobic MCB degradation are available (Kaschl et al., 2005), whereas factors retrieved under laboratory conditions for anaerobic MCB degradation are missing. Although MCB degradation under anaerobic conditions is expected in the wetland, Eq. (7-3) allows estimating the significance of

degradation along the flow path throughout the wetland applying the highest isotope fractionation factors (α C) retrieved for aerobic MCB degradation by a dioxygenase reaction pathway (α C = 1.0004) (Kaschl et al., 2005). The quantification was carried out based on the isotope signature of MCB measured at the planted side at day 0. The lowest $\delta^{13}\text{C}$ value measured (inflow point, 0 m: MCB $\delta^{13}\text{C} = -27.0 \text{ ‰}$) was used as the initial isotope composition of the source (R_0). The estimated percentage of biodegradation was 60% of the inflowing MCB mass at 3.5 m, whereas the observed contaminant mass decrease reached only 38% at 4 m from the inflow. Processing of isotope and concentration data of the second sampling campaign gave almost identical results.

The use of too low fractionation factors for the quantification of MCB degradation would lead to an overestimation of the effective MCB mass depletion at the planted segment. The observed absence of significant MCB mass depletions along with a higher isotopic shift at the unplanted segment obviously direct to the same conclusion. Therefore, higher fractionation factors are expected, which would be more in concordance with the observed MCB concentration values. The use of higher isotope fractionation factors would point to a dominating anaerobic fractionation process, which is compatible with the observation of anoxic conditions in the wetland. The fractionation factors would then be comparable to the ones retrieved by Kaschl et al. 2005 (Kaschl et al., 2005) (α C = 1.0005) in the local anaerobic aquifer with MCB contamination or by Griebler et al. (2004a) and Mancini et al. (2003), obtaining significantly higher fractionation factors for the anaerobic degradation of benzene or trichlorobenzenes.

7.5.3 *In situ* microcosm experiment

[$^{13}\text{C}_6$] labelled MCB was used as stable isotope tracer in the *in situ* microcosm experiment in the wetland. The BACTRAPs were incubated directly in both soil compartments, at several distances from the inflow, as well as in the ponds. The incorporation of labelled carbon derived from the labelled substrate into bacterial fatty acids provided evidence for MCB degradation in the wetland system. In addition, [$^{13}\text{C}_6$] benzene, a possible intermediate, found on the BACTRAPs from both segments demonstrated the occurrence of reductive dehalogenation of MCB (Fig. 7-3). Even though the employed [$^{13}\text{C}_6$] MCB contained 0.02 % of [$^{13}\text{C}_6$] benzene as impurity, significantly higher amounts of [$^{13}\text{C}_6$] benzene were detected by GC-MS.




Fig. 7-3: GC-MS chromatogram (total ion current) of the *in situ* microcosm extract showing the ^{13}C -labelled benzene metabolite at retention time of 1.73 min and corresponding mass spectrum (a) with the molecule mass of $m/z = 84$ for [$^{13}\text{C}_6$] benzene and $m/z 78$ for [^{12}C] benzene (b).

Fatty acid composition

The composition of total fatty acid fractions extracted from *in situ* microcosms was compared to investigate variation in the microbial community. No systematic differences were observed between the oxic ponds and the anoxic segments of the wetland. However, the lowest quantity of fatty acids was retrieved from the microcosms exposed at 4.5 m in the soil compartments, and the highest accumulated biomass was retrieved from the microcosms deployed in the pond of the unplanted side of the wetland (Fig. 7-4).

The fatty acid patterns were dominated by high amounts of the saturated hexa- (C16:0) and octadecanoic (18:0) and the monounsaturated hexa- (C16:1) and octadecenoic (C18:1) acids which are common fatty acids in bacteria. The tetra- (C14:0), pentadecanoic (C15:0) and eicosanoic (C20:0) acids, the *iso* and *anteiso* isomers of C15:0 as well as the unsaturated C18:2 were present in lower abundance and could not be detected in all samples (Fig. 7-4).




Fig. 7-4: Absolute abundance of extracted fatty acids [μg FA per gram Bio-Sep® beads] from *in situ* microcosms exposed in the soil compartment at different distance from the inflow and ponds at the unplanted (a) and planted (b) segments of the constructed wetland.

The variation of the geochemical parameters down gradient of the inflow were not related to a distinct change in fatty acid composition in both segments. No systematic differences in the fatty acid patterns were found between the planted and unplanted segments. Globally, no clear indications of microbial community changes were obtained on the basis of the fatty acid composition. In this experiment, fatty acid patterns were probably not sensitive enough to reflect changes in the microbial communities as a function of variation in geochemical conditions within the constructed wetland.

Isotope signatures of fatty acids

The total lipid fatty acids extracted from the *in situ* microcosms displayed some differences in the incorporation of ^{13}C into fatty acids. This was particularly obvious when comparing the

samples from the soil compartments and the ponds. The isotope composition of fatty acids ($\delta^{13}\text{C}_{\text{FA}}$) extracted from the BACTRAPs amended with [$^{13}\text{C}_6$] MCB ranged between -39‰ and 7244‰ (Tab. 7-2). An enriched ^{13}C signature of fatty acids (> 0 ‰) can only stem from the microbial utilisation of [$^{13}\text{C}_6$] MCB as a carbon source. In contrast, fatty acids with an isotope signature lower than -20 ‰ showed the typical natural abundance of $\delta^{13}\text{C}_{\text{FA}}$ found in soil and aquifer material (Pelz et al., 2001b; Pombo et al., 2002). Fatty acids derived from parallel microcosms with non labelled MCB or from non amended *in situ* microcosms displayed an isotope signature ranging from -24‰ to -54‰ (data not shown). This represents a typical isotope composition of lipids derived from microorganisms feeding on organic substrates with natural isotope composition.

Tab. 7-2: Carbon isotope composition of fatty acids extracted from *in situ* microcosms incubated for 6 weeks with $^{13}\text{C}_6$ labelled monochlorobenzene at 1.5, 2.5 and 4.5 m from the inflow point as well as at the pond.

[m] from inflow	Planted $\delta^{13}\text{C}_{\text{FA}} [\text{\textperthousand}]$				Unplanted $\delta^{13}\text{C}_{\text{FA}} [\text{\textperthousand}]$			
	1.5	2.5	4.5	Pond	1.5	2.5	4.5	Pond
C13:0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	118
C14:0	-6	40	n.d.	n.d.	138	204	116	238
iC15:0	121	89	n.d.	n.d.	n.d.	742	n.d.	1597
aC15:0	23	68	n.d.	n.d.	n.d.	3	n.d.	n.d.
C15:0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	-37	252
C16:1	411	453	n.d.	7244	n.d.	1832	n.d.	806
C16:0	2	9	93	1711	260	95	192	639
C17:0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	242
C18:2	-31	-30	n.d.	65	n.d.	-27	-30	216
C18:1	-30	-29	-28	276	-23	-23	-24	175
C18:0	-27	-28	-24	449	-22	-27	-26	266
C20:0	-37	-39	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

n.d. – not detected

Generally, all fatty acids extracted from the *in situ* microcosms exposed in the ponds showed enrichment in ^{13}C (65‰ to 7244‰), which was generally of higher intensity than in the soil samples (-39‰ to 1832‰). In contrast, only fatty acids with up to 16 carbon atoms showed ^{13}C incorporation in the soil samples with highest enrichment in C16 species. Fatty acids with longer carbon chains such as C18:0 or C18:1 displayed no enrichment in ^{13}C (-22‰ to -39‰), indicating that the microbial community was not exclusively growing on the [$^{13}\text{C}_6$] MCB. Comparing the samples from the planted and unplanted segments no significant differences in

$\delta^{13}\text{C}_{\text{FA}}$ were observed. Therefore, the putative impact of plants on the microbial community involved in MCB degradation could not be assessed using our test system.

Some fatty acids can be used as biomarker to identify specific groups of microorganisms (Kaur et al., 2005; Zelles, 1999). In some of the samples labelled *iso* and *anteiso* branched fatty acids with 15 carbon atoms could be identified, indicating that Gram-positive bacteria were involved in the biodegradation of MCB. Labelled C18:2 was only found in samples of the ponds. Linoleic acids (C18:2) can serve as a biomarker for fungi or other eukaryotic organisms (Lösel, 1988) and their presence may lead to the hypothesis that grazing organisms, not involved in the biodegradation of MCB, may feed on the microbial biofilm. However, the applied GC-MS procedure did not allow conclusive identification of the position of the double bond.

7.6 Discussion

The geochemical parameters indicated the overall prevalence of anoxic conditions associated with iron mobilisation in the soil parts of the wetland, whereas an aerobic milieu characterised the ponds. In the *in situ* microcosm experiments, the level of incorporation of labelled carbon into bacterial biomass was used as direct indicator of *in situ* MCB degradation. Interestingly, the analysis of the BACTRAPs incubated in the ponds revealed fatty acids patterns and ^{13}C incorporation levels differing from the ones retrieved from the soil compartments. The higher ^{13}C incorporation level observed in the ponds of both segments is indicative of a more effective microbial transformation of the [$^{13}\text{C}_6$] MCB under the prevailing aerobic conditions. Along with a significant accumulation of biomass on the microcosms retrieved from the ponds, these results suggest that some change in the microbial community dynamics may operate between the anoxic soil compartments and the more aerobic ponds. Moreover, the fact that all the extracted fatty acids showed incorporation in ^{13}C suggests that the microbial community established on the microcosms were mostly involved in contaminant degradation. However, it should be considered that cross feeding by metabolites or recycling of dead biomass within the biofilm may also channel labelled carbon into individual members of the microbial community, explaining the variation in the labelling of specific fatty acids. In general, fatty acids displaying a higher incorporation of ^{13}C were very likely synthesised by organisms feeding on [$^{13}\text{C}_6$] MCB, whereas organisms synthesising non-labelled fatty acids were likely not involved in the degradation of the labelled MCB and use different carbon sources. The analysis of the composition of total fatty acid fractions showed that this method might not be sensitive enough for investigating in detail changes in microbial communities between soil and water compartments as well as down gradient the flow path.

Additionally, the MCB degradation processes in the wetland were investigated by carbon stable isotope composition analysis. A correlation between decreasing MCB concentration and a shift in the carbon isotope signature towards the heavier isotope was visible along the flow path,

suggesting the degradation of MCB. Toxic effects of MCB on the MCB-degrading population can be reasonably excluded at the observed range of concentration values (Fritz et al., 1992; Vogt et al., 2002; Vogt et al., 2004).

In the transition from the soil compartments to the ponds, a substantial contaminant mass depletion without concomitant isotope enrichment was observed. In this open system, it is likely that part of the contaminant may partition into the atmosphere, affecting the MCB concentration values without generating a significant isotope shift. Moreover, under oxic-prevailing conditions, several bacteria have the ability to use MCB as sole carbon and energy source, and may putatively adopt the well-known and described aerobic degradation pathways (Van Agteren et al., 1998). These bacteria may degrade MCB at faster rate than the degrading bacteria associated with anoxic conditions, contributing to the observed contaminant mass decrease. These oxygen-driven degradation reactions would lead to MCB mass decrease without a significant associated isotope effect (Kaschl et al., 2005). In parallel, biogeochemical processes such as oxidation of ferrous iron or mineral surfaces may compete for oxygen (Ehrlich, 1998; Søgaard et al., 2001; Warren and Haack, 2001), leading to transient conditions and oxygen gradients, which may affect the composition of the existing microbial community and rate of degradation reactions. However, the anaerobic degradation pathway of MCB is not elucidated yet.

In the anoxic soil compartments, two major hypothetical degradation pathways would come into consideration: 1) reductive dechlorination of MCB, and 2) degradation of MCB as an electron donor molecule. First, an initial dechlorination of MCB, followed by the degradation of benzene, is likely to occur in the soil compartments. Indeed, the presence of ¹³C-labelled benzene detected in all the microcosms along with the detection of low benzene concentrations suggested that MCB is degraded reductively to benzene under anoxic conditions prior mineralisation. This pathway would suggest a similar pattern as observed previously by Nowak et al. (1996). The pH values were, however, constant, and the relatively high background level of chloride in the groundwater hindered the direct verification of MCB degradation by an increasing chloride concentration along the flow path. Bacterial breakdown of benzene could lead to the formation of benzoate or phenol as intermediates (Chakraborty and Coates, 2005; Edwards and Grbic-Galic, 1992; Lovley, 2000; Phelps et al., 2001; Ulrich et al., 2005), which were not detected in this study. Soluble organic carbon species, which could serve as electron donors for dechlorination processes, were found in low concentration in the unplanted segment, which might effectively limit the extent of degradation. Conversely, dechlorination activity in the planted segment may be partly related to the abundance of hydrogen and reduced organic acids such as acetate and propionate (Holliger et al., 1992; Middeldorp et al., 1997).

Reductive dechlorination reaction of MCB to benzene is expected to be associated with a pronounced primary isotope effect (Griebler et al., 2004a). In contrast, the isotope composition presented in this study displayed a slight but significant enrichment ranging from 0.4 to 0.7 δ units, in the soil compartments of the planted and the unplanted segments, respectively. However, the isotope effect at zones of preferential *in situ* degradation reaction can be substantially higher. If other non fractionating processes such as sorption and volatilisation also contribute to a decrease in MCB concentrations, the isotope effect upon *in situ* degradation is expected to be relatively high. The observed isotope effect points to dominating anaerobic processes, as inferred by the estimation of biodegradation levels over the flow path. In the planted, the oxygen supplied by the plant at the rizosphere level may favour the establishment of aerobic zones. Although this process is not relevant for the electron budget, oxygen may contribute to the MCB degradation reactions, leading to MCB decrease without concomitant isotope effect. A fractionating anaerobic process and a less fractionating aerobic process may both contribute to *in situ* degradation, resulting in a mixed overall fractionation at the planted segment. Conversely, the isotopic composition shift observed at the unplanted segment suggests the occurrence of a more fractionating process.

Alternatively, MCB may be degraded as an electron donor molecule under ferric iron- or sulphate-reducing conditions. Anaerobic oxidation of benzene under these conditions has been previously observed (Anderson and Lovley, 2000; Anderson et al., 1998). For instance, geochemical footprints of iron reduction processes were found, and the ferrous iron mobilisation was increasing as a function of the flow path. Dissolved Fe in pore waters can be a result of different processes such as Fe(III) reduction (Lovley, 1991; Lovley, 1997), pyrite oxidation (Lord and Church, 1983), or Fe complexation (Luther et al., 1996). Fe(II) may be precipitated with sulphide originating from sulphate reduction activity or form complexes. Therefore the concentrations may not reflect the true extent of iron reduction in the presence of sulphate reduction. A low extent of sulphate reduction process and the availability of reactive iron may prevent accumulation of H₂S in the near-neutral conditions of the wetland. A MCB mineralisation by sulphate reduction contributing to the contaminant mass decrease is feasible and may be possible. However, due to the high concentrations of sulphate in the supplied water, a reliable estimation of the extent of sulphate reduction could not be carried out. However, a reduction of ferric iron directly linked stoichiometrically to MCB oxidation in the wetland may theoretically account for a MCB mass decrease of about 18% between the inflow and the outflow of the system at both the planted and unplanted segment.

Additionally, other unknown degradation pathways can not be excluded. These hypotheses will require further detailed investigations, along with further isolation and identification of microorganisms involved in the anaerobic MCB degradation in wetlands treating MCB contaminated water.

7.7 Conclusion

The integrated approach provided evidence for *in situ* MCB biodegradation in both, soil compartments and ponds of the planted and unplanted segments of a horizontal subsurface flow constructed wetland. This was supported by isotopic fractionation analysis, combined with *in situ* microcosm experiments, which can be utilized to document further the *in situ* degradation of MCB and other contaminants in wetland systems. Further investigations to elucidate the microbial degradation of MCB, facilitated by an integrated approach and combined with a high resolution sampling, are required to evaluate zones of enhanced *in situ* biodegradation of MCB and to optimise wetland systems treating contaminated groundwater.

7.8 Acknowledgements

The Department of Groundwater Remediation, the SAFIRA Project, the ANANAS Project and the Department of Analytical Chemistry of the UFZ are acknowledged for assistance in the field and laboratory work. We are grateful to S. Täglich, J. Ahlheim, O. Thiel, G. Mirschel, I. Mäusezahl and T. Nullmeyer. We are thankful to Dr. A. Miltner, Dr. M. Gehre, U. Günther, K. Ethner, K. Puschendorf, and A. Fischer, for their technical support in our laboratory and isotope measurements. Dr. Kerry Sublette, University of Tulsa, USA provided the Bio-Sep® beads for this work. We thank the EU Marie Curie Host Fellowships for Early Stage Training Project "AXIOM", the Virtual Institute for Isotope Biogeochemistry (VIBE) and the Deutsche Bundesstiftung Umwelt (20004/751) for financial support.

8 Zusammenfassung

Die Untersuchungen, die im Rahmen dieser Promotion durchgeführt wurden, stehen im Kontext von *Natural Attenuation* (NA), den natürlichen Abbau- und Rückhalteeigenschaften eines Grundwasserleiters, die immer häufiger zur Überwachung und Sicherung kontaminierte Standorte genutzt werden. Da insbesondere der mikrobielle Schadstoffabbau zu einer destruktiven und vor allem nachhaltigen Reduzierung der Schadstoffmengen im Aquifer beiträgt, ist der Nachweis des biologischen Abbaus von Grundwasserkontaminanten ein wesentliches Element für die Implementierung von NA als Standortsicherungskonzept. Es gibt nur wenige Methoden, die einen direkten Nachweis der Biodegradation unter *in situ* Bedingungen ermöglichen. Inhalt der vorliegenden Dissertation ist daher die Entwicklung eines Testsystems (*in situ* Mikrokosmen, BACTRAP[®]) zum direkten Nachweis des mikrobiellen Schadstoffabbaus im kontaminierten Grundwasser sowie die Integration des Verfahrens in Monitoringstrategien für NA-Konzepte.

8.1 Beurteilung der *in situ* Mikrokosmen zum Nachweis der Biodegradation

Mit der Entwicklung von *in situ* Mikrokosmen (BACTRAP[®]) steht neben der substratspezifischen Isotopenfraktionierungsanalyse (CSIA) und *in situ* Tracerversuchen entlang einer Fließstrecke oder in Form von Push- und Pull Tests ein weiteres direktes Verfahren zur Charakterisierung des mikrobiellen *in situ* Abbaupotenzials in kontaminierten Aquiferen zur Verfügung. Mit *in situ* Mikrokosmen kann der mikrobielle Abbau von Grundwasserkontaminanten mit wenig Aufwand und ohne spezielle Vorkenntnisse zu den biogeochemischen Milieubedingungen direkt in einer Grundwassermessstelle untersucht werden. Die Methode eignet sich gleichermaßen für Anwendungen in konventionellen Grundwassermessstellen und für Multilevel-Untersuchungen, bei denen tiefenorientierte Aussagen bezüglich des *in situ* Abbaupotenzials gewonnen werden können (siehe Kapitel 2, 3, 5). Darüber hinaus ist der Einsatz von *in situ* Mikrokosmen nicht nur auf den Grundwasserleiter beschränkt. In Untersuchungen zum MCB Abbau in einer Pflanzenkläranlage wurden BACTRAPS erstmals direkt im Sedimentkörper inkubiert (Kapitel 7, Anhang H).

Die Verwendung ¹³C-markierter Verbindungen und die Analyse der Fettsäurefraktionen mittels GC-C-IRMS ermöglichen einen sehr sensitiven Nachweis der mikrobiellen Abbauaktivität (Boschker and Middelburg, 2002; Zengler et al., 1999), so dass selbst geringe Stoffumsätze im unteren Nanogramm Bereich empfindlich detektiert werden können (siehe Kapitel 4, 5, 6). Damit sich ausreichend Mikroorganismen auf dem Testsystem ansiedeln und eine substanziale ¹³C-Anreicherung der Biomasse erfolgen kann, sollte die Inkubationsdauer der *in situ* Mikrokosmen im Grundwasser circa zwei bis drei Monate betragen. Es kann also

bereits innerhalb weniger Monate eine Charakterisierung des Abbaupotenzials an einem kontaminierten Standort vorgenommen werden, weshalb sich *in situ* Mikrokosmen insbesondere zur Untersuchung der sehr langsam ablaufenden anaeroben Prozesse, wie für den Nachweis der Biodegradation persistenter Kontaminanten (z.B. MCB), eignen (Kapitel 5 bis 7).

Bisher galt MCB unter anoxischen Bedingungen als nicht abbaubare Verbindung (Field and Sierra-Alvarez, 2008). Der Einsatz stabiler Isotope ermöglichte den sensitiven Nachweis der vollständigen Mineralisierung in Labormikrokosmen sowie die Transformation in die bakterielle Biomasse unter *in situ* Bedingungen. Dadurch konnte der anaerobe MCB-Abbau sowohl im Labor als auch unter *in situ* Bedingungen nachgewiesen werden (Kapitel 5 bis 7). Die Bestimmung der Reaktionsmechanismen sowie die Identifizierung der am anaeroben MCB-Abbau beteiligten Mikroorganismen sind Gegenstand zukünftiger Untersuchungen. Im Rahmen weiterer Tracerversuche mit isotopisch markiertem MCB könnten charakteristische Metabolite Informationen zum Abbauweg liefern und RNA- bzw. DNA SIP zur Analyse der am MCB Abbau beteiligten mikrobiellen Gemeinschaft genutzt werden.

Ziel der Untersuchungen mit *in situ* Mikrokosmen ist es, die Abbaubarkeit organischer Grundwasserkontaminanten unter möglichst natürlichen Bedingungen nachzuweisen. Entgegen den alternativen Labormethoden werden Versuche mit BACTRAPS daher direkt unter den vorherrschenden Aquiferbedingungen durchgeführt, in dem das Testsystem in einer Grundwassermessstelle installiert wird. Von Nachteil ist hierbei, dass eine Messstelle nur bedingt die Aquiferbedingungen repräsentieren kann, da das Pegelwasser über längere Zeit stagniert und deshalb nur eingeschränkt dem umgebenden Grundwasser entspricht. Eine optimierte Art der Probenahme könnte erfolgen, in dem das Testsystem stattdessen direkt im Aquifersediment inkubiert wird. Diese Art der Beprobung ist jedoch technisch aufwendig und für Routineuntersuchungen ungeeignet, insbesondere für tiefer liegende Grundwasserstockwerke.

In situ Mikrokosmen bestehen aus einem Aufwuchskörper, der den Mikroorganismen als Substratpool und als besiedelbare Oberfläche dient. Um die Ergebnisse der BACTRAP-Untersuchungen auf den Grundwasserleiter übertragen zu können, sollten die Aufwuchskörper den Bewuchs sowie die Abbaaktivität in keiner Weise stimulieren. Wie die Ergebnisse in Kapitel 4 gezeigt haben, konnten die dominierenden Organismen der Grundwassermikroflora mittels molekularbiologischer Methoden (SSCP) auch auf den *in situ* Mikrokosmen nachgewiesen werden (Fig. 4-6). Darüber hinaus stimmten die Ergebnisse der molekularbiologischen Untersuchungen sehr gut mit den geochemischen Milieubedingungen überein. Untersuchungen im Randbereich der BTEX-Fahne am Standort Zeitz zeigten eine Dominanz von Alpha- und Betaproteobakterien und die Sequenzierung ausgewählter Banden von Toluol beladenen BACTRAPS ergab gute Übereinstimmung mit *Azoarcus* sp.,

Zusammenfassung

bekannt als Toluol- beziehungsweise *m*-Xylol Abbauer unter nitratreduzierenden Bedingungen (Tab. 4-2). Diese Ergebnisse korrelierten sehr gut mit der lokalen Geochemie am Fahnenrand, wo Nitrat als relevanter Elektronenakzeptor in Frage kommt.

Vergleichbare Untersuchungen mittels DNA-Fingerprint Methoden zur Phylogenie der auf den *in situ* Mikrokosmen siedelnden mikrobiellen Gemeinschaft an einem weiteren BTEX kontaminierten Standort ergaben eine Dominanz von bisher noch nicht kultivierten Mikroorganismen mit nahem Verwandtschaftsgrad zu Eisen- und Sulfatreduzierern, insbesondere der *Geobacter* und *Desulfurumonas* Gruppen. Diese Organismen sind dafür bekannt, am anaeroben Abbau organischer Kontaminanten im Grundwasser beteiligt zu sein. Da die dominierenden terminalen Elektronenakzeptoren im Aquifer Sulfat und Eisen(III) sind, konnte auch an diesem Standort eine gute Übereinstimmung zwischen der Geochemie und den molekularbiologischen Untersuchungsergebnissen gefunden werden. Die Ergebnisse verschiedener Feldversuche zur Charakterisierung der am Schadstoffabbau beteiligten Mikroorganismen haben gezeigt, dass *in situ* Mikrokosmen geeignet sind, um in Kombination mit DNA Fingerprinting Methoden empfindlich die mikrobielle Gemeinschaft in Übereinstimmung mit der regionalen Geochemie abzubilden. Auch die Beladung der Aufwuchskörper mit verschiedenen Substraten wie Benzol und Toluol führte im Vergleich zu unbeladenen Testsystemen zu keiner signifikanten Veränderung hinsichtlich der Zusammensetzung der mikrobiellen Gemeinschaft, wie die Fettsäureprofile und insbesondere die SSCP Analysen gezeigt haben (Kapitel 3, 4). Dies verdeutlicht, dass sich das Testsystem nicht selektiv auf die bakterielle Gemeinschaft auswirkt und im Wesentlichen kein präferentieller Abbau stattfindet; eine wichtige Voraussetzung für die Anwendung der *in situ* Mikrokosmen zur Erfassung des Abbaupotenzials kontaminiert Standorte.

Basierend auf dem Einsatz ¹³C-markierter Substrate kann selbst in komplexen Systemen, mit multiplen Kontaminationsspektrum, anhand der isotopischen Markierung in Biomarkern die *in situ* Biodegradation für einen bestimmten Schadstoff nachgewiesen werden. Eine fundamentale Grundlage für die Anwendung des Testsystems ist jedoch, dass die Schadstoffe, deren Abbau untersucht werden soll, von den Mikroorganismen als Kohlenstoffquelle benutzt werden, da ansonsten die Biodegradation nicht mit einer Inkorporation ¹³C-markierten Kohlenstoffs in die Biomasse verbunden ist. Daher ist der Nachweis der reduktiven Dehalogenierung (RDH) von chlorierten, bromierten oder fluorierten Verbindungen, die während der RDH als Elektronenakzeptoren genutzt werden, mit dieser Methode nicht möglich.

Die mikrobielle Abbauaktivität wird anhand der Inkorporation ¹³C-markierten Kohlenstoffs in Biomarkermoleküle nachgewiesen. Hat eine ¹³C-Anreicherung in den Zellbestandteilen stattgefunden, so gilt dies als eindeutiger Indikator für *in situ* Abbau. Ist keine ¹³C-Anreicherung in der Biomasse nachweisbar, kann ein *in situ* Abbaupotenzial dennoch nicht

ausgeschlossen werden. Die Ergebnisse zeigen in diesem Fall nur, dass mithilfe der *in situ* Mikrokosmen ein Nachweis des Abbaus nicht erfolgen konnte. Verschiedene Ursachen hierfür sind denkbar. Da insbesondere unter anoxischen Bedingungen das Wachstum von Schadstoff abbauenden Mikroorganismen sehr langsam erfolgt, war möglicherweise die Inkubationszeit zu kurz, so dass sich die mikrobielle Gemeinschaft des Aquifers nicht auf dem Testsystem etablieren konnte und daher nicht ausreichend Schadstoff abbauende Mikroorganismen zur Verfügung standen. Weiterhin ist denkbar, dass Mikroorganismen die ¹³C-markierten Substrate primär für den Energiestoffwechsel und nicht zur Synthese von Biomasse (Zellwachstum) verwenden (Thauer et al., 1977), so dass keine Inkorporation des markierten Kohlenstoffs in die Biomarker erfolgen kann. Findet eine Umsetzung der ¹³C-markierten Substrate ausschließlich durch den Energiestoffwechsel ohne Zellwachstum statt, so könnten dennoch isotopisch markierte Metabolite als Indikator für Biodegradation auf dem Testsystem nachgewiesen werden.

Eine Quantifizierung des mikrobiellen Abbaus mithilfe von *in situ* Mikrokosmen ist schwierig, aber bedingt möglich (Kapitel 4). Anhand der Menge des ¹³C-markierten Kohlenstoffs in den Fettsäuren kann die Menge des dafür benötigten Substrats bestimmt werden. Unter weiterer Berücksichtigung des Anteils der Fettsäuren an der Gesamtbiomasse und unter der Annahme repräsentativer Ertragskoeffizienten für den anaeroben Abbau kann schließlich die Gesamtmenge des ¹³C-markierten Substrats ermittelt werden, die während der Inkubation des Testsystems mikrobiell umgesetzt wurde. Auf diese Art und Weise berechnete Umsatzraten für Versuche mit Benzol und Toluol ergaben, dass maximal 2,5 % des Substratverlustes auf mikrobiellen Abbau zurückgeführt werden konnten (Kapitel 4), so dass Verluste infolge von Desorption um Größenordnungen höher waren (Geyer et al., 2005). Alternativ sind semiquantitative Aussagen anhand der ¹³C-Anreicherungen in den Fettsäuren möglich, da die Intensität der Markierung als direktes Maß der mikrobiellen Aktivität betrachtet werden kann. Abgesehen von geringen Isotopenfraktionierungsprozessen bei der Fettsäuresynthese spiegeln die Isotopensignaturen der Fettsäuren im Wesentlichen das Isotopenverhältnis des Wachstumssubstrats (Kohlenstoffquelle) wider (Abraham et al., 1998; Deniro and Epstein, 1978). Pelz et al. (2001a) zeigten in Markierungsexperimenten gleiche Isotopensignaturen von Fettsäuren verschiedener Kettenlängen und auch ein wesentlicher Einfluss der Wachstumsphasen auf die Isotopensignaturen konnte nicht beobachtet werden (Abraham et al., 1998). Der Kohlenstoff wird also gleichermaßen zum Aufbau von Membranlipiden verwendet, so dass unterschiedlich markierte Fettsäuren auf eine diverse Besiedlung durch mikrobielle Konsortien hindeuten können. Da das Testsystem im Grundwasser nicht ausschließlich von Schadstoff abbauenden Mikroorganismen besiedelt wird, muss zusätzlich mit einem Verdünnungseffekt des Isotopensignals gerechnet werden, da Fettsäuren von Bakterien, die nicht am Abbau beteiligt sind, folglich auch keine ¹³C-

Zusammenfassung

Anreicherung aufweisen. Als alternative Kohlenstoffquellen stehen den Mikroorganismen zum Beispiel die unmarkierten Aquiferkontaminanten oder gelöster organischer Kohlenstoff (DOC) zur Verfügung. Trotzdem kann anhand der Intensität der Markierung auf die Abbauaktivität geschlossen werden. Eine intensivere Inkorporation ¹³C-markierten Kohlenstoffs deutet auf eine höhere Abbauaktivität hin. Auf dieser Ebene können an einem Standort Ergebnisse verschiedener Messstellen hinsichtlich ihrer Abbauaktivitäten miteinander verglichen werden. Beim Vergleich von Benzol- und Toluol BACTRAPS dokumentierte die deutlich höhere Anreicherung ¹³C-markierten Kohlenstoffs in den Fettsäuren der Mikroorganismen, die die Toluol BACTRAPS besiedelten, einen bevorzugten und intensiveren Toluolabbau (Kapitel 3, (Geyer et al., 2005)). Diese Resultate stimmen gut mit früheren Studien überein, die bereits gezeigt haben, dass Toluol deutlich leichter abbaubar ist als Benzol (Cunningham et al., 2001; Da Silva and Alvarez, 2004; Wiedemeier et al., 1999). Ebenso wurden signifikante Unterschiede hinsichtlich des Abbaupotenzials zwischen hoch- und niedrig kontaminierten Bereichen der Schadstofffahne ermittelt, wobei für die weniger stark belasteten Zonen eine höhere ¹³C-Anreicherung bestimmt wurde (Kapitel 3). Möglicherweise wirken sich die hohen Schadstoffkonzentrationen im Kontaminationszentrum hemmend auf den bakteriellen Metabolismus aus. Darüber hinaus ist zu berücksichtigen, dass das Angebot unmarkierten Substrats im hoch kontaminierten Bereich wesentlich höher ist, als am Fahnenrand oder im Anstrom. Unmarkierter Schadstoff der den Mikroorganismen als Wachstumssubstrat zur Verfügung steht, kann zu einem Verdünnungseffekt der Markierung in den Fettsäuren führen. Prinzipiell zeigen die Ergebnisse jedoch, dass das Testsystem signifikante Unterschiede in Bezug auf die Intensität des Abbaus widerspiegeln kann, so dass *in situ* Mikrokosmen zur Charakterisierung des Abbaupotenzials geeignet sind.

Die Gesamtbiomasse eines *in situ* Mikrokosmos kann sowohl aus lebender als auch toter Biomasse sowie aus abbauaktiven und nicht aktiven Mikroorganismen bestehen. Da der Nachweis der *in situ* Aktivität anhand der ¹³C-Inkorporation in die Biomasse erfolgt, werden statt den PLFA bevorzugt die Gesamtlipide (TLFA) extrahiert. PLFA dienen als Biomarker für die lebende Biomasse (Tab. 1-2, (Green and Scow, 2000; Kaur et al., 2005)). TLFA hingegen berücksichtigen nicht nur Fettsäuren der Zellmembranen sondern auch der Speicherstoffe und eine Differenzierung zwischen lebender und toter Biomasse erfolgt nicht, so dass theoretisch die Fettsäuren aller Organismen, die am Schadstoffabbau beteiligt waren, extrahiert werden können. Da die Fettsäureprofile aus den BACTRAP-Untersuchungen (Kapitel 3 bis 7) ein geringes taxonomisches Potenzial aufwiesen, werden zur Charakterisierung der bakteriellen Gemeinschaft eher Nukleinsäure-basierende Fingerprint Verfahren empfohlen (Kapitel 3, 4).

Neben dem Nachweis der Biodegradation anhand isotopisch markierter Biomasse als direkter Indikator für mikrobielle Assimilation der Schadstoffe unter *in situ* Bedingungen, kann zusätzlich die Analyse charakteristischer Metabolite zum Nachweis des Schadstoffabbaus genutzt werden. Darüber hinaus bietet die Analyse von Metaboliten die Möglichkeit Abbauwege zu rekonstruieren. Im Rahmen eines Feldversuchs mit ¹³C-Toluol beladenen BACTRAPS unter anoxischen Aquiferbedingungen konnte ¹³C-markiertes Benzylsuccinat vom Aufwuchsmaterial der *in situ* Mikrokosmen extrahiert werden. Benzylsuccinat ist ein typischer Metabolit des anaeroben Toluolabbaus *via* Fumarataddition, katalysiert durch die Benzylsuccinatsynthase (Evans et al., 1992; Heider et al., 1999; Spormann and Widdel, 2000; Widdel and Rabus, 2001). Der Nachweis von Benzylsuccinat wurde bereits zuvor als Indikator für den anaeroben Toluolabbau genutzt (Beller, 2000; Beller, 2002; Elshahed et al., 2001; Griebler et al., 2004b; Reusser and Field, 2002). Des Weiteren wurde im Rahmen eines BACTRAP Experimentes zur Bestimmung der *in situ* Biodegradation von MCB ¹³C-markiertes Benzol (Fig. 7-3) auf den *in situ* Mikrokosmen nachgewiesen, ein Hinweis auf reduktive Dehalogenierung des MCB (Kapitel 7). Somit können zusätzlich Informationen zu den biogeochemischen Milieubedingungen während des mikrobiellen Abbaus sowie den Reaktionsmechanismen gewonnen werden, was eine bessere Charakterisierung der *in situ* Prozesse und die Rekonstruktion der Abbauwege ermöglicht. In diesem Zusammenhang ist die Reinheit der verwendeten ¹³C-markierten Substanzen ein wichtiger Aspekt, der beachtet werden muss, damit Rückschlüsse auf potenzielle Metabolite möglich sind. Bereits geringe Unreinheiten in den ¹³C-markierten Substraten könnten zu Fehlinterpretationen führen. So konnte beispielsweise festgestellt werden, dass ¹³C-markiertes MCB auch Spuren von ¹³C-markiertem Benzol enthält, was die Identifizierung von Benzol als potenziellen Metaboliten des MCB Abbaus *via* RDH erschweren kann (Kapitel 5, 7). Insofern auf den Mikrokosmen die nachweisbaren Anteile ¹³C-markierter Metabolite im Vergleich zu den Unreinheiten der Ausgangssubstanzen höher sind, können Aussagen zu den Reaktionsmechanismen getroffen werden.

Im Rahmen dieser Promotion konnte gezeigt werden, dass *in situ* Mikrokosmen für vielfältige Anwendungen geeignet sind. Es handelt sich um ein Testsystem, dass den Nachweis der Aktivität, Struktur und Funktion von am Abbau beteiligten Mikroorganismen ermöglicht. Darüber hinaus können *in situ* Mikrokosmen als Präinokulum für Anreicherungskulturen im Labor zur Verfügung stehen, um die Aktivität aus dem Feld ins Labor zu übertragen, mit dem Ziel langfristig die Schadstoff abbauenden Organismen zu kultivieren (Kapitel 5, 6).

8.2 Vergleich mit anderen Methoden

Kapitel 1.4 gibt einen Überblick über die Methoden, die zum Nachweis der *in situ* Biodegradation zur Verfügung stehen: Überwachung der räumlichen und zeitlichen Abnahme

Zusammenfassung

der Schadstoffkonzentrationen; Ausbildung charakteristischer Redoxzonen; Anreicherung spezifischer Metabolite; Labormikroskopienstudien; molekularbiologische Techniken wie Nukleinsäure basierende Fingerprint Verfahren; substratspezifische Isotopenanalyse (CSIA) sowie *in situ* Tracerversuche mit radioaktiven oder stabilen Isotopen markierten Substraten. Mit Ausnahme der zwei zuletzt genannten Verfahren ist mit dem Großteil dieser Methoden nur ein indirekter Nachweis des *in situ* Abbaupotenzials möglich, was zur Entwicklung der *in situ* Mikrokosmen motiviert hat. Deshalb soll dieses Kapitel im Wesentlichen dazu dienen, anhand der Ergebnisse dieser Promotion eine Evaluierung und Gegenüberstellung der *in situ* Mikrokosmen mit den anderen genannten Methoden vorzunehmen.

Es ist schwierig in einer Schadstofffahne mit komplexem Kontaminantenspektrum abnehmende Schadstoffkonzentrationen mit dem Verbrauch von Elektronenakzeptoren und der damit in Verbindung stehenden Ausbildung typischer Redoxzonen zu korrelieren, um den Abbau einzelner Schadstoffe nachzuweisen. Häufig stellt die Zehrung von Elektronenakzeptoren in kontaminierten Aquiferen ein komplexes Zusammenspiel unterschiedlicher Abbauprozesse dar. Ein gleichzeitiges Auftreten von signifikanten Intermediaten könnte die Charakterisierung der biogeochemischen Prozesse erleichtern, insofern ein ursprünglicher Eintrag der Metabolite am Untersuchungsstandort ausgeschlossen werden kann. Alternativ werden häufig mikrobiologische Untersuchungen an Standortmaterial (Grundwasser, Sediment) im Rahmen von Labormikroskopienstudien durchgeführt. Ein wesentlicher Vorteil der Laborstudien ist, dass in geschlossenen Laborsystemen Stoffbilanzen einfacher erfasst werden können. Der wesentliche Nachteil liegt in der eingeschränkten Übertragbarkeit der Ergebnisse aus dem Labor auf die weitaus komplexeren *in situ* Bedingungen kontaminiert Aquifere. Außerdem ist die Vielzahl der Schadstoffabbauer noch nicht kultivierbar und Laborversuche sowie Säulenexperimente unter quasi *in situ* Bedingungen sind sehr schwierig sowie arbeits- und zeitaufwändig, womit die Limitationen dieser Methoden deutlich werden.

Die Most-Probable-Number (MPN) und DNA-Fingerprint Methoden sind zwar für die Bestimmung von Gesamtzellzahlen bzw. zur Charakterisierung der mikrobiellen Gemeinschaften geeignet, aber allein der Nachweis über die Anwesenheit bestimmter Mikroorganismen erlaubt noch keine Rückschlüsse auf deren Abbauaktivität, so dass diese Methoden nur Aussagen zum Abbaupotenzial liefern können, ein direkter Nachweis des Abbaus jedoch nicht erfolgen kann. Ein direkter Nachweis der vollständigen Mineralisierung sowie der Transformation von Schadstoffen in die Biomasse ist allein mithilfe von stabilen bzw. radioaktiven Isotopen als Tracer möglich. Aufgrund des erhöhten Risikos, ausgehend von radioaktiven Stoffen, für den Mensch und die Umwelt werden bevorzugt Verbindungen mit stabilen Isotopen in offenen Systemen genutzt.

Eine etablierte Methode zur Charakterisierung der Abbauprozesse im Aquifer stellen *in situ* Tracerversuche dar, die entlang einer Grundwasserfließstrecke oder als *Pusch- und Pull* Tests in einer Grundwassermessstelle durchgeführt werden (Kapitel 1.4.2). Diese Methoden eignen sich, um den Schadstoffabbau anhand isotopisch markierter Biomasse, Metabolite oder Mineralisierungsprodukte nachzuweisen. Die Aussagen, die mit *in situ* Mikrokosmen oder *in situ* Tracerversuchen gewonnen werden können, sind vergleichbar. Während Tracerversuche entlang einer Fließstrecke mit umfangreichen Probenahmen zur Bestimmung der Tracer Durchbruchskurven verbunden sind, erfordern *in situ* Mikrokosmen neben dem Ein- und Ausbau keine weiteren Feldarbeiten, so dass der Arbeitsaufwand bei der Durchführung von Tracerversuchen deutlich größer ist. Darüber hinaus werden für *in situ* Tracerversuche größere Mengen isotopisch markierter Substanzen benötigt, als für die Analyse mit *in situ* Mikrokosmen. Das führt zu deutlich höheren Kosten. Ein Vorteil der Tracerversuche liegt darin, dass der Aquifer den tatsächlichen Reaktionsraum darstellt, während *in situ* Mikrokosmen innerhalb von Grundwassermessstellen inkubiert werden. Weiterhin erlauben die Tracerversuche auch eine quantitative Abschätzung der Biodegradation im Aquifer. Beide Methoden basieren auf dem Einsatz isotopisch markierter Substrate, was einen direkten Nachweis des mikrobiellen Abbaus bestimmter Kontaminanten anhand des Verbleibs des ¹³C-markierten Kohlenstoffs ermöglicht.

Eine alternative Möglichkeit zum Nachweis der *in situ* Biodegradation bietet die Analyse der substratspezifischen Isotopenfraktionierung (CSIA) (Kapitel 1.4.1). Tracerversuche (Fischer et al., 2006; Reusser and Field, 2002) und CSIA (Anhang I) sind die derzeit einzigen Verfahren, die neben *in situ* Mikrokosmen die direkte Erfassung des mikrobiellen Schadstoffabbaus im kontaminierten Aquifer erlauben.

Unter der Voraussetzung, dass der mikrobielle Abbau zu einer signifikanten Veränderung der Isotopensignaturen in der residualen Schadstofffraktion führt, kann die *in situ* Biodegradation mittels CSIA über einen Fließweg auch quantitativ erfasst werden. Dazu ist in der Regel ein signifikanter Abbau mit einem deutlichen Konzentrationsgradienten notwendig, der vorwiegend in den Randbereichen von Schadstofffahnen beobachtet wird. Sind die hydrogeochemischen Milieubedingungen des Aquifers bekannt, kann der biologische Abbau unter Annahme charakteristischer Isotopenfraktionierungsfaktoren auch quantifiziert werden. Zur Analyse des Abbaupotenzials im hoch kontaminierten Bereich der Schadstoffquelle reicht die Empfindlichkeit dieser Methode oft nicht aus, da das Abbausignal sehr stark durch die hohen Anteile von nicht abgebauten Schadstoffen überlagert wird, wie im Rahmen eines Multilevel-Experimentes zur Untersuchung der vertikalen Struktur einer Benzolfahne gezeigt werden konnte (Kapitel 2,3). In den zentralen Bereichen der Fahne mit maximalen Schadstoffkonzentrationen lieferten die Benzolisotopensignaturen keinen Hinweis auf Benzolabbau, während mithilfe von *in situ* Mikrokosmen anhand der Inkorporation ¹³C-

Zusammenfassung

markierten Kohlenstoffs auch im hoch kontaminierten Bereich der Schadstofffahne Biodegradation nachweisbar war.

Für Schadstoffe mit großen Molekülen (> 12 Kohlenstoffatome), z.B. höher molekulare PAKs, ist CSIA nicht geeignet, da der messbare Isotopeneffekt zu gering ist, um sensitiv genug nachgewiesen werden zu können (Meckenstock et al., 2004a). Eine weitere Schwierigkeit stellen heterogene Schadstoffquellen und überlagerte Schadstofffahnen dar, weil es auch zu einer Überlagerung der Isotopensignale kommen kann. Abbaureaktionen, die prinzipiell nicht mit einer signifikanten Isotopenfraktionierung verbunden sind, wie teilweise für den aeroben MCB- und Toluolabbau beobachtet wurde (Kaschl et al., 2005; Morasch et al., 2002; Sherwood Lollar et al., 1999), können folglich nicht mittels CSIA erfasst werden. In diesem Fall können *in situ* Mikrokosmen einen wesentlichen Beitrag zur Charakterisierung der *in situ* Biodegradation leisten. Wie weiterhin in Kapitel 6 für chlorierte Benzole gezeigt wurde, stellen sequenzielle Abbauwege wie die RDH ein zusätzliches Problem für die Anwendung des CSIA Konzeptes dar, weil Intermediate sowohl durch Bildung als auch mikrobiellen Abbau, verbunden mit einem gegensätzlichen Isotopeneffekt, beeinflusst werden. In diesem Fall können durch die Berechnung einer kumulativen Isotopenbilanz Informationen zum Abbaupotenzial erhalten werden. Dennoch sind ergänzende Untersuchungen zur Verifizierung der Ergebnisse hilfreich, insbesondere da für die Beurteilung des *in situ* Abbaupotenzials kontaminiert Standorte mehrere Beweislinien gefordert werden.

Wie in den verschiedenen Feldstudien zur Evaluierung und Verifizierung des Testsystems gezeigt werden konnte, liefern *in situ* Mikrokosmen Ergebnisse, deren Plausibilität anhand der Übereinstimmung mit Ergebnissen anderer Methoden überprüft wurde. Die Analyse der vertikalen Struktur einer Benzolfahne mit geochemischen und isotopenchemischen Methoden ergab ein deutliches Abbaupotenzial für Benzol, was durch den Einsatz von *in situ* Mikrokosmen verifiziert werden konnte (Kapitel 2,3). Ist der Einsatz alternativer Methoden wie CSIA oder *in situ* Tracerversuchen unter bestimmten Umständen limitiert, so können *in situ* Mikrokosmen weiterführende Informationen zur Charakterisierung der *in situ* Biodegradation liefern, insbesondere für den hoch kontaminierten Bereich von Schadstofffahnen oder an komplexen Standorten. Abgesehen von Tracerversuchen existieren keine Methoden, mit denen der biologische Schadstoffabbau an kontaminierten Standorten mit vergleichbarer Sensitivität nachgewiesen werden kann.

8.3 Implementierung in die Altlastenpraxis

Verschiedene Prototypen des Testsystems wurden bisher an unterschiedlichen Untersuchungsstandorten im Rahmen von Industriekooperationen erfolgreich erprobt und es

liegen Erfahrungen zum Einsatz der BACTRAPS für verschiedene Kontaminanten wie Benzol, Toluol, MCB, MTBE und PAK vor (Tab. 8-1).

Die Zusammenarbeit mit der Industrie und Umweltbehörde ermöglichte es, dieses Forschungsvorhaben bereits von Beginn an in einen angewandten Kontext zu stellen und so die Belange von Wissenschaft und Praxis miteinander zu verknüpfen. Untersuchungen mit *in situ* Mikrokosmen, die in Kooperation mit einem Industriepartner an einem ehemaligen Kokereistandort durchgeführt wurden, haben dazu beigetragen, den Standort in ein *Natural Attenuation* Vorhaben zu überführen. In diesem Vorhaben wird auf aktive Sanierungsmaßnahmen über einen bestimmten Zeitraum verzichtet, um die Stabilität der Schadstofffahne und die *in situ* Abbauprozesse zu untersuchen sowie ein Grundwassermanagement- und Überwachungskonzept zu entwickeln. *Natural Attenuation* wurde für einen Zeitraum von 5 Jahren genehmigt, mit der Auflage weitere Monitoring-Untersuchungen durchzuführen, um den Standort zu erkunden. Im Rahmen dieser umfangreichen Standorterkundungen wurden BACTRAP Untersuchungen zum geforderten Nachweis der *in situ* Biodegradation eingesetzt. Auch die Untersuchungen zum MCB Abbau (Kapitel 5) erfolgten im Einverständnis mit der verantwortlichen Umweltbehörde. Die Methode der *in situ* Mikrokosmen ist bereits Bestandteil verschiedener Leitfäden und Handlungsempfehlungen (KORA, 2007; KORA, 2008; LABO, 2005), was die Bedeutung dieses Nachweisverfahrens im Bereich der Altlastenbewertung verdeutlicht. Anhand dieser Beispiele kann gezeigt werden, dass es seitens der Behörden bereits eine Akzeptanz für diese neuartige Methode zum geforderten Nachweis der *in situ* Biodegradation an kontaminierten Standorten gibt.

Entsprechend der verschiedenen Handlungsempfehlungen und Arbeitshilfen zur Anwendung von *Monitored Natural Attenuation* (MNA) als Standortsicherungsmaßnahme kontaminierte Aquifere werden mehrere Beweislinien über den Nachweis der *in situ* Biodegradation gefordert (Hessisches Landesamt für Umwelt und Geologie, 2004; ITVA, 2004; KORA, 2007; US-EPA, 1999). *In situ* Mikrokosmen können in diesem Zusammenhang einen wertvollen Beitrag zur Charakterisierung der natürlichen Abbauprozesse liefern, um *Monitored Natural Attenuation* als Standortsicherungsmaßnahme zu implementieren. Im Gegensatz zu einer Vielzahl herkömmlicher Verfahren (siehe Kapitel 1.4) handelt es sich bei dem Testsystem um ein innovatives Nachweisverfahren, welches innerhalb weniger Monate einen direkten Nachweis der *in situ* Biodegradation ermöglicht. Die Methode kann problemlos in das Monitoring-Programm implementiert werden, ohne Erfordernis zusätzlicher Maßnahmen.

Zusammenfassung

Tab. 8-1: Anwendungsbeispiele von *in situ* Mikrokosmen (BACTRAP®)

Kontaminante	Standort	Forschungsinstitut/ Industriepartner	Referenz
Benzol	Zeitz	UFZ	(Büning et al., 2005); (Geyer et al., 2005); (Kästner et al., 2006); (Stelzer et al., 2006a); (Stelzer et al., 2006b) Kapitel 2, 3, 4
Benzol	Gneisenau	DMT	(Büning et al., 2005); (Stelzer et al., 2006a) Dokumentation in internen Gutachten, (2007)
Benzol	Hansemann	DMT	Dokumentation in internen Gutachten, (2005, 2008)
Benzol	C.Rauxel	DMT	Dokumentation in internem Gutachten, (2008)
Benzol	Duisburg	HPC	in Bearbeitung, (2008)
Benzol	Gölzau	UFZ	in Bearbeitung, (2008)
Toluol	Zeitz	UFZ	(Geyer et al., 2005); (Kästner et al., 2006); (Stelzer et al., 2006a) Kapitel 3, 4
Toluol	Gneisenau	DMT	(Büning et al., 2005); (Stelzer et al., 2006a)
Toluol	C. Rauxel	DMT	Dokumentation in internem Gutachten, (2008)
Naphthalin	Hansemann	DMT	Dokumentation in internem Gutachten, (2007)
Naphthalin	Gneisenau	DMT	Dokumentation in internem Gutachten, (2007)
Naphthalin	C. Rauxel	DMT	Dokumentation in internem Gutachten, (2008)
Naphthalin	Hanau	HPC	Dokumentation in internem Gutachten, (2005)
Naphthalin	Karlsruhe	Roth und Partner	Dokumentation in internem Gutachten, (2007)
Naphthalin	Duisburg	HPC	in Bearbeitung, (2008)
Acenaphthen	Gneisenau	DMT	Dokumentation in internem Gutachten, (2007)
Acenaphthen	Karlsruhe	Roth und Partner	Dokumentation in internem Gutachten, (2007)
Anthracen	Hanau	HPC	Dokumentation in internem Gutachten, (2005)
Phenanthren	Hanau	HPC	Dokumentation in internem Gutachten, (2005)
MTBE	Leuna	UFZ	Dokumentation in internem Gutachten, (2004)
MTBE	vertraulich	GICON	Dokumentation in internem Gutachten, (2005)
MCB	Bitterfeld	UFZ	(Nijenhuis et al., 2007) Kapitel 5 (Braeckeveld et al., 2007a; Braeckeveld et al., 2007b) Kapitel 7, Anhang H
MCB	Hamburg	Roth und Partner	(Stelzer et al., 2008) Kapitel 6
Hexadekan	Gölzau	UFZ	in Bearbeitung, (2008)

8.4 Ausblick

Im Rahmen dieser Promotion wurde gezeigt, dass mithilfe von *in situ* Mikrokosmen die Aktivität, Struktur und Funktion von Schadstoff abbauenden Mikroorganismen analysiert werden kann. Damit steht ein weiteres innovatives Verfahren zum Nachweis der *in situ* Biodegradation kontaminierten Grundwasserleiter in der Altlastenpraxis zur Verfügung. Der Großteil der bisherigen Untersuchungen und Feldanwendungen wurde für Benzol, Toluol und MCB durchgeführt (Kapitel 2 bis 7, Tab. 8-1). Zukünftig soll das Testsystem für Anwendungen an PAK oder MTBE kontaminierten Standorten optimiert werden. Diese Verbindungen zählen zu den häufigsten Grundwasserkontaminanten und insbesondere für höher molekulare PAKs gibt es derzeit keine alternativen Methoden zum direkten Nachweis der *in situ* Biodegradation. Die verschiedenen Schadstoffe unterscheiden sich deutlich hinsichtlich ihrer Eigenschaften (z.B. Sorption, Wasserlöslichkeit, Bioverfügbarkeit). Schwierigkeiten zeigten sich hierbei vor allem infolge der stark sorptiven Eigenschaften der PAKs auf den aktivkohlehaltigen Trägermaterialien, was die Extraktion und Analyse der Biomasse von den *in situ* Mikrokosmen erschwert, so dass alternative Trägermaterialien benötigt werden. Erste Untersuchungen zum Einsatz der *in situ* Mikrokosmen zum Nachweis des PAK Abbaus wurden durchgeführt, wobei Teflonfilter als Aufwuchskörper dienten. Anhand der ¹³C-Anreicherung in den Fettsäuren und Aminosäuren konnte der *in situ* Abbau von Naphthalin unter anoxischen Aquiferbedingungen nachgewiesen werden.

Ein weiterer Schwerpunkt zukünftiger Untersuchungen mit *in situ* Mikrokosmen ist die Identifikation der am Schadstoffabbau beteiligten Mikroorganismen mithilfe der Methode des DNA- oder RNA-SIP. In bisherigen Feldversuchen war die ¹³C-Anreicherung in den Biomarkern, der auf den *in situ* Mikrokosmen siedelnden Mikroorganismen, nicht ausreichend. Für eine optimale Auftrennung von schwerer und leichter Nukleinsäurefraktion mithilfe der Dichtegradientenzentrifugation wird nach bisherigem Stand des Wissens eine deutlich höhere Inkorporation ¹³C-markierten Kohlenstoffs von rund 30 Atomprozent benötigt (Radajewski et al., 2003).

9 Literatur

- Abraham, W.R., Hesse, C. and Pelz, O., 1998. Ratios of Carbon Isotopes in Microbial Lipids as an Indicator of Substrate Usage. *Applied Environmental Microbiology*, 64(11): 4202-4209.
- Abrajano, T.A., Murphy, D.E., Fang, J., Comet, P. and Brooks, J.M., 1994. $^{13}\text{C}/^{12}\text{C}$ ratios in individual fatty acids of marine mytilids with and without bacterial symbionts. *Organic Geochemistry*, 21(6-7): 611-617.
- Adrian, L. and Gorisch, H., 2002. Microbial transformation of chlorinated benzenes under anaerobic conditions. *Research in Microbiology*, 153: 131-137.
- Adrian, L., Szewzyk, U. and Gorisch, H., 2000a. Bacterial growth based on reductive dechlorination of trichlorobenzenes. *Biodegradation*, 11(1): 73-81.
- Adrian, L., Szewzyk, U., Wecke, J. and Görisch, H., 2000b. Bacterial dehalorespiration with chlorinated benzenes. *Nature*, 408: 580-583.
- Alfreider, A., Vogt, C., Hoffmann, D. and Babel, W., 2003. Diversity of ribulose-1, 5-bisphosphate carboxylase/oxygenase large-subunit genes from groundwater and aquifer microorganisms. *Microbial Ecology*, 45: 317-328.
- Amann, R.I., Ludwig, W. and Schleifer, K.H., 1995. Phylogenetic identification and in situ detection of individual microbial cells without cultivation. *Microbiological Reviews*, 59: 143-169.
- Anderson, R.T. and Lovley, D.R., 2000. Anaerobic biodegradation of benzene under sulfate-reducing conditions in a petroleum-contaminated aquifer. *Environmental Science and Technology*, 34: 2261-2266.
- Anderson, R.T., Rooney-Varga, J.N., Gaw, C.V. and Lovley, D.R., 1998. Anaerobic benzene oxidation in the Fe(III) reduction zone of petroleum-contaminated aquifers. *Environmental Science and Technology*, 32: 1222-1229.
- Anderson, T.A. and Walton, B.T., 1995. Comparative fate of [^{14}C] trichloroethylene in the root zone of plants from a former solvent disposal site. *Toxicology and Chemistry*, 14: 2041-2047.
- Andrade, E.M.A., Palacio, H.A.Q., Souza, I.H., Leao, R.A.O. and Guerreiro, M.J., 2008. Land use effects in groundwater composition of an alluvial aquifer (Trussu River, Brazil) by multivariate techniques. *Environmental Research*, 106(2): 170-177.
- Anonymous, 1995. IAEA-TECHDOC-825, International Atomic Energy Agency.
- Arneth, J.D., Milde, G., Kerndorff, H. and Schleyer, R., 1989. Waste deposit influences on groundwater quality as tool for waste type and site selection for final storage quality. In: P. Baccini (Editor), *The Landfill*, Lecture Notes in Earth Science. Springer-Verlag, Berlin, Germany.
- Baker, L., 1998. Design considerations and applications for wetland treatment of high-nitrate waters. *Water Science and Technology*, 31: 389-395.
- Barragan, M.J.L. et al., 2004. The *bzd* gene cluster, coding for anaerobic benzoate catabolism in *Azoarcus* sp. strain CIB. 186: 5762-5774.
- Bayrisches Landesamt für Wasserwirtschaft, 2004. Natürliche Schadstoffminderung bei Grundwasserverunreinigungen durch Altlasten und schädliche Bodenveränderungen - Natural Attenuation - Merkblatt Nr. 3.8/3 vom 05.11.2004, München.
- Bekins, B.A. et al., 2001. Progression of natural attenuation processes at a crude oil spill site: II. Controls on spatial distribution of microbial population. *Journal of Contaminant Hydrology*, 53: 387-406.

- Beller, H.R., 2000. Metabolic indicators for detecting in situ anaerobic alkylbenzene degradation. *Biodegradation*, 11(2-3): 125-139.
- Beller, H.R., 2002. Analysis of benzylsuccinates in groundwater by liquid chromatography/tandem mass spectrometry and its use for monitoring in situ BTEX biodegradation. *Environmental Science and Technology*, 36: 2742-2728.
- Bianchin, M., Smith, L., Barker, J.F. and Beckie, R., 2006. Anaerobic degradation of naphthalene in a fluvial aquifer: A radiotracer study. *Journal of Contaminant Hydrology*, 84: 178-196.
- Bisswanger, H., 2000. Enzymkinetik. Wiley-VCH, Weinheim, 268 pp.
- Blessing, M., Jochmann, M.A. and Schmidt, T.C., 2008. Pitfalls in compound-specific isotope analysis of environmental samples. *Analytical and Bioanalytical Chemistry*, 390: 591-603.
- Bligh, E.G. and Dyer, W.J., 1959. A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37(8): 911-917.
- Blumberg, M., Seifert, R., Nauhaus, K., Pape, T. and Michaelis, W., 2005. In vitro study of lipid biosynthesis in an anaerobically methaneoxidising microbial mat. *Applied and Environmental Microbiology*, 71: 4345-4351.
- BMBF, 2000. Richtlinien über die Förderung von Forschungs- und Entwicklungsvorhaben der Altlastensanierung zum Thema "Kontrollierter natürlicher Rückhalt und Abbau von Schadstoffen bei der Sanierung kontaminierten Boden und Grundwässer" im Programm der Bundesregierung "Forschung für die Umwelt". 1.2.2000; Bonn.
- Borjesson, G., Sundh, I., Tunlid, A. and Svensson, B.H., 1998. Methane oxidation in landfill cover soils as revealed by potential oxidation measurements and PLFA analysis. *Soil Biology and Biochemistry*, 30: 1423-1433.
- Boschker, H.T.S. and Middelburg, J.J., 2002. Stable isotopes and biomarkers in microbial ecology. *FEMS Microbiology Ecology*, 40(2): 85-95.
- Boschker, H.T.S. et al., 1998. Direct linking of microbial populations to specific biogeochemical processes by ¹³C-labelling of biomarkers. *Nature*, 392: 801-805.
- Bowman, J.P., Sly, L.I., Nichols, P.D. and Hayward, A.C., 1993. Revised taxonomy of the methanotrophs: description of *Methylobacter* gen. nov., emendation of *Methylococcus*, validation of *Methylosinus* and *Methylocystis* species, and a proposal that the family Methylococcaceae includes only the group I methanotrophs. *International journal of systematic bacteriology*, 43: 735-753.
- Boyle, A.W., Hagglom, M.M. and Young, L.Y., 1999. Dehalogenation of lindane ([gamma]-hexachlorocyclohexane) by anaerobic bacteria from marine sediments and by sulfate-reducing bacteria. *FEMS Microbiol Ecol*, 29(4): 379.
- Bradley, P.M., Landmeyer, J. and Chapelle, F.H., 2002. TBA biodegradation in surface-water sediments under aerobic and anaerobic conditions. *Environmental Science and Technology*, 36: 4087-4090.
- Braeckeveldt, M. et al., 2007a. Assessment of in situ biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland. *Environmental Pollution*, 148(2): 428-437.
- Braeckeveldt, M. et al., 2007b. Biodegradation of chlorobenzene in a constructed wetland treating contaminated groundwater. *Water Science and Technology*, 56(3): 57-62.
- Brenna, J.T., Corso, T.N., Tobias, H.J. and Caimi, R.J., 1997. High-precision continuous-flow isotope ratio mass spectrometry. *Mass Spectrometry Reviews*, 16(5): 227-258.

Literatur

- Breukelen, B.M. and Griffioen, J., 2004. Biogeochemical processes at the fringe of a landfill leachate pollution plume: potential for dissolved organic carbon, Fe(II), Mn(II), NH₄, and CH₄ oxidation. *Journal of Contaminant Hydrology*, 73(1-4): 181-205.
- Brinch-Iversen, J. and King, G.M., 1990. Effects of substrate concentration, growth rate, and oxygen availability on relationships among bacterial carbon, nitrogen and phospholipid phosphorus content. *FEMS Microbiology Ecology*, 74: 345-356.
- Bull, I.D., Parekh, N.R., Hall, G.H., Ineson, P. and Evershed, R.P., 2000. Detection and classification of atmospheric methane oxidising bacteria in soil. *Nature*, 405: 175-178.
- Büning, C. et al., 2005. *In situ* Mikrokosmenuntersuchungen mit ¹³C-markiertem Benzol und Toluol zum Nachweis des natürlichen biologischen Abbaus im Grundwasser und zur molekularbiologischen Analyse der abbauaktiven Mikroflora. *Altlastenspektrum*, 14(3): 137-146.
- Butler, J.L., Williams, M.A., Bottomley, P.J. and Myrold, D.D., 2003. Microbial community dynamics associated with rhizosphere carbon flow. *Applied and Environmental Microbiology*, 69: 6793-6800.
- Cerniglia, C.E., 1984. Microbial transformation of aromatic hydrocarbons. In: R.M. Atlas (Editor), *Petroleum microbiology*. Macmillan Publishing Co., New York, pp. 99-128.
- Chakraborty, R. and Coates, J.D., 2004. Anaerobic degradation of monoaromatic hydrocarbons. *Applied Microbiology and Biotechnology*, 64(4): 437-446.
- Chakraborty, R. and Coates, J.D., 2005. Hydroxylation and carboxylation: crucial steps of anaerobic benzene degradation by *Dechloromonas* strain RCB. *Applied and Environmental Microbiology*, 71: 5427-5432.
- Chang, Y.J. et al., 2005. Microbial Incorporation of ¹³C-Labeled Acetate at the Field Scale: Detection of Microbes Responsible for Reduction of U(VI). *Environmental Science and Technology*, 39(23): 9039-9048.
- Chapelle, F.H., Bradley, P.M., Lovley, D.R. and Vroblesky, D.A., 1996. Measuring rates of biodegradation in a contaminated aquifer using field and laboratory methods. *Ground Water*, 34(4): 691-698.
- Chartrand, M.M.G., Morrill, P.L., Lacrampe-Couloume, G. and Lollar, B.S., 2005. Stable isotope evidence for biodegradation of chlorinated ethenes at a fractured bedrock site. *Environmental Science and Technology*, 39: 4848-4856.
- Cichocka, D., Imfeld, G., Richnow, H.H. and Nijenhuis, I., 2008. Variability in microbial carbon isotope fractionation of tetra- and trichloroethene upon reductive dechlorination. *Chemosphere*, 71(4): 639-648.
- Clark, I. and Fritz, P., 1997. *Environmental Isotopes in Hydrogeology*. Lewis-Publisher, New York, 328 pp.
- Clement, T.P., Truex, M.J. and Lee, P., 2002. A case study for demonstrating the application of US EPA's monitored natural attenuation screening protocol at a hazardous waste site. *Journal of Contaminant Hydrology*, 59: 133-162.
- Cline, J.D., 1969. Spectrophotometric determination of hydrogen sulfide in natural waters. *Limnol Oceanogr*, 14: 454-458.
- Coates, J.D. et al., 2001. Anaerobic benzene oxidation coupled to nitrate reduction in pure culture by two strains of *Dechloromonas*. *Nature*, 411(6841): 1039-1043.
- Coplen, T.B. et al., 2006. After two decades a second anchor for the VPDB δ¹³C scale. *Rapid Communication Mass Spectrometry*, 20: 3165-3166.

- Cozzarelli, I.M., Eganhouse, R.P. and Baedecker, M.J., 1990. Transformation of Monoaromatic Hydrocarbons Organic Acid in Anoxic Groundwater Environment. *Environmental Geology*, 16(2): 135-141.
- Craig, H., 1957. Isotopic standards for carbon and oxygen and correction for mass-spectrometric analysis of carbon dioxide. *Geochimica et Cosmochimica Acta*, 12: 133-149.
- Crossmann, Z.M., Ineson, P. and Evershed, R.P., 2005. The use of ^{13}C labelling of bacterial lipids in the characterisation of ambient methane-oxidising bacteria in soils. *Organic Geochemistry*, 36: 769-778.
- Cunningham, J.A., Rahme, H., Hopkins, G.D., Lebron, C. and Reinhard, M., 2001. Enhanced in situ bioremediation of BTEX-contaminated groundwater by combined injection of nitrate and sulphate. *Environmental Science and Technology*, 35: 1663-1670.
- Da Silva, M.L.B. and Alvarez, P.J.J., 2004. Enhanced anaerobic biodegradation of benzene-toluene-ethylbenzene-xylene-ethanol mixtures in bioaugmented aquifer columns. *Applied and Environmental Microbiology*, 70: 4720-4726.
- Dahmke, A., Dethlefsen, F. and Wachter, T., 2004. Geochemische Untersuchungen des Aquifers. Teil des Projektendberichts RETZINA: REferenzTestfeld Zeitz zur Implementierung des "Natural Attenuation" Ansatzes, Institut für Geowissenschaften, Christian-Albrechts Universität Kiel.
- Dempster, H.S., Lollar, B.S. and Feenstra, S., 1997. Tracing organic contaminants in groundwater: A new methodology using compound-specific isotopic analysis. *Environmental Science and Technology*, 31(11): 3193-3197.
- Deniro, M.J. and Epstein, S., 1978. Influence of diet on distribution of carbon isotopes in animals. *Geochimica et Cosmochimica Acta*, 42: 495-506.
- Dethlefsen, F., Bliss, F., Wachter, T. and Dahmke, A., 2004. Reaktive Tracer zur Bestimmung der sedimentären Aquifer-Oxidationskapazität im Labor- und Feldversuch. *Grundwasser*, 9(1): 12-20.
- DIN, 1985. 38402-Teil 13 Probenahme aus Grundwasserleitern, Deutsches Institut für Normung e. V.
- DIN, 1996. EN ISO 5667-3 (A 21) Anleitung zur Konservierung und Handhabung von Proben, Deutsches Institut für Normung e. V.
- Dohrmann, A.B. and Tebbe, C.C., 2004. Microbial community analysis by PCR-single-strand conformation polymorphism (PCR-SSCP). In: G.A. Kowalchuk, F.J. de Bruijn, I.M. Head, A.D. Akkermans and J.D. van Elsas (Editors), *Molecular Microbial Ecology Manual*. Kluwer Academic Publisher, pp. 809-838.
- Dolfing, J. and Harrison, K., 1993. Redox and reduction potentials as parameters to predict the degradation pathway of chlorinated benzenes in anaerobic environments. *FEMS Microbiology Ecology*, 13: 23-30.
- Doll, A., 1998. Entscheidungsgrundlagen für Sicherungs- und Sanierungskonzepte für militärische Tanklager, Anlage I: Recherche zum Stand der Technik bei der Sicherung und Sanierung ehemals genutzter Tanklager insbesondere in den USA. Umweltforschungsplan des Bundesminister für Umwelt, Naturschutz und Reaktorsicherheit, FKZ 203 40 831, im Auftrag des Umweltbundesamtes; Berlin.
- Dowling, N.J.E., Widdel, F. and White, D.C., 1986. Phospholipid Ester-linked Fatty Acid Biomarkers of Acetate-oxidizing Sulphate-reducers and Other Sulphide-forming Bacteria. *Journal of General Microbiology*, 132: 1815-1825.
- Dumont, M.G. and Murell, J.C., 2005. Stable Isotope Probing - linking microbial identity to function. *Nature Reviews Microbiology*, 3: 499-504.

Literatur

- Dunbabin, J. and Bowmer, K., 1992. Potential use of constructed wetlands for treatment of industrial wastewaters containing metals. *Science of the Total Environmental Forensics*, 111: 151-168.
- Edlung, A., Nichols, P.D., Roffey, R. and White, D.C., 1985. Extractable and lipopolysaccharide fatty acid and hydroxy acid profiles from *Desulfovibrio* species. *Journal of Lipid Research*, 26: 982-988.
- Edwards, E.A. and Grbic-Galic, D., 1992. Complete mineralization of benzene by aquifer microorganisms under strictly anaerobic conditions. *Applied and Environmental Microbiology*, 58: 2663-2666.
- Ehrlich, H.L., 1998. Geomicrobiology: its significance for geology. *Earth-Science Review*, 45: 45-60.
- Elshahed, M.S., Gieg, L.M., McInerney, M.J. and Sufita, J.M., 2001. Signature metabolites attesting to the in situ attenuation of alkylbenzenes in anaerobic environments. *Environmental Science and Technology*, 35: 682-689.
- Elsner, M., McKelvie, J.R., Lacrampe-Couloume, G. and Lollar, B.S., 2007. Insight into Methyl *tert*-Butyl Ether (MTBE) Stable Isotope Fractionation from Abiotic Reference Experiments. *Environmental Science and Technology*, 41(16): 5693-5700.
- Elsner, M., Zwank, L., Hunkeler, D. and Schwarzenbach, R.P., 2005. A new concept linking observable stable isotope fractionation to transformation pathways of organic pollutants. *Environmental Science and Technology*, 39(18): 6896-6916.
- Evans, P.J., Ling, W., Goldschmidt, B., Ritter, E.R. and Young, L.Y., 1992. Metabolites formed during anaerobic transformation of toluene and o-xylene and their proposed relationship to the initial steps of toluene mineralization. *Applied and Environmental Microbiology*, 58: 496-501.
- Evershed, R.P. et al., 2006. ¹³C-Labelling of lipids to investigate microbial communities in the environment. *Current Opinion in Biotechnology*, 17(1): 72-82.
- Farhadiana, M., Vachelarda, C., Ducheza, D. and Larrochea, C., 2008. In situ bioremediation of monoaromatic pollutants in groundwater: A review. *Bioresource Technology*, in press, doi:10.1016/j.biortech.2007.10.025.
- Federle, T.W., 1986. Microbial distribution in soil. In: F. Megusar and M. Ganthaer (Editors), *Perspectives in Microbial Ecology*. Solveme Society for Microbiology, Ljubljana, pp. 493-498.
- Fennell, D.E., Nijenhuis, I., Wilson, S.F., Zinder, S.H. and Haggblom, M.M., 2004. *Dehalococcoides ethenogenes* strain 195 Reductively Dechlorinates Diverse Chlorinated Aromatic Pollutants. *Environ Sci Technol*, 38(7): 2075-2081.
- Fent, G., Hein, W.J., Moendel, M.J. and Kubiak, R., 2003. Fate of ¹⁴C-bisphenol A in soils. *Chemosphere*, 51(8): 735-746.
- Field, F.A. and Sierra-Alvarez, R., 2008. Microbial degradation of chlorinated benzenes. *Biodegradation*, in press, DOI 10.1007/s10532-007-9155-1.
- Fischer, A. et al., 2005. Konzept zur Quantifizierung des anaeroben in situ Schadstoffabbaus in BTEX-kontaminierten Grundwasserleitern mittels Deuterium-markierter Substanzen. *Altlastenspektrum*, 14(1): 1-12.
- Fischer, A. et al., 2006. A Multitracer Test Proving the Reliability of Rayleigh Equation-Based Approach for Assessing Biodegradation in a BTEX Contaminated Aquifer. *Environmental Science and Technology*, 40(13): 4245-4252.
- Fischer, A. et al., 2008. Combined carbon and hydrogen isotope fractionation investigations for elucidating benzene biodegradation pathways. *Environmental Science and Technology*, in press, ES702468f.

- Fischer, A. et al., 2007. Applicability of Stable Isotope Fractionation Analysis for the Characterization of benzene Biodegradation in a BTEX-contaminated Aquifer. Environmental Science and Technology, 41(10): 3689-3696.
- Fischer, A. et al., 2004. Charakterisierung des mikrobiellen Schadstoffabbaus mit Hilfe von isotopenchemischen Methoden. Grundwasser, 9(3): 159-172.
- Friedrich, M.W., 2006a. Stabile Isotopenbeprobung von Nukleinsäuren – ein Schlüssel zur Ökophysiologie nicht kultivierter Mikroorganismen. Biospektrum, 01/06: 39-40.
- Friedrich, M.W., 2006b. Stable-isotope probing of DNA: insights into the function of uncultivated microorganisms from isotopically labeled metagenomes. Current Opinion in Biotechnology, 17(1): 59-66.
- Fritz, H., Reineke, W. and Schmidt, E., 1992. Toxicity of chlorobenzene on *Pseudomonas* sp. strain RHO1, a chlorobenzene-degrading strain. Biodegradation, 2: 165-170.
- Frostegard, A. and Bååth, E., 1996. The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil. Biology and Fertility of Soils, 22: 59-65.
- Galimov, E., 1985. The Biological Fractionation of Isotopes. Academic Press Inc., London, 261 pp.
- Geyer, R. et al., 2005. In situ assessment of biodegradation potential using biotraps amended with ¹³C-labeled benzene or toluene. Environmental Science and Technology, 39(13): 4983-4989.
- Gibson, D.T. and Subramanian, V., 1984. Microbial degradation of aromatic hydrocarbons. In: D.T. Gibson (Editor), Microbial degradation of organic compounds. Dekker, New York, N.Y., pp. 181-252.
- Gilbert, E.S. and Crowley, D.E., 1997. Plant compounds that induce polychlorinated biphenyl biodegradation by *Arthrobacter* sp. strain B1B. Applied Environmental Microbiology, 63: 1933-1938.
- Gödeke, S. et al., 2004a. Strömungs- und Tracer-Transportmodellierung am Natural Attenuation-Standort Zeitz. Grundwasser, 9(1): 3-11.
- Gödeke, S. et al., 2004b. Tracertest zur Analyse des reaktiven Transports im Aquifer am Industriestandort Zeitz. Altlastenspektrum, 13(4): 181-189.
- Goodmann, K.J. and Brenna, J.T., 1992. High sensitivity tracer detection using high-precision gas chromatography-combustion isotope ratio mass spectrometry and highly enriched [U-¹³C]-labeled precursors. Analytical Chemistry, 64(10): 1088-1096.
- Grandel, S. and Dahmke, A., 2004. Monitored natural attenuation of chlorinated solvents: assessment of potential and limitations. Biodegradation, 15: 371-386.
- Grbic-Galic, D. and Vogel, T., 1987. Transformation of toluene and benzene by mixed methanogenic cultures. Applied and Environmental Microbiology, 53: 254-260.
- Green, C.T. and Scow, K.M., 2000. Analysis of phospholipid fatty acids (PLFA) to characterize microbial communities in aquifers. Hydrogeology Journal, 8(1): 126-141.
- Griebler, C., Adrian, L., Meckenstock, R.U. and Richnow, H.H., 2004a. Stable carbon isotope fractionation during aerobic and anaerobic transformation of trichlorobenzene. FEMS Microbiology Ecology, 48: 313-321.
- Griebler, C., Safinowski, M., Vieth, A., Richnow, H.H. and Meckenstock, R.U., 2004b. Combined application of stable carbon isotope analysis and specific metabolites determination for assessing in situ degradation of aromatic hydrocarbons in a tar oil-contaminated aquifer. Environmental Science and Technology, 38(2): 617-631.

Literatur

- Grossmann, E.L., Cifuentes, L.A. and Cozzarelli, I.M., 2002. Anaerobic Methane Oxidation in a Landfill-Leachate Plume. *Environmental Science and Technology*, 36(11): 2436-2442.
- Guckert, J.B., Antworth, C.P., Nichols, P.D. and White, D.C., 1985. Phospholipid, ester-linked fatty-acid profiles as reproducible assays for changes in prokaryotic community structure of estuarine sediments. *FEMS Microbiology Ecology*, 31: 147-158.
- Gumbrecht, T., 1993. Nutrient removal processes in fresh-water submersed macrophyte systems. *Ecological Engineering*, 2: 1-30.
- Hanson, J.R., Macalady, J.L., Harris, D. and Scow, K.M., 1999. Linking toluene degradation with specific microbial populations in soil. *Applied and Environmental Microbiology*, 65(12): 5403-5408.
- Harrington, R.R., Poulsen, S.R., Drever, J.I., Colberg, P.J.S. and Kelly, E.F., 1999. Carbon isotope systematics of monoaromatic hydrocarbons: vaporization and adsorption experiments. *Organic Geochemistry*, 30(8A): 765-775.
- Harwood, D.L. and Russell, N.J., 1984. Lipids in plants and microbes. George Allen and Unwin, London.
- He, Z. et al., 2007. GeoChip: a comprehensive microarray for investigating biogeochemical, ecological and environmental processes. *The ISME Journal*, 1: 67-77.
- Heider, J., Spormann, A.M., Beller, H.R. and Widdel, F., 1999. Anaerobic bacterial metabolism of hydrocarbons. *FEMS Microbiology Reviews*, 22(5): 459-473.
- Heidrich, S. et al., 2004a. Regionally contaminated aquifers - toxicological relevance and remediation options (Bitterfeld case study). *Toxicology*, 205(3): 143-155.
- Heidrich, S., Weiß, H. and Kaschl, A., 2004b. Attenuation reactions in a multiple contaminated aquifer in Bitterfeld (Germany). *Environmental Pollution*, 129(2): 277-288.
- Hendrickx, B. et al., 2005. Dynamics of an Oligotrophic Bacterial Aquifer Community during Contact with a Groundwater Plume Contaminated with Benzene, Toluene, Ethylbenzene, and Xylenes: An In Situ Mesocosm Study. *Applied and Environmental Microbiology*, 71(7): 3815-3825.
- Herrmann, S., Kleinstuber, S., Neu, T.R., Richnow, H.H. and Vogt, C., 2008. Enrichment of anaerobic benzene-degrading microorganisms by *in situ* microcosms. *Microbial Ecology*, 63: 94-106.
- Hessisches Landesamt für Umwelt und Geologie, 2004. Arbeitshilfe zu überwachten natürlichen Abbau- und Rückhalteprozessen im Grundwasser (Monitored Natural Attenuation MNA), Wiesbaden, Hessisches Landesamt für Umwelt und Geologie. Handbuch Altlasten, Bd.8, T.1.
- Hoefs, J., 1997. Stable Isotope Gechemistry. Springer-Verlag, Berlin, Heidelberg, 201 pp.
- Holliger, C., Schraa, G., Stams, A. and Zehnder, A., 1992. Enrichment and property of an anaerobic mixed culture reductively dechlorinating 1,2,3-trichlorobenzene to 1,3-dichlorobenzene. *Applied and Environmental Microbiology*, 58: 1636-1644.
- Holscher, T. et al., 2004. Multiple nonidentical reductive-dehalogenase-homologous genes are common in *Dehalococcoides*. *Applied and Environmental Microbiology*, 70: 5290-5297.
- Hunkeler, D., Anderson, N., Aravena, R., Bernasconi, S.M. and Butler, B.J., 2001. Hydrogen and carbon isotope fractionation during aerobic biodegradation of benzene. *Environmental Science and Technology*, 35(17): 3462-3467.
- Hunkeler, D., Aravena, R. and Butler, B.J., 1999a. Monitoring microbial dechlorination of tetrachloroethene (PCE) in groundwater using compound-specific stable carbon isotope ratios: microcosm and field studies. *Environmental Science and Technology*, 33(16): 2733-2738.

- Hunkeler, D., Aravena, R. and Cox, E., 2002. Carbon isotopes as tool to evaluate the origin and fate of vinyl chloride: Laboratory experiments and modeling of isotope evolution. *Environmental Science and Technology*, 36: 3378-3384.
- Hunkeler, D., Höhener, P., Bernasconi, S. and Zeyer, J., 1999b. Engineered in situ bioremediation of a petroleum hydrocarbon-contaminated aquifer: assessment of mineralization based on alkalinity, inorganic carbon and stable carbon isotope balances. *Journal of Contaminant Hydrology*, 37(3-4): 201-223.
- Imfeld, G. et al., 2008. Assessment of in situ degradation of chlorinated ethenes and bacterial community structure in a complex contaminated groundwater system. *Water Research*, 42(4-5): 871-882.
- Istok, J.D., Humphrey, M.D., Schroth, M.H., Hyman, M.R. and O'Reilly, K.T., 1997. Single-well, "push&pull" test for in situ determination of microbial activities. *Ground Water*, 35(4): 619-631.
- Istok, J.D. et al., 2004. In Situ Bioreduction of Technetium and Uranium in a Nitrate-Contaminated Aquifer. *Environmental Science and Technology*, 38: 468-475.
- ITVA, 2004. Monitored Natural Attenuation. Ingenieurtechnischer Verband Altlasten e.V., ITVA-Arbeitshilfe –H1-12, Dezember 2004, Berlin, 35 S.
- Jayachandran, G., Gorisch, H. and Adrian, L., 2003. Dehalorespiration with hexachlorobenzene and pentachlorobenzene by *Dehalococcoides* sp strain CBDB1. *Archives of Microbiology*, 180: 411-416.
- Johnsen, A.R., Winding, A., Karlson, U. and Roslev, P., 2002. Linking of Microorganisms to Phenanthrene Metabolism in Soil by Analysis of ¹³C-Labeled Cell Lipids. *Applied and Environmental Microbiology*, 68(12): 6106-6113.
- Jordahl, J.L., Foster, L., Schnoor, J.L. and Alvarez, P.J.J., 1997. Effect of hybrid popular trees on microbial populations important to hazardous waste bioremediation. *Environmental Toxicology Chemistry*, 16: 1318-1321.
- Jørgensen, K.S., 2007. In Situ Bioremediation. *Advances in Applied Microbiology*, 61: 285-305.
- Kanaly, R.A. and Harayama, S., 2000. Biodegradation of high-molecular-weight polycyclic aromatic hydrocarbons by bacteria. *Journal of Bacteriology*, 182(8): 2059-2067.
- Kaschl, A. et al., 2005. Isotopic fractionation indicates anaerobic monochlorobenzene biodegradation. *Environmental Toxicology and Chemistry*, 24(6): 1315-1324.
- Kästner, M. et al., 2006. Assessment of microbial in situ activity in contaminated aquifers. *Engineering in Life Science*, 6(3): 234-251.
- Kaur, A., Chaudhary, A., Kaur, A., Choudhary, R. and Kaushik, R., 2005. Phospholipid fatty acid - A bioindicator of environment monitoring and assessment in soil ecosystem. *Current Science*, 89(7): 1103-1112.
- Kopinke, F.D., Georgi, A., Voskamp, M. and Richnow, H.H., 2005. Carbon isotope fractionation of organic contaminants due to retardation on humic substances: implications for natural attenuation studies in aquifers. *Environmental Science and Technology*, 39(16): 6052-6062.
- KORA, 2007. Mikrobiologische NA-Untersuchungsmethoden: Fachliche Grundlagen für die Anwendung von Methoden zur Erfassung des natürlichen mikrobiellen Schadstoffabbaus im Aquifer / ProcessNet, Expertengruppe "Intrinsisches Abbaupotential" im Arbeitsausschuss "Ressourcenmanagement Boden und Grundwasser". - Stand: März 2007. Frankfurt M. DECHEMA, 2007. - 73 S.: Ill., graph. Darst.; 30 cm.
- KORA, 2008. Methodensammlung zur Untersuchung natürlicher Schadstoffminderungsprozesse. KORA-Handlungsempfehlungen, in press.

Literatur

- Kreuzer-Martin, H.W., 2007. Stable isotope probing: Linking functional activity to specific members of microbial communities. *Soil Science Society of America Journal*, 71(2): 611-619.
- Kube, M. et al., 2005. Genome sequence of the chlorinated compound-respiring bacterium *Dehalococcoides* species strain CBDB1. *Nature Biotechnology*, 23: 1269–1273.
- Kuder, T. et al., 2005. Enrichment of stable carbon and hydrogen isotopes during anaerobic biodegradation of MTBE: microcosm and field evidence. *Environmental Science and Technology*, 39(1): 213-220.
- Kuntze, K. et al., 2008. 6-Oxocyclohex-1-ene-1-carbonyl-coenzyme A hydrolases from obligately anaerobic bacteria: characterization and identification of its gene as a functional marker for aromatic compounds degrading anaerobes. *Environmental Microbiology*, in press, doi:10.1111/j.1462-2920.2008.01570.x.
- LABO, 2005. Berücksichtigung natürlicher Schadstoffminderungsprozesse bei der Altlastenbearbeitung -Bund-/Länderarbeitsgemeinschaft Bodenschutz (LABO) - Altlastenausschuss (ALA) Ad-hoc Unterausschuss „Natural Attenuation“, vom 01.06.2005.
- Lee, J.Y., Lee, J.Y.C., Lee, K.K., Lee, S.Y. and Lee, M.H., 2001. Statistical evaluation of geochemical parameter distribution in a ground water system contaminated with petroleum hydrocarbons. *Journal of Environmental Quality*, 35(5): 1548-1563.
- Liang, L.N. and Gribic-Galic, D., 1990. Anaerobic transformation of chlorobenzene and mixtures with aromatic acids by aquifer microorganisms and stable suspended consortia. *Abstracts of the Annual Meeting of the American Society for Microbiology*, 90: 198-202.
- Löffler, F.E., Cole, J.R., Ritalahti, K.M. and Tiedje, J.M., 2003. Diversity of dechlorinating bacteria. In: M.M. Häggblom and I.D. Bossert (Editors), *Microbial processes and environmental applications*. Kluwer Academic Publishers, Boston, pp. 53-87.
- Lollar, B.S. et al., 2001. Stable carbon isotope evidence for intrinsic bioremediation of tetrachloroethene and trichloroethene at area 6, Dover Air Force Base. *Environmental Science and Technology*, 35: 261-269.
- Lopez-Arilla, A.I., Gerard, E., Moreira, D. and Lopez-Garcia, P., 2004. Macrofilamentous microbial communities in the metal-rich and acidic River Tinto, Spain. *FEMS Microbiology Letters*, 235: 221-228.
- Lorah, M.M. and Olsen, L.D., 1999. Natural attenuation of chlorinated volatile organic compounds in a freshwater tidal wetland: field evidence of anaerobic biodegradation. *Water Resources Research*, 35: 3811-3827.
- Lord, C. and Church, T.A., 1983. Quantitative model for pyritization in salt-marsh sediments. *Estuaries*, 6: 295-296.
- Lösel, D.M., 1988. Fungal lipids. In: C. Ratledge and S.G. Wilkinson (Editors), *Microbial Lipids*. Academic Press, London, pp. 699-806.
- Lovley, D.R., 1991. Dissimilatory Fe(III) and Mn(IV) reduction. *Microbiological Reviews*, 55: 259-287.
- Lovley, D.R., 1995. Benzene oxidation coupled to sulfate reduction. *Applied and Environmental Microbiology*, 61: 953-958.
- Lovley, D.R., 1997. Microbial Fe(III) reduction in subsurface environments. *FEMS Microbiology Reviews*, 20: 305-313.
- Lovley, D.R., 2000. Anaerobic benzene degradation. *Biodegradation*, 11: 107-116.
- Lovley, D.R. et al., 1989. Oxidation of aromatic contaminants coupled to microbial iron reduction. *Nature*, 339: 297-300.

- Lovley, D.R. and Phillips, E.J.P., 1988. Novel mode of microbial energy metabolism: organic carbon oxidation coupled to dissimilatory reduction of iron or manganese. *Applied and Environmental Microbiology*, 54: 1472-1480.
- Lovley, D.R., Woodward, J.C. and Chapelle, F.H., 1994. Stimulated anoxic biodegradation of aromatic hydrocarbons using Fe(III) ligands. *Nature*, 370: 128-131.
- Lovley, D.R., Woodward, J.C. and Chapelle, F.H., 1996. Rapid anaerobic benzene oxidation with a variety of chelated Fe(III) forms. *Applied and Environmental Microbiology*, 62(1): 288-291.
- Lu, Y., Abraham, W.R. and Conrad, R., 2007. Spatial variation of active microbiota in the rice rhizosphere revealed by *in situ* stable isotope probing of phospholipid fatty acids. *Environmental Microbiology*, 9(2): 474-481.
- Lu, Y., Murase, J., Watanabe, A., Sugimoto, A. and Kimura, M., 2004. Linking microbial community dynamics to rhizosphere carbon flow in a wetland rice soil. *FEMS Microbiology Ecology*, 48: 179-186.
- Ludvigsen, L., Albrechtsen, H.J., Holst, H. and T.H., C., 1997. Correlating phospholipid fatty acids (PLFA) in a landfill leachate polluted aquifer with biogeochemical factors by multivariate statistical methods. *FEMS Microbiology Reviews*, 20: 447-460.
- Lueders, T., Manefield, M. and Friedrich, M.W., 2004. Enhanced sensitivity of DNA- and rRNA-based stable isotope probing by fractionation and quantitative analysis of isopycnic centrifugation gradients. *Environmental Microbiology*, 6(1): 73-78.
- Luther, G.W., Shellenbarger, P.A. and Brendel, P.J., 1996. Dissolved Fe(III) and Fe(II) complexes in salt marsh porewaters. *Geochimica et Cosmochimica Acta*, 60: 951-960.
- Macek, T., Mackova, M., Burkhard, J. and Demnerova, K., 1998. Introduction of green plants for the control of metals and organics in environmental remediation. In: F.W. Holm (Editor), *Effluents from Alternative Demilitarisation Technologies*. Kluwer Academic Publishers, Dordrecht, pp. 71-84.
- Madsen, E.L., 2000. Nucleic-acid characterization of the identity and activity of subsurface microorganisms. *Hydrogeology Journal*, 8: 112-125.
- Madsen, E.L., 2006. The use of stable isotope probing techniques in bioreactor and field studies on bioremediation. *Current Opinion in Biotechnology*, 17: 92-97.
- Major, D.W. et al., 2002. Field demonstration of successful bioaugmentation to achieve dechlorination of tetrachloroethene to ethene. *Environmental Science and Technology*, 36: 5106-5116.
- Makula, R.A., 1978. Phospholipid composition of methane utilizing bacteria. *Journal of Bacteriology*, 134: 771-777.
- Mancini, S.A. et al., 2002. Hydrogen isotopic enrichment: An indicator of biodegradation at a petroleum hydrocarbon contaminated field site. *Environmental Science and Technology*, 36(11): 2464-2470.
- Mancini, S.A. et al., 2003. Carbon and hydrogen isotopic fractionation during anaerobic biodegradation of benzene. *Applied and Environmental Microbiology*, 69(1): 191-198.
- Mandelbaum, R.T., Shati, M.R. and Ronen, D., 1997. *In situ* microcosms in aquifer bioremediation studies. *FEMS Microbiology Reviews*, 20: 489-502.
- Manefield, M., Whiteley, A.S. and Bailey, M.J., 2004. What can stable isotope probing do for bioremediation? *International Biodeterioration & Biodegradation*, 54: 163-166.

Literatur

- Manefield, M., Whiteley, A.S., Griffiths, R.I. and Bailey, M.J., 2002a. RNA stable isotope probing, a novel means of linking microbial community function to phylogeny. *Applied and Environmental Microbiology*, 68: 5367-5373.
- Manefield, M., Whiteley, A.S., Ostle, N., Ineson, P. and Bailey, M.J., 2002b. Technical considerations for RNA-based stable isotope probing: an approach to associating microbial diversity with microbial community function. *Rapid Communications in Mass Spectrometry*, 16: 2179-2183.
- Mariotti, A. et al., 1981. Experimental determination of nitrogen kinetic isotope fractionation: some principles, illustration for the denitrification and nitrification processes. *Plant and Soil*, 62: 413-430.
- Martin, H.M. and Ertl, S., 2005. Personal communication and presentation on the occasion of the DECHEMA Workshop 29.30.9.2005 at the FAL Braunschweig.
- Mauclaire, L., Pelz, O., Thullner, M., Abraham, W.R. and Zeyer, J., 2003. Assimilation of toluene carbon along a bacteria-protist food chain determined by C-13-enrichment of biomarker fatty acids. *Journal of Microbiological Methods*, 55(3): 635-649.
- McKelvie, J.R. et al., 2007. Evaluation of TCE and MTBE in situ biodegradation: integrating stable isotope, metabolic intermediate, and microbial lines of evidence. *Ground Water Monitoring & Remediation*, 27(4): 63-73.
- Meckenstock, R.U., Morasch, B., Griebler, C. and Richnow, H.H., 2004a. Stable isotope fractionation analysis as a tool to monitor biodegradation in contaminated aquifers. *Journal of Contaminant Hydrology*, 75(3-4): 215-255.
- Meckenstock, R.U., Morasch, B., Kästner, M., Vieth, A. and Richnow, H.H., 2002. Assessment of bacterial degradation of aromatic hydrocarbons in the environment by analysis of stable carbon isotope fractionation. *Water, Air and Soil Pollution Focus*, 2(3): 141-152.
- Meckenstock, R.U. et al., 1999. C-13/C-12 isotope fractionation of aromatic hydrocarbons during microbial degradation. *Environmental Microbiology*, 1(5): 409-414.
- Meckenstock, R.U., Safinowski, M. and Griebler, C., 2004b. Anaerobic degradation of polycyclic aromatic hydrocarbons. *FEMS Microbiology Ecology*, 49(1): 27-36.
- Meselson, M. and Stahl, F.W., 1958. The replication of DNA in *E. coli*. *Proceedings of the National Academy of Sciences of the United States of America*, 44: 671-682.
- Middeldorp, P., deWolf, J., Zehnder, A. and Schraa, G., 1997. Enrichment and properties of a 1,2,4-trichlorobenzene-dechlorinating methanogenic microbial consortium. *Applied and Environmental Microbiology*, 63: 1225-1229.
- Miltner, A., Richnow, H.H., Kopinke, F.D. and Kästner, M., 2004. Assimilation of CO₂ by soil microorganisms and transformation into soil organic matter. *Organic Geochemistry*, 35: 1015-1024.
- Mohn, W.W. and Tiedje, J.M., 1990. Strain DCB-1 conserves energy for growth from reductive dechlorination coupled to formate oxidation. *Archives of Microbiology*, 153(3): 267-271.
- Morasch, B., Höhener, P. and Hunkeler, D., 2007. Evidence for *in situ* degradation of mono- and polyaromatic hydrocarbons in alluvial sediments based on microcosm experiments with ¹³C-labeled contaminants. *Environmental Pollution*, 148(3): 739-748.
- Morasch, B., Richnow, H.H., Schink, B. and Meckenstock, R.U., 2001. Stable hydrogen and carbon isotope fractionation during microbial toluene degradation: Mechanistic and environmental aspects. *Applied Environmental Microbiology*, 67: 4842-4849.

- Morasch, B., Richnow, H.H., Schink, B., Vieth, A. and Meckenstock, R.U., 2002. Carbon and hydrogen stable isotope fractionation during aerobic bacterial degradation of aromatic hydrocarbons. *Applied Environmental Microbiology*, 68: 5191-5194.
- Morasch, B., Richnow, H.H., Vieth, A., Schink, B. and Meckenstock, R.U., 2004. Stable isotope fractionation caused by glycyl radical enzymes during bacterial degradation of aromatic compounds. *Applied and Environmental Microbiology*, 70(5): 2935-2940.
- Mordaunt, C.J., Gevao, B., Jones, K.C. and Semple, K.T., 2005. Formation of non-extractable pesticide residues: observations on compound differences, measurement and regulatory issues. *Environmental Pollution*, 133: 25-34.
- Morrill, P.L. et al., 2005. Quantifying chlorinated ethene degradation during reductive dechlorination at Kelly AFB using stable carbon isotopes. *Journal of Contaminant Hydrology*, 76: 279-293.
- Mottram, H. and Evershed, R.P., 2003. Practical considerations in GC/C/IRMS of ^{13}C -enriched compounds. *Rapid Communication Mass Spectrometry*, 17: 2669-2674.
- Neufeld, J. et al., 2007. DNA stable-isotope probing. *Nature Protocols*, 2(4): 860-866.
- Nichols, P.D., Mancuso, C.A. and White, D.C., 1987. Measurement of methanotroph and methanogen signature phospholipids for use in assessment of biomass and community structure in model systems. *Organic Geochemistry*, 11: 451-461.
- Nichols, P.D., Smith, G.A., Antworth, C.P., Hanson, R.S. and White, D.C., 1985. Phospholipid and lipopolysaccharide normal and hydroxyl fatty acids as potential signatures for the methaneoxidizing bacteria. *FEMS Microbiology Ecology*, 31: 327-335.
- Nijenhuis, I. et al., 2005. Stable isotope fractionation of tetrachloroethene during reductive dechlorination by *Sulfurospirillum multivorans* and *Desulfitobacterium* sp. strain PCE-S and abiotic reactions with cyanocobalamin. *Applied and Environmental Microbiology*, 71(7): 3413-3419.
- Nijenhuis, I., Stelzer, N., Kästner, M. and Richnow, H.H., 2007. Sensitive detection of anaerobic monochlorobenzene degradation using stable isotope tracers. *Environmental Science and Technology*, 41(11): 3836-3842.
- Nikolausz, M., Nijenhuis, I., Ziller, K., Richnow, H.H. and Kästner, M., 2006. Stable carbon isotope fractionation during degradation of dichloromethane by methylotrophic bacteria. *Environ. Microbiol.* 8. *Environmental Microbiology*, 8: 156-164.
- North, N.N. et al., 2004. Change in bacterial community structure during in situ biostimulation of subsurface sediment cocontaminated with uranium and nitrate. *Applied and Environmental Microbiology*, 70: 4911-4920.
- Nowak, J., Kirsch, N.H., Hegemann, W. and Stan, H.-J., 1996. Total reductive dechlorination of chlorobenzenes to benzene by a methanogenic mixed culture enriched from Saale river sediment. *Applied Microbiology and Biotechnology*, 45: 700-709.
- O'leary, W.M. and Wilkinson, S.G., 1988. Gram-positive bacteria. In: C. Ratledge and S.G. Wilkinson (Editors), *Microbial lipids*. Academic Press, London, pp. 117-201.
- Pardue, J.H., Kongara, W.A. and Jones, W.J., 1996. Effect of cadmium on reductive dechlorination of trichloroaniline. *Environmental Toxicology Chemistry*, 15: 1083-1088.
- Parker, J.H., Smith, G.A., Fredrickson, H.L., Vestal, J.R. and White, D.C., 1982. Sensitive assay based on hydroxy fatty acids from lipopolysaccharide lipid A, for Gram-negative bacteria in sediments. *Environmental Microbiology*, 44: 1170-1177.
- Parkes, R.J. and Calder, A.G., 1985. The cellular fatty acids of three strains of *Desulfobulbus*, a priopionate-utilising sulphatereducing bacterium. *FEMS Microbiology Ecology*, 31: 361-363.

Literatur

- Peacock, A.D. et al., 2004. Utilization of microbial biofilms as monitors of bioremediation. *Microbial Ecology*, 47: 284-292.
- Pelz, O. et al., 2001a. Use of isotopic and molecular techniques to link toluene degradation in denitrifying aquifer microcosms to specific microbial populations. *Archives of Microbiology*, 175(4): 270-281.
- Pelz, O., Chatzinotas, A., Zarda-Hess, A., Abraham, W.R. and Zeyer, J., 2001b. Tracing toluene-assimilating sulfate-reducing bacteria using C-13-incorporation in fatty acids and whole-cell hybridization. *FEMS Microbiology Ecology*, 38(2-3): 123-131.
- Peter, A. et al., 2004. Assessing microbial degradation of o-xylene at field-scale from the reduction in mass flow rate combined with compound-specific isotope analyses. *Journal of Contaminant Hydrology*, 71: 127-154.
- Peters, S., Koschinsky, S., Schwieger, F. and Tebbe, C.C., 2000. Succession of microbial communities during hot composting as detected by PCR-SSCP-based genetic profiles of small subunit rRNA genes. *Applied and Environmental Microbiology*, 66: 930-936.
- Pfeifer, F., 2004. Bedeutung und Möglichkeiten zur Stimulierung der natürlichen Selbstreinigungs Kräfte in kontaminierten Grundwässern ehemaliger Bergbaustandorte. DMT-Berichte aus Forschung und Entwicklung, 158.
- Pfeifer, F., Podwojewski, E., Quecke, W. and Richnow, H.H., 2003. Charakterisierung des natürlichen biologischen Abbaus von BTEX im Grundwasser des ehemaligen Kokereistandortes Gneisenau 1/2/3/4 mittels Isotopenfraktionierung. Altlastenspektrum, 3: 126-135.
- Phelps, C.D., Zhang, X. and Young, L.Y., 2001. Use of stable isotopes to identify benzoate as a metabolite of benzene degradation in a sulphidogenic consortium. *Environmental Microbiology*, 3: 600-603.
- Phillips, T.M., Seech, A.G., Lee, H. and Trevors, J.T., 2005. Biodegradation of hexachlorocyclohexane (HCH) by microorganisms. *Biodegradation*, 16(4): 363-392.
- Plumb JR, R.H. and Pitchford, A.M., 1985. Volatile organic scans: implications for groundwater monitoring, Proceedings of the National Water Well Association/American Petroleum Institute Conference on Petroleum Hydrocarbons and Organic Chemicals in Groundwater, Houston (Texas), pp. 1-15.
- Pombo, S.A., Pelz, O., Schroth, M.H. and Zeyer, J., 2002. Field-scale C-13-labeling of phospholipid fatty acids (PLFA) and dissolved inorganic carbon: tracing acetate assimilation and mineralization in a petroleum hydrocarbon-contaminated aquifer. *FEMS Microbiology Ecology*, 41(3): 259-267.
- Popp, P., Bruggemann, L., Keil, P., Thuss, U. and Weiss, H., 2000. Chlorobenzenes and hexachlorocyclohexanes (HCHs) in the atmosphere of Bitterfeld and Leipzig (Germany). *Chemosphere*, 41(6): 849-855.
- Radajewski, S., Ineson, P., Parekh, N.R. and Murrell, J.C., 2000. Stable-isotope probing as a tool in microbial ecology. *Nature*, 403: 646-649.
- Radajewski, S., McDonald, I.R. and Murrell, J.C., 2003. Stable-isotope probing of nucleic acids: a window to the function of uncultured microorganisms. *Current Opinion in Biotechnology*, 14(3): 296-302.
- Ramsburg, C.A. et al., 2004. Stimulated microbial reductive dechlorination following surfactant treatment at the Bachman road site. *Environmental Science and Technology*, 38: 5902-5914.
- Reusser, D.E. and Field, J.A., 2002. Determination of benzylsuccinic acid in gasoline-contaminated groundwater by solid-phase extraction coupled with gas chromatography-mass spectrometry. *Journal of Chromatography A*, 953: 215-225.

- Richnow, H.H., Annweiler, E., Michaelis, W. and Meckenstock, R.U., 2003a. Microbial in situ degradation of aromatic hydrocarbons in a contaminated aquifer monitored by carbon isotope fractionation. *Journal of Contaminant Hydrology*, 65(1-2): 101-120.
- Richnow, H.H. et al., 1999a. Formation of nonextractable soil residues: A stable isotope approach. *Environmental Science and Technology*, 33: 3761-3767.
- Richnow, H.H. et al., 1998. The use of ^{13}C -labeled polycyclic aromatic hydrocarbons for the analysis of their transformation in soils. *Chemosphere*, 36: 2211-2224.
- Richnow, H.H. and Meckenstock, R.U., 1999b. Isotopen-geochemisches Konzept zur In-situ-Erfassung des biologischen Abbaus in kontaminiertem Grundwasser. *TerraTech*: 38-41.
- Richnow, H.H. et al., 2003b. In situ biodegradation determined by carbon isotope fractionation of aromatic hydrocarbons in an anaerobic landfill leachate plume (Vejen, Denmark). *Journal of Contaminant Hydrology*, 64(1-2): 59-72.
- Ringelberg, D.B., Sutton, S. and White, D.C., 1997. Biomass, bioactivity and biodiversity: microbial ecology of the deep subsurface: analysis of esterlinked phospholipid fatty acids. *FEMS Microbiology Reviews*, 20: 371-377.
- Rittmann, B.E., 2004. Definition, objectives, and evaluation of natural attenuation. *Biodegradation*, 15: 349-357.
- Rosell, M. et al., 2007a. Variations in $^{13}\text{C}/^{12}\text{C}$ and D/H enrichment factors of aerobic bacterial fuel oxygenate degradation. *Environmental Science and Technology*, 41(6): 2036-2043.
- Rosell, M., Häggblom, M.M. and Richnow, H.H., 2007b. Compound specific isotope analysis (CSIA) to characterise degradation pathways and to quantify in situ degradation of fuel oxygenates and other fuel derived contaminants. In: D. Barceló (Editor), *Fuel Oxygenates, The Handbook of Environmental Chemistry*. Springer-Verlag, Berlin, Heidelberg, Germany, pp. 99-119.
- Rügge, K., Bjerg, P.L., Pedersen, J.K., Mosbaek, H. and Christensen, T.H., 1999. An anaerobic field injection experiment in a landfill leachate plume, Grindsted, Denmark: 1. Experimental setup, tracer movement, and fate of aromatic and chlorinated compounds. *Water Resources Research*, 35(4): 1231-1246.
- Scheuerbrandt, G. and Bloch, K., 1962. Unsaturated fatty acids in microorganisms. *Journal of Biological Chemistry*, 237: 2064–2068.
- Scheunert, I., Vockel, D. and Schmitzer, J., 1987. Biomineralization rates of ^{14}C -labelled organic chemicals in aerobic and anaerobic suspended soil. *Chemosphere*, 16: 1031-1041.
- Schiedeck, T., Grathwohl, P. and Teutsch, G., 1997. Literaturstudie zum natürlichen Rückhalt/Abbau von Schadstoffen im Grundwasser. Texte und Berichte zur Altlastenbearbeitung 35/37, 35/37. State Agency for Environmental Protection Baden-Württemberg, Karlsruhe, Germany.
- Schildkraut, C.L., 1967. The preparation and properties of 15N-deuterated DNA from bacteria. In: L. Grossman and K. Moldave (Editors), *Methods in Enzymology*. Academic Press, New York, pp. 695-699.
- Schirmer, M. et al., 2006. Natural attenuation research at the contaminated megasite Zeitz. *Journal of Hydrology*, 328(3-4): 393-407.
- Schirmer, M., Jones, I., Teutsch, G. and Lerner, D.N., 1995. Development and testing of multi-level sock samplers for groundwater. *Journal of Hydrology*, 171(3-4): 239-257.
- Schlegel, H.G., 1992. *Allgemeine Mikrobiologie*. Thieme Verlag, Stuttgart, New York.
- Schmalenberger, A., Schwieger, F. and Tebbe, C.C., 2001. Effect of primers hybridizing to different evolutionarily conserved regions of the small subunit rRNA gene in PCR-based microbial

Literatur

- community analyses and genetic profiling. *Applied and Environmental Microbiology*, 67: 3557-3563.
- Schmidt, I. et al., 2003. New concepts of microbial treatment processes for the nitrogen removal in wastewater. *FEMS Microbiology Reviews*, 27(4): 481-492.
- Schmidt, T.C., Morgenroth, E., Schirmer, M., Effenberger, M. and Haderlein, S.B., 2002. Use and occurrence of fuel oxygenates in Europe. In: A.F. Diaz and D.L. Drogos (Editors), *Oxygenates in Gasoline: Environmental Aspects*. ACS, Washington, pp. 58- 79.
- Schmidt, T.C. et al., 2004. Compound-specific stable isotope analysis of organic contaminants in natural environments: a critical review of the state of the art, prospects, and future challenges. *Analytical and Bioanalytical Chemistry*, 378(2): 283-300.
- Schnoor, J.L., Licht, L.A., McCutcheon, S.C., Wolfe, N.L. and Carreira, L.H., 1995. Phytoremediation of organic and nutrient contaminants. *Environmental Science and Technology*, 29(7): 318-323.
- Schroth, M.H., Kleikemper, J., Bolliger, C., Bernasconi, S.M. and Zeyer, J., 2001. In situ assessment of microbial sulfate reduction in a petroleum-contaminated aquifer using push-pull tests and stable sulfur isotope analyses. *Journal of Contaminant Hydrology*, 51: 179-195.
- Schüth, C., Taubald, H., Bolano, N. and Maciejczyk, K., 2003. Carbon and hydrogen isotope effects during sorption of organic contaminants on carbonaceous materials. *Journal of Contaminant Hydrology*, 64(3-4): 269-281.
- Schwieger, F. and Tebbe, C.C., 1998. A new approach to utilize PCR-single-strand conformation polymorphism for 16S rRNA gene-based microbial community analysis. *Applied and Environmental Microbiology*, 64: 4870-4876.
- Schwieger, F. and Tebbe, C.C., 2000. Effect of field inoculation with *Sinorhizobium meliloti* L33 on the composition of bacterial communities in rhizospheres of a target (*Medicago sativa*) and a non-target plant (*Chenopodium album*) - Linking 16S rRNA gene based SSCP-community profiles and diversity of cultivated isolates. *Applied and Environmental Microbiology*, 66: 3556-3565.
- Scow, K.M. and Hicks, K.A., 2005. Natural attenuation and enhanced bioremediation of organic contaminants in groundwater. *Current Opinion in Biotechnology*, 16(3): 246-253.
- Sherwood Lollar, B. and Slater, G.-F., 2001. Stable carbon isotope evidence for intrinsic bioremediation of tetrachlorethene and trichlorethene at Area 6, Dover Air Force Base. *Environmental Science and Technology*, 35(2): 261-269.
- Sherwood Lollar, B. et al., 1999. Contrasting carbon isotope fractionation during biodegradation of trichloroethylene and toluene: implications for intrinsic bioremediation. *Organic Geochemistry*, 30: 813-820.
- Shimp, J.F. et al., 1993. Beneficial effects of plants in the remediation of soil and groundwater contaminated with organic materials. *Environmental Science and Technology*, 23: 41-77.
- Slater, G.F., Ahad, J.M.E., Lollar, B.S., Allen-King, R. and Sleep, B., 2000. Carbon isotope effects resulting from equilibrium sorption of dissolved VOCs. *Analytical Chemistry*, 72(22): 5669-5672.
- Søgaard, E.G., Remigijus, A., Abraham-Peskir, J. and Koch, B.C., 2001. Conditions for biological precipitation of iron by *Gallionella ferruginea* in a slightly polluted groundwater. *Applied Geochemistry*, 16: 1129-1137.
- Song, D.L., Conrad, M.E., Sorenson, K.S. and Alvarez-Cohen, L., 2002. Stable carbon isotope fractionation during enhanced in situ bioremediation of trichloroethene. *Environmental Science and Technology*, 36: 2262-2268.

- Spiegelman, D., Whissel, G. and Greer, C.W., 2005. A survey of the methods for the characterization of microbial consortia and communities. Canadian Journal of Microbiology, 51(5): 355-386.
- Spormann, A.M. and Widdel, F., 2000. Metabolism of alkylbenzenes, alkanes, and other hydrocarbons in anaerobic bacteria. Biodegradation, 11(2-3): 85-105.
- Stein, L.Y. et al., 2002. Intriguing microbial diversity associated with metal-rich particles from a freshwater reservoir. FEMS Microbiology Ecology, 42: 431-440.
- Steinbach, A., Seifert, R., Annweiler, E. and Michaelis, W., 2004. Hydrogen and carbon isotope fractionation during anaerobic biodegradation of aromatic hydrocarbons – A field study. 38: 609-616.
- Stelzer, N. et al., 2006a. In situ microcosms (BACTRAP) to evaluate natural attenuation potentials in contaminated aquifers. Organic Geochemistry, 37(10): 1394-1410.
- Stelzer, N., Fischer, A., Kästner, M. and Richnow, H.H., 2006b. Analyse des anaeroben *in situ* Benzolabbaus: Vergleich von *In-situ*-Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen. Grundwasser, 11(4): 247-258.
- Stelzer, N. et al., 2008. Integrative approach to delineate natural attenuation of chlorinated benzenes in anoxic aquifers. Environmental Science and Technology, submitted.
- Sundh, I., Borga, P., Nilsson, M. and Svensson, B.H., 1995. Estimation of cell numbers of methanotrophic bacteria in boreal peatlands based on analysis of specific phospholipid fatty acids. FEMS Microbiology Ecology, 18: 103-112.
- Taylor, J. and Parkes, R.J., 1983. The cellular fatty acids of the sulphate-reducing bacteria Desulfobacter sp., Desulfovibulus sp. and Desulfovibrio desulfuricans. Journal of General Microbiology, 129(11): 3303-3309.
- Tebbe, C.C., Schmalenberger, A., Peters, S. and Schwieger, F., 2001. Single-strand conformation polymorphism (SSCP) for microbial community analysis. In: P.A. Rochelle (Editor), Environmental Molecular Microbiology: Protocols and Applications. Horizon Scientific Press, Wymondham, UK, pp. 161-175.
- Teutsch, G., Grathwohl, P. and Schiedek, C., 1997. Literaturstudie zum natürlichen Rückhalt/Abbau von Schadstoffen im Grundwasser, Handbuch Altlasten und Grundwasserschadensfälle. Landesanstalt für Umweltschutz Baden-Württemberg, Karlsruhe.
- Thauer, R.K., Jungermann, K. and Decker, K., 1977. Energy conservation in chemotrophic anaerobic bacteria. Bacteriological Reviews, 41: 100-180.
- Thiel, V. et al., 2001. Molecular signals for anaerobic methane oxidation in Black Sea seep carbonates and a microbial mat. Marine Chemistry, 73(2): 97-112.
- Thierrin, J., Davis, G.B. and Barber, C., 1995. A ground water tracer test with deuterated compounds for monitoring *in situ* biodegradation and retardation of aromatic hydrocarbons. Ground Water. Ground Water, 3: 469-475.
- Thulliner, M. and Schäfer, W., 1999. Modeling of a field experiment on bioremediation of chlorobenzenes in groundwater. Bioremediation Journal, 3(3): 247-267.
- Trapp, S., 2000. Aspekte der Phytoremediation organischer Schadstoffe. USWF - Z. Umweltchemie und Ökotoxikologie, 12(5): 246-255.
- Tratnyek, P. et al., 2001. Visualizing redox chemistry: probing environmental oxidation-reduction reactions with indicator dyes. Chemical Ed, 6(3): 172-179.
- Treonis, A.M. et al., 2004. Identification of groups of metabolically-active rhizosphere microorganisms by stable isotope probing of PLFAs. Soil Biology & Biochemistry, 36: 533-537.

Literatur

- Tunlid, A. and White, D.C., 1992. Biochemical analysis of biomass, community structure, nutritional status and metabolic activity of microbial community in soil. In: C. Stotzky and J.M. Bollag (Editors). Marcel Dekker, New York, pp. 229-262.
- Ulrich, A.C., Beller, H.R. and Edwards, E.A., 2005. Metabolites detected during biodegradation of $^{13}\text{C}_6$ -benzene in nitrate-reducing and methanogenic enrichment cultures. Environmental Science and Technology, 39(17): 6681-6691.
- Ulrich, A.C. and Edwards, E.A., 2003. Physiological and molecular characterization of anaerobic benzene-degrading mixed cultures. Environmental Microbiology, 5(2): 92-102.
- Umweltbundesamt, 2007. Bundesweite Übersicht zur Altlastenstatistik. Stand 06/2007.
- US-EPA, 1999. Use of monitored natural attenuation at superfund, RCRA corrective action and underground storage tank sites. Directive Number: 9200.4-17P.
- US-EPA, 2004. Performance monitoring of MNA remedies for VOCs in ground water. EPA/600/R-04/027, April 2004.
- Van Agteren, M.H., Keuning, S. and Janssen, D.B., 1998. Handbook on biodegradation and biological treatment of hazardous organic compounds. Kluwer Academic Publishers, Dordrecht.
- Van de Pas, B.A. et al., 1999. Purification and molecular characterization of ortho-chlorophenol reductive dehalogenase, a key enzyme of halorespiration in *Desulfobacterium dehalogenans*. J Biol Chem, 274(29): 20287-20292.
- Van Doesburg, W. et al., 2005. Reductive dechlorination of beta-hexachlorocyclohexane (beta-HCH) by a *Dehalobacter* species in coculture with a *Sedimentibacter* sp. FEMS Microbiol Ecol, 54(1): 87-95.
- Van Eekert, M.H.A. et al., 1998. Anaerobic transformation of beta-hexachlorocyclohexane by methanogenic granular sludge and soil microflora. Environmental Science and Technology, 32: 3299-3304.
- Vestal, J.R. and White, D.C., 1989. Lipid analysis in microbial ecology. Bioscience, 39: 535-541.
- Vieth, A., Kästner, M., Morasch, B., Meckenstock, R.U. and Richnow, H.H., 2001. Charakterisierung des biologischen Abbaus von BTEX im Grundwasser über Isotopenfraktionierung - Feldstudien. TerraTech, 10(5): 37-41.
- Vieth, A. et al., 2005. Monitoring in situ biodegradation of benzene and toluene by stable carbon isotope fractionation. Environmental Toxicology and Chemistry, 24(1): 51-60.
- Vieth, A. et al., 2003. In-situ biodegradation of tetrachloroethene and trichloroethene in contaminated aquifers monitored by stable isotope fractionation. Isotopes in Environmental and Health Studies, 39: 113-124.
- Vogt, C. et al., 2002. Two pilot plant reactors designed for the in situ bioremediation of chlorobenzene-contaminated ground water: hydrogeological and chemical characteristics and bacterial consortia. Water, Air and Soil Pollution, 2: 161-170.
- Vogt, C., Simon, D., Alfreider, A. and Babel, W., 2004. Microbial degradation of chlorobenzene under oxygen-limited conditions leads to accumulation of 3-chlorocatechol. Environmental Toxicology and Chemistry, 23: 265-270.
- Wachter, T., 2004. Geostatistische Analyse der Variabilität hydrogeochemischer Parameter und Quantifizierung der NA-Kapazität am Beispiel des BTEX-kontaminierten Aquifers in Zeitz / Sachsen-Anhalt. Dissertation Thesis, Christian-Albrechts-Universität, Kiel, 98 pp.

- Wagner, M., Nielsen, P.H., Loy, A., Nielsen, J.L. and Daims, H., 2006. Linking microbial community structure with function: fluorescence *in situ* hybridization-microautoradiography and isotope arrays. *Current Opinion in Biotechnology*, 17: 83-91.
- Warren, L.A. and Haack, E.A., 2001. Biological control on metal behaviour in freshwater environments. *Earth-Science Review*, 44: 261-320.
- Weiss, H., Schirmer, M., Teutsch, G. and Merkel, P., 2001. Sanierungsforschung in regional kontaminierten Aquiferen (SAFIRA)-2. Projektüberblick und Pilotanlage. *Grundwasser*, 3: 135-139.
- Weiss, H., Schirmer, M., Teutsch, G. and Merkel, P., 2002. Sanierungsforschung in regional kontaminierten Aquiferen (SAFIRA) - 2. Projektübersicht und Pilotanlage. *Grundwasser*, 3: 135-139.
- Weiss, J.V. and Cozzarelli, I.M., 2008. Biodegradation in Contaminated Aquifers: Incorporating Microbial/Molecular Methods. *Ground Water*, in press.
- White, D.C., 1988. Validation of quantitative analysis for microbial biomass, community structure, and metabolic activity. *Advanced Limnology*, 31: 1-18.
- White, D.C., 1994. Is there anything else you need to understand about the microbiota that cannot be derived from analysis of nucleic acids? *Microbial Ecology*, 28: 163-166.
- White, D.C. et al., 2003. Forensic analysis by comprehensive rapid detection of pathogens and contamination concentrated in biofilms in drinking water systems for water resource protection and management. *Environmental Forensics*, 4(1): 63-74.
- White, D.C. et al., 1983. The groundwater aquifer microbiota: biomass, community structure, and nutritional status. *Developments in Industrial Microbiology*, 24: 201–211.
- White, D.C., Stair, J.O. and Ringelberg, D.B., 1996. Quantitative comparisons of *in situ* microbial biodiversity by signature biomarker analysis. *Journal of Industrial Microbiology & Biotechnology*, 17: 185-196.
- Whiteley, A.S., Manefield, M. and Lueders, T., 2006. Unlocking the microbial black box using RNA-based stable isotope probing technologies. *Current Opinion in Biotechnology*, 17(1): 67-71.
- Whiteley, A.S., Thomson, B., Lueders, T. and Manefield, M., 2007. RNA stable-isotope probing. *Nature Protocols*, 2(4): 838-844.
- Whiticar, M.J., 1999. Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane. *Chemical Geology*, 161(1-3): 291-314.
- WHO, 2006. Guidelines for Drinking-water Quality: incorporating first addendum., Vol. 1, Recommendations. - 3rd ed.
- Widada, J., Nojiri, H. and Omori, T., 2002. Recent developments in molecular techniques for identification and monitoring of xenobiotic-degrading bacteria and their catabolic genes in bioremediation. *Applied Microbiology and Biotechnology*, 60: 45-59.
- Widdel, F. and Rabus, R., 2001. Anaerobic biodegradation of saturated and aromatic hydrocarbons. *Current Opinion in Biotechnology*, 12(3): 259-276.
- Wiedemeier, T.H., Rifai, H.S., Newell, C.J. and Wilson, J.T., 1999. Natural attenuation of fuels and chlorinated solvents in the subsurface. John Wiley & Sons, INC., New York, 617 pp.
- Wilkinson, S.G., 1988. Gram-negative bacteria. In: C. Ratledge and S.G. Wilkinson (Editors), *Microbial lipids*. Academic Press, London, pp. 299-488.

Literatur

- Winderl, C., Schaefer, S. and Lueders, T., 2007. Detection of anaerobic toluene and hydrocarbon degraders in contaminated aquifers using benzylsuccinate synthase (*bssA*) genes as a functional marker. *Environmental Microbiology*, 9(4): 1035-1046.
- Wöstmann, U., 2007. Natürliche Selbstreinigung und Immobilisierung bei schädlichen Bodenveränderungen und Altlasten: Praxishandbuch zur Sanierungsprüfung. Bodenschutz & Altlasten; Bd. 15. Erich Schmidt Verlag GmbH & Co., Berlin.
- Wycisk, P., Weiss, H. and Kaschl, A., 2004. Groundwater pollution and remediation options for multisource-contaminated aquifer (Bitterfeld/Wolfen, Germany). *Toxicology Letters*, 140: 343-351.
- Zelles, L., 1999. Fatty acid patterns of phospholipids and lipopolysaccharides in the characterisation of microbial communities in soil: A review. *Biology and Fertility of Soils*, 29: 111-129.
- Zengler, K., Richnow, H.H., Rossello-Mora, R., Michaelis, W. and Widdel, F., 1999. Methane formation from long-chainalkanes by anaerobic microorganisms. *Nature*, 401: 266-269.
- Zhang, C.L., 2002. Stable carbon isotopes of lipid biomarkers: analysis of metabolites and metabolic fates of environmental microorganisms. *Current Opinion in Biotechnology*, 13(1): 25-30.
- Zhang, C.L. et al., 2003. Carbon isotope signatures of fatty acids in *Geobacter metallireducens* and *Shewanella algae*. *Chemical Geology*, 195(1-4): 17-28.
- Zhou, J.Z., 2003. Microarrays for bacterial detection and microbial community analysis. *Current Opinion in Microbiology*, 6: 288-294.
- Zhou, J.Z., Fries, M.R., Chee-Sanford, J.C. and Tiedje, J.M., 1995. Phylogenetic analyses of a new group of denitrifiers capable of anaerobic growth on toluene and description of *azoorcus-tolulyticus* sp-nov. *International Journal of Systematic Bacteriology*, 45: 500-506.
- Zinder, S.H., 1998. Methanogens. In: R.S. Burlage, R. Atlas, D. Stahl, G. Geesey and G. Sayler (Editors), *Techniques in Microbial Ecology*. Oxford University Press, New York, pp. 113-136.
- Zwank, L. et al., 2005. New evaluation scheme for two-dimensional isotope analysis to decipher biodegradation processes: Application to groundwater contamination by MTBE. *Environmental Science and Technology*, 39: 1018-1029.

10 Acknowledgements

Such a work would not be possible without the help, support, guidance, friendship and understanding of many people to whom I am very grateful. It would be impossible to name them all, but nevertheless I like to acknowledge some persons in particular.

Many thanks especially go to Hans Richnow for his supervision of this thesis and also for all his support and critical discussions during the last 5 years. Special thanks also go to Prof. Andreas Dahmke for good cooperation and for giving me the chance to submit and defend my thesis at the Institute of Applied Geosciences within the Faculty of Mathematics and Natural Sciences at the Christian-Albrecht University in Kiel. Further, I like to acknowledge the Deutsche Bundesstiftung Umwelt (20004/751) for financial support of this work and the organisation of interesting seminars at wonderful locations.

Since the first day I enjoyed my time in the UFZ within the Department of Isotope Biogeochemistry and in this respect I like to thank all my colleagues and friends especially from the ISOBIO, BIOREM, UMB departments who made everyday life at work but also after work really delightful. I think we are a great team from all over the world. But beside the fun we had I also appreciate the time you spend to help me in the laboratory, with your discussions and feedback whenever I needed it. Especially, I like to acknowledge Anko Fischer and Ivonne Nijenhuis for all their critical and fruitful discussions and help with corrections of manuscripts, abstracts, presentations etc.; Prof. Dr. Matthias Kästner for his support and cooperation; Matthias Gehre, Ursula Günther, Kerstin Ethner and Ines Mäusezahl to keep the things running in the laboratories; Liane Paul our wonderful secretary; all students who came to work with me during an internship or for a diploma thesis; my colleagues from the wonderful AXIOM group, namely Paula Martinez, Gwael Imfeld, Danuta Cichocka, Makeba Kampara for not only sharing cookies and coffee in the office; the team in Bitterfeld, Werner Kletzander and Hansjörg Weiß for technical assistance in the field; Stefanie Weber, Petra Bombach, Carsten Vogt, Reimo Kindler and many others more.

My sincere thanks go to my family and closest friends who were always there if I needed them. And in particular I want to say thank you to Christian, Lei and Paula!

Eidesstattliche Erklärung

11 Eidesstattliche Erklärung

Hiermit erkläre ich, dass die vorliegende Arbeit über die *Entwicklung eines Testsystems zum Nachweis der in situ Biodegradation im Grundwasser* von mir selbstständig angefertigt wurde, abgesehen von der Beratung durch meine Betreuer und unter Verwendung der entsprechend gekennzeichneten Hilfsmittel und Literaturangaben. Mit Ausnahme der benannten Teile ist die Arbeit noch nicht veröffentlicht und wurde bisher nicht an anderer Stelle im Rahmen eines Prüfungsverfahrens eingereicht.

Die Arbeit ist unter Einhaltung der Regeln guter wissenschaftlicher Praxis entstanden.

Leipzig, 20.05.2008

12 Persönlicher Werdegang

Name: Nicole Stelzer
Geburtsdatum: 17.06.1979
Geburtsort: Borna
Familienstand: ledig
Nationalität: deutsch

BILDUNGSWEG

seit 01/2005	Promotion mit Stipendium der Deutschen Bundesstiftung Umwelt (DBU) Doktorandin am Helmholtz-Zentrum für Umweltforschung Leipzig im Department für Isotopengeochemie und der Arbeitsgruppe Angewandte Geologie an der Christian-Albrechts-Universität zu Kiel
10/1998-06/2004	Universität Leipzig, Institut für Geophysik und Geologie Studium im Fach Geologie/ Paläontologie Abschluss des Studiums mit der Note 1,4 08/2000 Vordiplom mit der Note 1,6
10/2003-04/2004	Diplomarbeit am Helmholtz-Zentrum für Umweltforschung Leipzig, Department Bioremediation Thema: In-situ Kosmen zur Erfassung des mikrobiologischen Abbaus verschiedener geochemischer Zonen eines BTEX-kontaminierten Aquifers
09/1992-07/1998	„Wiprecht-Gymnasium“ Groitzsch Abitur 1998, Prädikat: 1,7
09/1987-07/1992	Oberschule „Otto Grotewohl“ Neukieritzsch

PRAKТИСHE TÄTIGKEITEN

09/2007-12/2007	Isodetect GmbH
06/2007	Knauf Gips KG Iphofen, Abteilung Rohstoffsicherung Geologische Kartierung eines Gipsvorkommens in Spanien
09/2005	Knauf Gips KG Iphofen, Abteilung Rohstoffsicherung Geologische Kartierung eines Gipsvorkommens in Spanien
07/2004-12/2004	Helmholtz-Zentrum für Umweltforschung Wissenschaftliche Mitarbeit im Department Bioremediation
01/2003-03/2003	Knauf Gips KG Iphofen, Abteilung Rohstoffsicherung Erkundung eines Gipsvorkommens auf Sizilien: Geologische Kartierung, Lagerstättenkundliche Auswertung, Abbauplanung und Kostenschätzung
06/2001-12/2003	FUGRO CONSULT GmbH Bearbeitung von hydro- und ingenieurgeologischen Aufgaben Durchführung von Feld- und Laborversuchen
03/2001	DMT-Gesellschaft für Geotechnik und Umweltschutz mbH Mitarbeit bei der Erstellung einer bergschadenkundlichen Gefährdungsanalyse
10/2000-12/2002	Universität Leipzig, Institut für Geophysik und Geologie Ausbau und Pflege der paläobotanischen Sammlung

13 Publikationsliste

13.1 Artikel

1. Stelzer, N., Fischer, A., Kästner, M., Richnow, H.-H. (2006): Analyse des anaeroben *in situ* Benzolabbaus: Vergleich von In-situ-Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen. *Grundwasser* 11(4), S. 247-258.
2. Stelzer, N., Büning, C., Pfeifer, F., Tebbe, C. C., Nijenhuis, I., Kästner, M., Richnow, H.-H. (2006): *In situ* microcosms (BACTRAP) to evaluate natural attenuation potentials in contaminated aquifers. *Organic Geochemistry* 37(10), S. 1394-1410.
3. Fischer, A., Stelzer, N., Eisenmann, H., Richnow, H.-H. (2006): Nachweis des mikrobiellen Schadstoffabbaus in Grundwasserleitern: Innovative Isotopenverfahren erfassen Abbauprozesse direkt im kontaminierten Grundwasser. *Terra Tec* 15(1-2), S. 14-17.
4. Kästner, M., Fischer, A., Nijenhuis, I., Geyer, R., Stelzer, N., Bombach, P., Tebbe, C. C., Richnow, H.-H. (2006): Assessment of Microbial *In Situ* Activity in Contaminated Aquifers. *Engineering in Life Science* 6(3), S. 234-251.
5. Braeckeveldt, M., Rokadia, H., Imfeld, G., Stelzer, N., Paschke, H., Kuschk, K., Kästner, M., Richnow, H.-H., Weber, S. (2007): Assessment of *in situ* biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland. *Environmental Pollution* 148(2), S. 428-437.
6. Nijenhuis, I., Stelzer, N., Kästner, M., Richnow, H.-H. (2006): Sensitive detection of anaerobic monochlorobenzene degradation using stable isotope tracers. *Environmental Science and Technology* 41(11), S. 3836-3842.
7. Braeckeveldt, M., Rokadia, H., Mirschel, G., Weber, S., Imfeld, G., Stelzer, N., Kuschk, K., Kästner, M., Richnow, H.-H. (2007): Biodegradation of chlorobenzene in a constructed wetland treating contaminated groundwater. *Water Science & Technology* 56(3), S. 57-62.
8. Fischer, A., Theuerkorn, K., Stelzer, N., Gehre, M., Thullner, M., Richnow, H.-H. (2007): Applicability of Stable Isotope Fractionation Analysis for the Characterization of Benzene Biodegradation in a BTEX-contaminated Aquifer. *Environmental Science and Technology* 41(10), S. 3689-3696.
9. Stelzer, N., Imfeld, G., Thullner, M., Lehmann, J., Poser, A., Richnow, H.-H., Nijenhuis, I. (2008): Integrative approach to delineate Natural Attenuation of chlorinated benzenes in anoxic aquifers. *Environmental Science and Technology*, submitted.

13.2 Konferenzvorträge

1. Nijenhuis, I., Stelzer, N., Fischer, A., Kästner, M., Richnow, H.-H.: Isotope concepts to assess natural attenuation in contaminated aquifers. The General Assembly of the European Geosciences Union (EGU), 24.-29.04.2005, Vienna, Austria.
2. Stelzer, N., Geyer, R., Kästner, M., Nijenhuis, I., Weber, S., Büning, C., Pfeifer, F., Tebbe, C., Richnow, H.-H.: Erfahrungen aus dem Einsatz von ^{13}C -beladenen

- Aufwuchskörpern (BACTRAP®) bei BTEX-Schadensfällen. KORA-Workshop mit Posterpräsentation "Perspektiven molekularer und isotopischer Methoden zum Nachweis des natürlichen Schadstoffabbaus in Böden, 29.-30.09.2005, FAL, Braunschweig, Deutschland.
3. Stelzer, N., Weber, S., Nijenhuis, I., Kästner, M., Richnow, H.-H.: Monitoring of *in situ* biodegradation of groundwater contaminants using a test system (BACTRAP®) with ¹³C-labelled substrates. GASIR Jahrestagung 10.-12.10.2005 in Jena, Deutschland.
 4. Stelzer, N., Fischer, A., Nijenhuis, I., Gehre, M., Kästner, M., Richnow, H.-H.: Aktuelle Entwicklungen im Bereich MNA: Nachweis des mikrobiellen Schadstoffabbaus mit Isotopenmethoden. Jahrestreffen der ITVA Regionalgruppe Bayern, 16.02.2006 in München, Deutschland.
 5. Kästner, M., Fischer, A., Nijenhuis, I., Geyer, R., Stelzer, N., Bombach, P., Tebbe, C., Richnow, H.-H.: Assessment of microbial *in situ*-activity in contaminated aquifers. International Conference on Environmental Biotechnology (ISEB, ESEB, JSEB), 09.-13.07.2006, Leipzig, Germany.
 6. Stelzer, N., Richnow, H.-H.; Nijenhuis, I.: Lines of Evidence for anaerobic MCB Degradation in contaminated Groundwater based on Stable Isotope Tools. EGU European Geosciences Union, 15.-20.04.2007, Vienna, Austria.
 7. Kästner, M., Fischer, A., Nijenhuis, I., Geyer, R., Stelzer, N., Bombach, P., Tebbe, C.C., Richnow, H.-H.: *In situ* microbial activity assessment in contaminated aquifers. EGU European Geosciences Union, 15.-20.04.2007, Vienna, Austria.
 8. Stelzer, N., Nijenhuis, I., Lehman, J., Richnow, H.-H.: Isotopenmethoden zur Evaluierung des *Natural Attenuation* Potenzials an einem MCB kontaminierten Standort. 9. Symposium Natural Attenuation, 21.-22.11.2007 in Frankfurt, Deutschland.
 9. Stelzer, N., Weber, S., Nijenhuis, I., Richnow, H.-H.: Monitoring of *in situ* biodegradation in contaminated aquifers using *in situ* microcosms (BACTRAP®) with ¹³C-labelled substrates. AXIOM-VIBE-eTRAP-HIGRADE Workshop: "Electron transfer processes at biogeochemical gradients", 04.-07.03.2008, Leipzig, Germany.
 10. Stelzer, N., Nijenhuis, I., Weber, S., Fischer, A., Richnow, H.-H.: Einsatz stabiler Isotope zum qualitativen und quantitativen Nachweis des biologischen Schadstoffabbaus im kontaminierten Grundwasser. Tagung der FH-DGG, 21.05.-25.05.2008 in Göttingen, Deutschland.

13.3 Poster

1. Stelzer, N., Geyer, R., Fischer, A., Kästner, M., Weber, S., Richnow, H.-H.: Monitoring of *in-situ* biodegradation using "BACTRAPS" with ¹³C-labelled substrate. Poster presentation "1st German-American Workshop on Biogeochemical Gradients" 04.-06.05.2005 in Tübingen, Germany.
2. Stelzer, N., Kästner, M., Fischer, A., Richnow, H.-H.: Monitoring of *in-situ* biodegradation using a test system (BACTRAP) with ¹³C-labelled substrates. Poster presentation "2nd European Conference on Natural Attenuation, Soil and Groundwater Risk Management" 18.-20.05.2005 in Frankfurt, Germany.
3. Geyer, R.; Stelzer, N., Peacock, A., White D. C., Tebbe, C. C., Richnow, H.-H., Kästner, M.: Assessment of microbial *in situ* activity using microcosm amended with

Publikationsliste

- ¹³C-labelled substrates. Poster presentation "2nd European Conference on Natural Attenuation, Soil and Groundwater Risk Management" 18.-20.05.2005 in Frankfurt, Germany.
4. Richnow, H.-H., Stelzer, N., Nijenhuis, I., Kästner, M., Weber, S., Pfeifer, F., Tebbe, C. C.: In-Situ Microcosms to Assess Natural Attenuation in Contaminated Aquifers. Poster presentation International Symposia for Subsurface Microbiology (ISSM 2005) and Environmental Biogeochemistry (ISEB XVII), 14.-19.08.2005, Jackson Hole, Wyoming, USA.
 5. Stelzer, N., Geyer, R., Kästner, M., Fischer, A., Weber, S., Richnow, H.-H. "Monitoring of *in situ* biodegradation in contaminated aquifers using BACTRAPS with ¹³C-labelled substrates. Poster presentation International Summer School: "Biomonitoring, bioavailability and microbial transformation of pollutants in sediments and approaches to stimulate their biodegradation". 12-14.09.2005, Genoa, Italy.
 6. Stelzer, N., Kästner, M., Weber, S., Büning, C., Pfeifer, F., Richnow, H.-H. (2005): BACTRAP®: *In situ* Testsystem zum Nachweis des mikrobiellen Schadstoffabbaus in kontaminierten Grundwasserleitern. Posterpräsentation zum Workshop „Perspektiven molekularer und isotopischer Methoden zum Nachweis des natürlichen Schadstoffabbaus in Böden“, 29.-30.09.2005, FAL, Braunschweig, Deutschland.
 7. Fischer, A.; Vogt, C.; Herrmann, S.; Stelzer, N.; Theuerkorn, K.; Eisenmann, H.; Richnow, H. H.: Monitoring *in situ*-benzene biodegradation within an anoxic BTEX contaminated aquifer using stable isotope fractionation analysis (SIFA). VAAM-Jahrestagung, 19.-22.03.2006, Jena, Deutschland.
 8. Stelzer, N., Nijenhuis, I., Kästner, M., Richnow, H.-H.: *In situ* microcosms to evaluate natural attenuation potentials in contaminated aquifers. Poster presentation at the AXIOM-Virtual Institute Spring School: „Microbial Activity at Biogeochemical Gradients“. 03.-06.04.2006, Leipzig, Germany.
 9. Stelzer, N., Nijenhuis, I., Kästner, M., Richnow, H.-H.: *In situ* microcosms to evaluate natural attenuation potentials in contaminated aquifers. Poster presentation at the ISEB, ESEB, JSEB. 09.-13.07.2006, Leipzig, Germany.
 10. Nijenhuis, I., Stelzer, N., Nikolausz, M., Kästner, M., Richnow, H.-H.: Evidence for microbial Degradation of Monochlorobenzene under anaerobic conditions. Poster presentation at the 11th International Symposium on Microbial Ecology (ISME), 20.-25.08.2006 in Vienna, Austria.
 11. Fischer, A., Vogt, C., Herrmann, S., Stelzer, N., Theuerkorn, K., Herklotz, I., Thullner, M., Richnow, H.-H.: Characterization and assessment of BTEX biodegradation within aquifers using compound-specific stable isotope analysis (CSIA). Poster presentation at the HydroEco, 11.-14.09.2006 in Carlsbad, Czech Republic.
 12. Stelzer, N., Weber, S., Nijenhuis, I., Kuntze, K., Kästner, M., Richnow, H.-H.: Monitoring of microbial activity, metabolites and degradation pathways in contaminated aquifers using *in situ* microcosms with ¹³C-labelled substrates. Poster presentation at the 3. European Conference on Natural Attenuation and In-Situ Remediation, 19.-21.11.2007 in Frankfurt, Germany.

14 Anhang (siehe CD)

- A Analyse des anaeroben Benzolabbaus: Vergleich von *In situ* Mikrokosmen, Elektronenakzeptorbilanzen und Isotopenfraktionierungsprozessen (2006)
(Grundwasser 11(4), S. 247-258)
- B Nachweis des mikrobiellen Schadstoffabbaus in Grundwasserleitern (2006)
(Terra Tec 15(1-2), S. 14-17)
- C In situ microcosms to evaluate natural attenuation potentials in contaminated aquifers (2006)
(Organic Geochemistry 37(10), S. 1394-1410)
- D Assessment of Microbial In Situ Activity in Contaminated Aquifers (2006)
(Review in Engineering in Life Science 6(3), S: 234-251)
- E Sensitive Detection of Anaerobic Monochlorobenzene Degradation Using Stable Isotope Tracers (2007) und Supporting Information
(Environmental Science and Technology 41(11), S. 3836-3842)
- F Integrative approach to delineate Natural Attenuation of chlorinated benzenes in anoxic aquifers (2008) und Supporting Information
(Environmental Science and Technology submitted, 18.04.2008)
- G Assessment of in situ biodegradation of monochlorobenzene in contaminated groundwater treated in a constructed wetland (2007)
(Environmental Pollution 148(2), S. 428-437)
- H Biodegradation of chlorobenzene in a constructed wetland treating contaminated groundwater (2007)
(Water Science and Technology 56(3), S. 57-62)
- I Applicability of Stable Isotope Fractionation Analysis for the Characterization of Benzene Biodegradation in a BTEX-contaminated Aquifer (2007) und Supporting Information
(Environmental Science and Technology 41(10), S. 3689-3696)

