

This is the accepted manuscript version of the contribution published as:

Guber, A., **Blagodatskaya, E.**, Kravchenko, A. (2022):
Are enzymes transported in soils by water fluxes?
Soil Biol. Biochem. **168** , art. 108633

The publisher's version is available at:

<http://dx.doi.org/10.1016/j.soilbio.2022.108633>

1 Are enzymes transported in soil by water fluxes?

2 Andrey Guber^{a,b*}, Evgenia Blagodatskaya^c, Alexandra Kravchenko^{a,b},

3
4 *^a Department of Soil, Plant and Microbial Sciences, Michigan State University, East Lansing,*
5 *MI, USA*

6 *^b DOE Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, MI,*
7 *USA*

8 *^c Department of Soil Ecology, Helmholtz Centre for Environmental Research – UFZ, Halle*
9 *(Saale), Germany*

10
11 * Corresponding author: A.K. Guber. E-mail: akguber@msu.edu. 1066 Bogue st., East Lansing,
12 MI, 48824, USA

14 **Highlights**

- 15 • Hydrolytic enzymes are transported convectively, attached to soil colloids
- 16 • This transport is soil and enzyme specific
- 17 • Soil colloids released jointly with ions which alter ionic strength in effluents
- 18 • Enzyme activity demonstrated a bell-shape relationship with the ionic strength

20 **ABSTRACT**

21 Transport of extracellular hydrolytic enzymes in soils has always been a subject of doubt. The
22 considerations against its importance are that (i) enzymes benefit their producers the most when
23 they remain in close proximity; and (ii) enzymes are large molecules with low mobility due to
24 high affinity to fine soil particles and organic matter. However, soil mineral colloids (SMC), to
25 which extracellular enzymes also have an affinity and which are known to facilitate transport of
26 a broad variety of chemicals and microorganisms in soils, can serve as vehicles for enzyme
27 transport as well. Since current literature lacks information on enzyme transport in soils, our goal
28 was to determine whether enzymes are transported and, if so, whether they are transported in a
29 free- or in a colloid-associated form. We conducted column transport experiments with four
30 hydrolytic enzymes, namely, β -glucosidase, acid-phosphatase, cellobiohydrolase, and
31 xylosidase, in soils with contrasting textures. The eluents containing enzymes were applied on

32 top of soil columns, while enzyme activities, SMC, and electrical conductivity were measured in
33 the effluents from the columns. Our results provided evidence of joint enzyme transport with soil
34 colloids. The enzymes associated with the coarse SMC ($1 \mu\text{m} < \text{Ø}$) contributed 52 - 88% of the
35 total enzyme activity in the effluents. The remaining enzyme activity was attributed to the
36 enzymes associated with organic colloids, fine SMC ($\text{Ø} < 1 \mu\text{m}$) and free enzymes in solution.
37 This study suggested a dual effect of ionic strength in the soil suspension on enzyme activity and
38 their release from soils with soil colloids.

39

40 *Keywords:* Soil hydrolytic enzymes, colloid-facilitated transport, column experiments, ionic
41 strength.

42

43 *Abbreviations:* IS, ionic strength; EC, electrical conductivity; POM, particulate organic matter;
44 SMC, soil mineral colloids; CMC, coarse mineral colloids; FMC, fine mineral colloids; TN, total
45 nitrogen; TC, total carbon; SL, Sandy loam; SL-S, sandy loam soil from summit; SL-D, sandy
46 loam soil from depression.

47

48 INTRODUCTION

49 Most plant and microbial cell debris present within soil as polymeric molecules are quickly
50 transformed by extracellular enzymes to oligo- and monomers, which then become readily
51 available to microbial decomposers. Like most proteins, soil extracellular enzymes are capable to
52 diffuse away from their parent cell in free solutions due to Brownian motion (Burns et al., 2013).
53 Moreover, several studies in artificial solutions have reported a self-propelled diffusion of
54 enzymes, which enhanced their movement by 30-80% during substrate catalysis (Yu et al., 2009;
55 Muddana et al., 2010; Riedel et al., 2015; Jee et al., 2018; Günther et al., 2018). Still, there is no
56 consensus on the enzyme diffusivity. On one hand, diffusivity of free enzymes should increase as
57 substrate availability decreases, thus the enzyme producer can potentially access more distant
58 substrates (Allison et al., 2011). On the other hand, competition for products between enzyme
59 producers suggests relatively low enzyme diffusivity (Burns et al., 2013). The amount of reaction
60 products captured by a microbial cell per unit of enzyme produced declines with increasing
61 distance between the cell and the produced enzymes due to diffusion losses of the product to the
62 environment, cell competition for reaction products, and decreasing enzyme and substrate

63 concentrations. These product losses increase exponentially with the distance between the
64 enzymes and microbial cells, and less than 4% of the product can reach the microbial cells
65 located at a distance $> 200 \mu\text{m}$ from the enzymes (Guber et al, 2021). Therefore, low enzyme
66 diffusion, and thus lower diffusion losses of the product might represent a beneficial strategy for
67 microbial cells.

68 The enzyme's capacity to move within an unrestricting volume of liquid does not imply that
69 enzymes can similarly freely diffuse through soil pores. Among factors restricting such diffusion
70 are: (i) enzyme attachment to soil particles and soil colloids (Nannipieri et al., 2003); (ii)
71 relatively low water retention and fast draining of large pores resulting in reduction of pore
72 volume available for enzyme movement (Allison et al., 2011); (iii) hydraulic discontinuity of
73 water pathways in partly saturated pores and layers of extracellular polymeric substances,
74 created by microorganisms on surface of soil particles, restricting free enzyme motion (Or et al.,
75 2007). Moreover, most free enzymes survive only briefly in pore solutions due to fast
76 denaturation by physical and chemical factors or consumption by proteolytic microorganisms
77 (Sarkar and Burns, 1984; Burns, 1986). However, adsorption confers protection against
78 microbial degradation (Lähdesmäki & Piispanen, 1992; Kedi et al., 2013). Therefore, most
79 viable enzymes and microorganisms in soil are bound to organic surfaces (Ahmed and Oades,
80 1984; Christensen and Bech-Andersen, 1989; Jocteur Monrozier et al., 1991) and fine soil
81 particles (Singh and Singh, 1995; Schulten et al., 1993; Stemmer et al., 1998; Kandeler et al.,
82 1999a). Association of microbial cells and enzymes with soil particles and organic surfaces
83 precludes their free diffusive transport due to relatively large size of these particles and low pore
84 volume available for the diffusion in partly water-saturated soil. However, rapid water
85 infiltration after heavy rainfalls or overland irrigation causes both physical and chemical
86 perturbations in the soil and results in a release of colloids from soil matrix and their transport
87 via large pores (Ryan and Elimelech, 1996). These colloids are composed of organic and mineral
88 particles (including clay and partly silt soil fractions) with effective diameters $< 10 \mu\text{m}$
89 (McCarthy and Zachara, 1989; Sposito, 2016). The colloids within size range $0.01\text{-}10 \mu\text{m}$ are
90 regarded as the most stable (Buffle and Leppard, 1995) and are common carriers of soil
91 microorganisms, organic substances (Smith et al., 1985; McCarthy & Zachara, 1989; Natsch et
92 al., 1996), and environmental contaminants (Ryan & Elimelech, 1996; de Jonge et al., 2004).
93 Strong adsorption of enzymes on fine soil particles and their low extractability in free form

94 (Fornasier et al., 2011), suggests possibility of their transport by moving colloids. However,
95 despite extensive study of enzyme interaction with clay minerals and soil colloids during last 4
96 decades, surprisingly, enzyme transport in soils has never been reported. Therefore, the goal of
97 this study was to explore the possibility of enzyme transport in intact soils with contrasting
98 textures under water flow conditions that mimic those during heavy rainfalls.

99 The activity of four hydrolytic enzymes involved in C and P acquisition were studied in the
100 transport experiment: β -glucosidase, acid-phosphatase, cellobiohydrolase, and xylosidase. The
101 former two participate in the last step of decomposition, i.e., release of monomers (glucose and
102 phosphate) that are easily available for microorganisms. The latter two are involved in the early
103 stage of decomposition destroying long polymeric chains of cellulose and hemicelluloses.

104

105 2. MATERIALS AND METHODS

106 2.1. *Soil properties*

107 Soil for the column experiments was collected at three experimental sites with contrasting
108 soil texture located in Michigan, USA. Sandy soil of Riddles-Hillsdale series (fine-loamy, mixed,
109 active, mesic Typic Hapludalfs) was obtained from Michigan State University's (MSU) Sandhill
110 farm site, East Lansing, MI. We further refer to this soil as Sand. Two sandy loam soils of Capac
111 series (fine-loamy, mixed active, mesic Aquic Glossudalf) were obtained from summit and
112 depression topographical positions at MSU's Mason experimental farm, East Lansing, MI. We
113 refer to them as sandy loam soils SL-S and SL-D, for summit and depression respectively.

114 Loamy soil of Kalamazoo series (fine-loamy, mixed, active, mesic Typic Hapludalfs) was from
115 Cellulosic Biofuel Diversity Experiment at Kellogg Biological Station Long-Term Ecological
116 Research site, Hickory Corners, MI, referred to further as Loam soil. At each site, undisturbed
117 soil cores ($\varnothing=2.5$ cm, height=10 cm) were taken from 5-15 cm depth and stored at 4°C for 2-3
118 days prior to experiments. In addition, disturbed soil samples in amounts of approximately 300 g
119 were taken from the immediate vicinity of the intact cores for basic soil analyses. At the time of
120 sampling, Sand soil was under corn, SL-S and SL-D soils were under corn-soybean rotation, and
121 Loam soil was under long-term (>10 years) native prairie vegetation. We collected 3 cores from
122 each site and one additional core from Loam soil site, for a total of 13 cores. The number of soil
123 cores was selected arbitrarily solely for exploring purposes.

124 The following soil analyses were conducted using disturbed soil samples. Soil texture and
125 soil mineral colloids (SMC), operationally defined as mineral particles $\text{\O} < 10 \mu\text{m}$, were
126 measured using the pipette method (Gee & Or, 2002). Total nitrogen (TN) and total carbon (TC)
127 were measured using an elemental analyzer ECS 4010 CHNSO (Costech Analytical
128 Technologies Inc., Valencia, CA, USA). Soil particulate organic matter (POM) was measured by
129 wet sieving (Cambardella and Elliot, 1992). Soil pH and electrical conductivity (EC) were
130 measured using SevenCompact Duo s213 meter (Mettler-Toledo LLC, Columbus, OH USA) in
131 Soil-DI water suspensions at 1:1 solid/liquid ratio. General characteristics of the studied soils are
132 given in Table 1.

133 *2.2 Applied suspensions*

134 The compositions of applied suspensions, referred to as eluents, for the enzyme transport
135 experiment were designed to minimize artificial effects of solution chemistry on the transport
136 and transformation processes within the soil cores. This composition mimicked soil suspensions
137 generated in the field during heavy rainfalls by kinetic energy of rain drops or runoff water.
138 Therefore, the eluents were prepared individually for each soil by adding 1 g of fresh soil to 100
139 ml of DI water followed by 5 min low-energy sonication using Fisher Scientific FS20 Ultrasonic
140 Cleaner (Thermo Fisher Scientific Inc., Waltham, MA, USA). The sonication settings were
141 chosen to break up soil aggregates, while preserving SMC (Stemmer et al., 1998), soil organic
142 colloids, microorganisms and enzymes (Kandeler et al., 1999b). Soil suspensions were kept for
143 30 min to allow settling of sand particles, and the supernatant solutions separated from the
144 sediment were used as eluents. Thus, the prepared eluents contained dissolved chemicals, soil
145 particles, microorganisms, and enzymes native to the respective soils. The activities of β -
146 glucosidase, phosphatase, xylosidase and cellobiohydrolase were measured in the eluents before
147 and after precipitating the coarse mineral colloids (CMC) as described in Sections 2.3 and 2.4.
148 Soil mineral colloids were measured in the eluents as described in Section 2.2.

149 *2.3 Column experiment*

150 Soil column experiments were conducted to quantify possible activity of the four hydrolytic
151 enzymes in the suspensions passing through the intact soil cores, which we will refer to as
152 effluents. We used undisturbed soil cores to preserve distribution of enzymes and water flow
153 pathways unchanged in the soils. For Sand soil the entire soil cores were used as experimental

154 columns, that is, the column height was 10 cm. For finer textured SL-S, SL-D, and Loam soils,
 155 due to their low water infiltration rates, the intact cores were cut to 5 cm experimental columns.
 156 Longer Sand columns as compared with those for SL-S, SL-D, and Loam soils reduced the water
 157 flow velocity in the Sand columns and prevented mechanical detachment of colloids from soil.
 158 The eluents were applied by a pipette to the top of the columns in 1 ml increments to prevent
 159 water ponding on the soil surface (Fig. 1). A pressure head of -30 kPa, which is an equivalent to
 160 the field capacity, was maintained at the bottom of the columns during the experiment to keep
 161 steady-state flow through the columns. Coarse porous filters with particle retention $\text{Ø} > 40 \mu\text{m}$
 162 (Filter Paper Grade 417, VWR[®], Radnor, PA, USA) were installed at the bottom of the columns.
 163 The filters, permeable for SMC, microorganisms, and enzymes, prevented detachment of soil
 164 particles from the columns. The effluents were collected in 4 ml increments from the bottom of
 165 Loam, SL-S and SL-D columns, and in 8 ml increments from Sand columns. Collected effluents
 166 were analyzed for activity of the four enzymes, SMC contents, pH, and EC. The enzyme
 167 activities were measured in the effluents with all colloids and in the supernatant solutions which
 168 contained only the colloidal particles smaller than $1 \mu\text{m}$ (Fig. 1). To precipitate the coarse
 169 mineral colloids (CMC) size of large $1 \mu\text{m}$, the effluents were centrifuged for 5 min at 5000 rpm
 170 using a Heraeus Megafuge 16 centrifuge (Thermo Fisher Scientific Inc., Waltham, MA, USA).
 171 Diameter of precipitated CMC was calculated (Gee and Or, 2002) as:

$$172 \quad d = \frac{1}{60} \sqrt{\frac{18\eta \ln\left(\frac{r_2}{r_1}\right)}{(\rho_s - \rho_l)\omega^2 t}} \quad (1)$$

173 where r_1 to r_2 are distances from the axis of centrifuge rotor to the particle [cm], η is the fluid
 174 viscosity [$\text{g cm}^{-1} \text{sec}^{-1}$], $\rho_s \sim 2.6 \text{ g cm}^{-3}$ is the density of soil mineral particles, ρ_l is the liquid
 175 density [g cm^{-3}], ω is the centrifuge angular velocity [rpm], t is time for particle of diameter d to
 176 settle from r_1 to r_2 [sec]. It was confirmed in a preliminary study that selected centrifugation
 177 settings had a negligible effect on the activity of colloid-free almond β -glucosidase and wheat
 178 germ acid phosphatase (CALZYME Laboratories, Inc., San Luis Obispo, CA U.S.A.) dissolved
 179 in DI water. Precipitating mineral colloids smaller than $1 \mu\text{m}$ (e.g., 0.45 mm) and organic
 180 colloids ($\rho_s < 2.6 \text{ g cm}^{-3}$) requires increasing of angular velocity of centrifugation proportionally
 181 to ω or increasing the duration of centrifugation as t^2 . Such changes in the centrifugation settings
 182 results in partial precipitation of bio colloids and free enzymes in the effluent suspension in

183 amounts that are generally unknown. Therefore, most organic colloids, bio colloids enzymes and
184 mineral colloids size of smaller 1 μm remained in supernatants, while CMC were precipitated by
185 centrifugation.

186 For comparison purposes the results of the column experiments were expressed in a
187 normalized form. Specifically, volumes of eluent that passed through the columns were
188 expressed in total pore volumes, where the total pore volume in each soil column was calculated
189 from its bulk density, particle density [$\sim 2.6 \text{ g cm}^{-3}$], and column's volume. The SMC contents,
190 enzyme activities and EC values in the effluents from the columns were normalized by the
191 corresponding values in the eluents. To evaluate the effect of ionic strength in the effluents on
192 the enzyme activity, the measured EC values were transformed using the Marion-Babcock
193 equation (Sposito, 2016):

$$194 \quad \log_{10}(I) = -1.841 + 1.009 \log_{10}(\text{EC}) \quad \text{for } I \leq 0.2 \text{ M} \quad (2)$$

195 where I is the ionic strength in the solution [M], and EC is the electrical conductivity in the
196 solution [dS m^{-1}].

197 2.4. Enzyme assay

198 The activity of β -glucosidase, acid-phosphatase, xylosidase and cellobiohydrolase were
199 measured in the soil, eluents, and effluents collected in the column experiments before and after
200 precipitating CMC as described in Section 2.3 (Fig. 1). For the activity measurements we used a
201 microplate fluorometric assay technique described in Saiya-Cork et al. (2002) and Deng et al.
202 (2011) with minor modifications of the substrate concentrations. Specifically, we used substrates
203 based on 4-methylumbelliferone (MUF) fluorescent reagent (i.e. MUF- β -D-glucopyranoside,
204 MUF-Phosphate, MUF- β -D-xylopyranoside, and MUF- β -D-cellobioside) in quantities of 40
205 nmol well^{-1} per 50 $\mu\text{l well}^{-1}$ soil suspensions with 5.0 $\mu\text{mol well}^{-1}$ of sodium MES buffer
206 ($\text{C}_6\text{H}_{13}\text{NO}_4\text{SNa}_{0.5}$, pH 6.1 at 25°C) solution. The calibration was performed using 0, 100, 200
207 and 400 pmol well^{-1} MUF solutions with the same aliquots of soil suspensions. The fluorescence
208 intensity in the plates was measured using a Multilabel Plate Reader Victor³ (PerkinElmer Inc.,
209 Waltham, MA, USA) every 15 min for 2 hours. The enzyme activities were calculated from
210 linear parts of the intensity time series with correction for the substrate autohydrolysis (Deng et
211 al., 2011).

212 The enzyme activity associated with CMC in the effluents v_c^{ef} was calculated as:

213
$$v_c^{ef} = v_t^{ef} - v_s^{ef} \quad (3)$$

214 were v_t^{ef} is the total enzyme activity in the effluent [$\text{pmol min}^{-1} \text{ ml}^{-1}$], and v_s^{ef} is the enzyme
215 activity in the supernatant from centrifuged effluent [$\text{pmol min}^{-1} \text{ ml}^{-1}$].

216 Since the mass of SMC differed in the effluents from the four soils and changed over time, to
217 evaluate the effect of ionic strength on the enzyme activity associated with SMC, the activity was
218 normalized by the dry mass of colloids in the effluent:

219
$$v_c^S = v_c^{ef} / S_c \quad (4)$$

220 where v_c^S is the activity of enzymes associated with CMC per mass of dry colloids [$\text{nmol min}^{-1} \text{ g}$
221 dry colloids^{-1}], and S_c is the dry mass of CMC in the effluent [g L^{-1}]. Note, that the dry mass of
222 colloids in supernatants from the effluents was within the accuracy range of analytical balance
223 (0.1 mg). Therefore, the dry mass of SMC measured in the effluents was attributed to CMC.

224 2.5. Data processing and statistical analysis.

225 Differences among the soils in terms of the studied properties were conducted via analysis of
226 variance, with soils as the only studied factor, using *lm* function of R (version 4.1.2). To compare
227 the enzyme activities within the soils we fitted the data with a statistical model that consisted of
228 (i) the fixed effects of the soils and the enzymes and their interaction, and (ii) a random effect of
229 the replicated samples, nested within the soils, also used as an error term for testing the soils'
230 effect. Model fitting was conducted using *lmer* function from R's *lme4* package. When the
231 effects of the studied factors were found to be statistically significant ($p < 0.05$) we performed t-
232 tests to conduct all pairwise comparisons among the means using *emmeans* package.

233 We used correlation analysis to explore possible relationships between enzyme activity
234 associated with CMC and colloid contents in the effluents from the soil columns using Analysis
235 tool of SigmaPlot software (Systat Software Inc., San Jose, California, USA). Pearson
236 correlation coefficients were reported as statistically significant and $p < 0.05$ and marginally
237 significant at $p < 0.1$.

238 3. RESULTS

239 3.1. Properties of the soils and applied suspensions.

240 The soils of the four locations differed in their properties. The bulk density was the lowest in
241 the sandy soil (Sand) due to its loose structure. The highest sand content was expectedly

242 observed in Sand, lowest in Loam, and intermediate in the sandy loam SL-S and SL-D soils
243 (Table 1). An inverse trend was observed for SMCs, which contents were the highest in Loam,
244 lowest in Sand, and intermediate, though close to Loam, in SL-S and SL-D soils (Table 1). The
245 POM content was rather low in all soils (0.11 % – 0.31%) but tended to be the highest in Sand.
246 TN and TC increased in the soils in the order of decreasing sand content. The pH values ranged
247 from 5.4 to 6.0. The EC values were 2 to 3 times higher in SL-S and SL-D as compared to Sand
248 and Loam, indicating overall higher contents of dissolvable ions in sandy loam soils.

249 The SMC content in the eluents increased in the order Sand < SL-D < SL-S < Loam. The pH
250 values were slightly higher than in the soil suspensions and ranged from 5.9 to 6.7, while the EC
251 values were much lower (6-9 $\mu\text{S cm}^{-1}$) in the eluents as compared to those in the soils, as an
252 expected result of a 100-fold dilution of the soil suspensions (Table 1).

253 The differences in the soil properties were mirrored by enzyme activities, which for β -
254 glucosidase and phosphatase were the highest in Loam and SL-S, and the lowest in Sand (Table
255 1). For xylosidase and cellobiohydrolase the highest activities were observed in SL-S, and the
256 lowest in Sand. The enzyme activity in the eluents followed the same general trend as that in the
257 soils, with much smaller differences between β -glucosidase and phosphatase activities as
258 compared with those in the soils. Among the four enzymes, the activities in most soil samples
259 and eluents increased in the order xylosidase < cellobiohydrolase < phosphatase < β -glucosidase
260 and were up to one order of magnitude higher for phosphatase and β -glucosidase than for
261 xylosidase and cellobiohydrolase (Table 2). Precipitation of CMC by centrifugation considerably
262 reduced enzyme activity in the eluents. The enzyme activity associated with CMC calculated
263 using Eq.(3) ranged from 33% to 100% in the eluents in the four soils. The lowest activities of
264 CMC-associated enzymes were observed in Sand (Table 2), while the highest activities were
265 observed in Loam, where the enzyme activities in the supernatants after eluent centrifugation
266 were below the detection limit.

267

268 *3.2. Column experiment*

269 The relative concentrations of SMC in the effluents were the highest in the first portions of
270 effluent collected from the columns and decreased with the relative volumes of the solution
271 passed through the columns (i.e. pore volumes) (Fig. 2 and Fig. S1). The recovered SMC
272 differed in the four soils and ranged from 7% of that applied in Loam to 49% in Sand, with

273 intermediate values for SL-S and SL-D (Table 2). The relative concentrations of SMC were the
274 highest for Sand and the lowest for Loam columns (Fig. 2ab). Noticeably, in the Sand columns
275 the relative concentrations of SMC from the first portions of the effluent were >1 (Fig. 2a),
276 indicating that the effluents contained more SMC than the applied eluents. For Loam (Fig. 2b),
277 SL-S (Fig. S1a), and SL-D (Fig. S1b) columns, the SMC contents in the effluents were always
278 smaller than in the eluents.

279 The EC dynamics in the effluent resembled that of SMC, though unlike SMC, the highest
280 relative EC were observed in SL-S and SL-D columns (Fig. S1). The relative EC values in the
281 effluents were $\gg 1$ in all samples of all studied soils, indicating much higher concentrations of
282 soluble chemicals in the effluents as compared to those in the applied eluents.

283 As was the case with SMC and EC, the relative activity of four enzymes in the effluents
284 decreased with the amount of eluent passing the columns (Fig. 3). Among the four enzymes, the
285 absolute values of activities in the effluents were the lowest for xylosidase and cellobiohydrolase
286 and the highest for phosphatase (Table 3). Enzymatic activity in the first portions of the effluents
287 1.5 to 8 times exceeded those in the eluents (Fig. 3 and Fig. S2), with exception of β -glucosidase
288 in SL-S and SL-D columns (Fig. S2a) and cellobiohydrolase in the Loam column (Fig. 3b).
289 Among the four enzymes, the relative activity in the first portions of effluents was the highest for
290 phosphatase (Fig. 3b and Fig. S2b), the smallest for β -glucosidase (Fig. 3a and Fig S2a), and
291 intermediate for xylosidase (Fig. 3c and Fig. S2c) except the Loam soil, where xylosidase and
292 cellobiohydrolase were not detected in the effluents. Among the four soils, the relative activities
293 in the effluent were the highest for Sand, the lowest for Loam (Fig. 3), and intermediate for SL-S
294 and SL-D columns (Fig. S2), which was consistent with the dynamics of SMC outflow from the
295 columns (Fig. 3 and Fig. S1). For example, β -glucosidase activities in the effluent corresponding
296 to 1 pore volume in Sand was >2 times higher than that in Loam, while for phosphatase it was >4
297 times higher than in Loam (Table 3).

298 For β -glucosidase and phosphatase the activity associated with CMC constituted a substantial
299 portion of the total enzyme activity (Fig. 4, Table 3). For example, for β -glucosidase in the
300 effluent corresponding to the first pore volume, the activity associated with CMC constituted
301 66% and 52% of the total activity for Loam and Sand, respectively (Table 3). For phosphatase, it
302 was even higher and constituted 79% and 88% for Loam and Sand, respectively. In most tested
303 enzymes, their activity associated with CMC was positively correlated with concentration of

304 SMC in the effluents (Fig. 5 and Fig. S3). The correlation coefficients between v_c^{ef} and
305 concentration of SMC in the effluents were high at $p < 0.1$ and $p < 0.05$ for all enzymes except
306 xylosidase and cellobiohydrolase in the Loam column where the activities were below the
307 detection level (Fig. 5), and except phosphatase, xylosidase and cellobiohydrolase in SL-D
308 columns (Fig. S3).

309 The relationships between the activity of enzymes associated with SMC v_c^{ef} and ionic
310 strength in the effluents, as derived from EC using Eq.(2), were rather scattered. Still, a sharp
311 increase of β -glucosidase and phosphatase activity associated versus not associated with CMC
312 can be seen in the Sand and Loam columns with increasing ionic strength from 0.1 to 2.3 mM
313 (Fig. 6). The slope of linear regression between the enzyme activity and IS ranged from 4.5 to 43
314 $\text{pmol min}^{-1} \text{g}^{-1} \text{mM}^{-1}$ and was steeper for Sand as compared with that in the Loam columns. For
315 the SL-S and SL-D columns the activity of all enzymes was somewhat decreasing within the
316 ionic strength range 1.4 - 21.4 mM. This trend was less expressed but still visible for xylosidase
317 (Fig. S4 a,c) and cellobiohydrolase (Fig. S4 b,d) in the SL-S and SL-D columns.

318 The pH values in the effluents moderately increased with pore volumes during the
319 experiment from 6.0 to 6.6, 6.3 to 7.1 and 5.2 to 6.3 in the Sand, Loam, and both SL-S and SL-D
320 columns, respectively. However, v_c^{ef} did not correlate with pH in the effluents.

321

322 4. DISCUSSION

323 The results of column experiments demonstrated possibility for soil hydrolytic enzymes to
324 be transported by water fluxes during heavy rainfall events in the soils with different soil texture.

325 4.1. Associations between enzymes and colloids in the eluents

326 Precipitation of CMC in the eluents reduced considerably activity of the four enzymes
327 suggesting their association with colloids (Table 2). CMC are comprised of clay and silt
328 particles, which hydrolytic enzymes are mostly associated with (Feller et al., 1994; Turner et al.,
329 2002; Sinsabaugh et al., 2008; Kandeler et al., 1999a; Kedi et al., 2013). Association of β -
330 glucosidase with soil mineral colloids in paddy soil has been reported by Yan et al. (2010 a,b).
331 These authors found approximately 50% higher β -glucosidase association with fine ($< 0.2 \mu\text{m}$)
332 than with the coarse ($0.2 - 2.0 \mu\text{m}$) SMC. On the contrary, 86% to 100% of β -glucosidase
333 activity was associated with CMC in our study. The discrepancy likely results from Yan et al

334 (2010 a,b) using highly diluted (1:100 solid to liquid ratio) soil suspensions, where all added
335 enzymes freely interacted with all resuspended soil particles. Yet, in intact soils, as used in our
336 study, enzyme-colloid interaction occurs primarily in hydrologically active pores occasionally
337 and not completely saturated by liquids.

338 It should be noted that enzyme-CMC associations in eluents were enzyme and soil specific.
339 While activity of all enzymes was completely associated with CMC in Loam soil, in the coarser
340 textured soils the percent of CMC associated activity was lower. Of the four studied enzymes, β -
341 glucosidase seemed to be the most associated with CMC, while activity of xylosidase and
342 cellobiohydrolase tended to be the least associated, especially in Sand (Table 2). While it is not
343 clear what is the cause for the differences among the enzymes, the results suggest high
344 possibility of joint transport of enzymes with CMC, especially in fine-textured soils, though
345 transport of not associated enzymes cannot be excluded.

346 4.2. Colloid and enzyme transport through the soil columns

347 The transport experiments demonstrated high mobility and recovery of SMC in sandy and
348 sandy loam soils and low mobility in the Loam (Fig. 2), likely caused by straining of colloids
349 within small pores of the finer textured soil. This result is consistent with other studies (Bradford
350 et al., 2002, 2003), where straining of colloids was shown to increase with increasing colloid size
351 and with decreasing soil grain size (Bradford et al., 2003).

352 Soil colloids appeared to be involved in enzyme transport, the result that was consistently
353 observed in all cases when the enzymes were present in the soil in appreciable amounts (Table
354 3). High significant correlation between the activity of these enzymes associated with CMC
355 (v_c^{ef}) and SMC contents in the effluent (Fig. 5, Fig. S3) imply that v_c^{ef} was proportional to the
356 mass of transported colloids for each portion of the solution passed the columns. Closeness of the
357 intercepts of the linear regressions to zero for all enzymes in Sand and Loam columns (Fig. 5),
358 and considerable (from 2 to 10-fold) decreases in the enzyme activity after precipitation of CMC
359 in the effluents (Table 3 and Fig. 5) suggested that large fraction of the enzymes was transported
360 convectively attached to SMC.

361 High correlation coefficients between enzyme activity and contents of SMC in the
362 effluents do not imply that all detected enzyme activity was associated solely with SMC. Non-
363 zero activities of all enzymes were detected in the supernatants after centrifugation of effluents

364 from SL-D columns (Fig. S3). The centrifugation removed mostly CMC ($\varnothing > 1 \mu\text{m}$) from the
365 effluents. Fine mineral colloids, organic colloids, microbial cells and macromolecules, with
366 density much smaller than that of soil mineral particles ($\rho_s < 2.6 \text{ g cm}^{-3}$), were only partly
367 precipitated by centrifugation, but could also act as colloidal carriers (Buffle and Leppard, 1995).
368 The number of fine colloid particles $< 0.2 \mu\text{m}$ (not necessarily mass) can constitute up to 70% of
369 the total number of mobile colloids during a heavy rainfall (Lehmann et al., 2021). Moreover, the
370 activity of β -glucosidase associated with fine ($< 0.2 \mu\text{m}$) SMC can be higher than that associated
371 with coarse ($0.2 - 2.0 \mu\text{m}$) SMC (Yan et al., 2010 a,b). Therefore, enzymes remaining in the
372 effluents after centrifugation were more likely associated with not-precipitated FMC, viable
373 microbial cells (Nannipieri et al., 2012), and/or organic colloids rather than being free floating
374 enzymes, whose lifetime in soil solutions is rather short (Burns, 1982; Ladd, 1985; Nannipieri,
375 1994, Nannipieri et al., 2002). Possible enzyme association with soil colloids suggest that in real-
376 world systems enzyme transport depends on the source and composition of moving colloids.
377 Specifically for agricultural environments, colloid-facilitated transport of enzymes can be
378 expected after irrigation by reclaimed wastewater or heavy rainfalls followed manure
379 application.

380 Colloidal transport is known to affect soil hydraulic properties, e.g., conductivity of soil
381 pores and their sizes (Miller and Baharuddin, 1986; McDowell-Boyer et al., 1986). Precipitation
382 and straining of colloids in soil pores during colloidal transport results in progressive straining of
383 colloids and in the associated decrease of pore volume available for transport. Therefore, it is
384 reasonable to suggest that, with time, the ratio between large and small colloids in the effluents
385 shifts toward small colloids. Since the proportion of organics and the surface area of colloids
386 increases with decreasing sizes of colloidal material (Buffle and Leppard, 1995), higher
387 association of enzymes with fine colloids is expected. This fact can be illustrated by β -
388 glucosidase (Fig. 4a) and xylosidase (Fig. 4c) activities in SL-S columns, where the ratio
389 between activities of enzymes associated (filled bars) and not associated (open bars) with CMC
390 decreased with the number of pore volumes.

391 The relationship between SMC content in effluents and activity of CMC-associated
392 enzymes was enzyme specific (Fig. 5 and Fig. S3) possibly due to mineralogical composition of
393 transported colloids and differences in enzyme-mineral interactions. The mineral composition of
394 Ap horizon in fine, mixed, mesic, Typic Hapludalfs, used in our study, varies greatly among soil

395 fractions. According to Sparks et al. (1979), quartz and mica dominate (>50%) in > 5 μm and < 2
396 μm mineral fractions of Typic Hapludalfs. The 2-5 μm soil fraction is composed of quartz, mica
397 and kaolinite in approximately equal amounts. Kaolinite contents vary from 7 to 28% in the soil
398 fractions with approximately two-fold greater contents in < 5 μm than in > 5 μm soil fractions.
399 Feldspar is present only in silt fractions, while vermiculite only in clay fractions, in the amounts
400 of less than 10%. Different enzyme affinity to the substrates in presence of soil minerals has
401 been reported in multiple studies (Ross, 1983; Makboul and Ottow, 1979; Haska, 1975; Pflug
402 (1982); Gianfreda and Bollag, 1994). Therefore, it is reasonable to suggest that SMC-enzyme
403 associations and enzyme activities in the effluents were affected by changing sizes and
404 mineralogy of transporting colloids. For example, the share of quartz particles in the total SMC
405 mass passing the columns likely decreased with time, while the share of mica and kaolinite
406 particles increased. Due to differences in soil texture, it is expected that the ratios between
407 contents of different minerals in the effluents was soil specific. Therefore, the differences in the
408 enzyme activities observed in Fig. 5 and Fig. S3 can likely be attributed to different association
409 of enzymes and their activity on SMC, which mineralogy changed in the effluents with respect to
410 the textural, mineralogical, and hydraulic properties of these soils. We realize that it is infeasible
411 to conduct particle and mineralogical analysis in the effluents, but it is worth to note that the
412 relationship between the enzyme activity and quantities of transported colloids is more
413 complicated than any adsorption isotherm used to model interactions between chemicals,
414 microorganisms, and soil particles.

415 *4.3. Whether ionic strength in solution affect enzyme transport?*

416 Soil's ionic strength affected recovery of SMC applied with eluent in the fine-textured soils
417 (Loam, SL-S and SL-D). The recovery was greater in the SL-S and SL-D columns (Table 2),
418 where soil EC and, respectively, IS were much higher than those in the Loam soils (Table 1).
419 Increasing IS causes coagulation, while decreasing IS causes disaggregation and mobilization of
420 soil colloids (Ryan and Elimelech, 1996). Rainfall water has typically much lower IS as
421 compared with that in soil solutions (McCarthy and Zachara, 1989). Therefore, decrease of soil
422 IS, due to dilution of pore solution by rainwater, results in a release of colloids from the soil
423 matrix at early stages of rainwater infiltration into the soil (Flury et al., 2002; Grolimund and
424 Borkovec, 1999). With time the difference between IS in soil pores and the applied solution
425 decreases slowing down soil disaggregation. However, soil, disaggregated by low IS, becomes

426 less conductive and filters out new infiltrated colloids via straining or physical-chemical
427 collection by attractive surfaces of the immobile soil matrix (McCarthy and Zachara, 1989)
428 Visually, this process manifested itself in this study via a noticeable reduction of the infiltration
429 rate in the fine textured soils after 1-1.5 pore volumes of the effluent were collected. The high
430 recovery of colloids from the Sand columns (49.1%) was less affected by soil disaggregation due
431 to much lower clay to sand ration as compared with the other soils (Table 1). Therefore, the
432 effect of IS on colloidal outflow was less pronounced in the Sandy, and more pronounced in
433 Loam and Sandy loam columns.

434 The IS also affects the activity of transported enzymes. In batch experiments with pure clay
435 minerals and in solutions with controlled pH and EC the relationships between enzyme activities
436 and IS could be well described quantitatively (e.g., Quiquampoix et al., 1993; Leprince &
437 Quiquampoix, 1996). In real soil and transport conditions this relationship is rather scattered
438 (Fig. 6) and likely altered by: (i) possible differences in mineralogy of soil colloids of the four
439 soils and effluents from the columns (Jaber et al., 2018; Nannipieri et al., 1996); (ii) different
440 affinity of enzymes to different minerals and colloids (Makboul and Ottow, 1979; Ross, 1983;
441 Sarkar et al., 1989); (iii) and different and likely changing during the transport experiment ionic
442 and colloidal composition of the effluents. Yet, we attempt to derive a relationship between the
443 activity of adsorbed enzymes and IS for out column studies using dependencies between
444 phosphatase catalytic activity and pH measured for montmorillonite at different IS levels
445 reported by Leprince & Quiquampoix (1996, Fig. 2, central column). The original data were
446 reorganized to depict relationships between the rate of the catalytic reaction (V) and IS at pH
447 levels within 5.0 - 7.0 interval (Fig. 7, symbols). The reorganized relationships were fitted using
448 an empirical equation:

$$449 \quad V = \frac{a}{IS} \exp[-b(\ln(IS) - c)^2] \quad (5)$$

450 where a , b , and c are the fitting parameters. The regression wizard of SigmaPlot software (Systat
451 Software Inc., San Jose, California, USA) was used to fit Eq.(5) to the data shown in Fig. 7. The
452 similarity in shapes between the regression curves and experimental data was assessed using the
453 two-sample Kolmogorov-Smirnov test in the Real Statistics Resource Pack software (Release
454 7.6, Copyright (2013 – 2021), Charles Zaiontz, www.real-statistics.com).

455 Despite a small number of experimental points (6 on each curve), Eq.(5) adequately ($p <$
456 0.05) reproduced shapes of the $V(IS)$ curves (Fig. 7). Therefore, Eq.(5) was applied to reconstruct

457 the shape of experimental curves describing relationships between activities of enzymes (i.e., β -
458 glucosidase and phosphatase associated (Fig. 6 a,b) and not associated (Fig. 6 c,d) with CMC)
459 and IS in effluents from the four soils. Based on the Kolmogorov-Smirnov test the fitted curves
460 reproduced adequately ($p < 0.05$) the shapes of β -glucosidase and phosphatase activity curves in
461 the Sand and Loam columns for enzymes not associated with CMC (Fig. 6 c,d). The shapes of
462 fitted curves differed significantly ($p < 0.05$) from those experimental for enzymes associated with
463 the CMC. Fitting Eq.(5) to xylosidase and cellobiohydrolase activity was not successful due to
464 scattered data on these curves (Fig. S4).

465 Several mechanisms can potentially explain a bell-shape relationship between enzyme
466 activity and IS. The first one is a competition between enzymes and cations in the effluents for
467 adsorption sites on mineral and organo-mineral colloids. Large IS values imply higher contents
468 of anions and cations in the effluents, and stronger competition with enzymes for the adsorption
469 sites on SMC. The increase in IS also alters repulsive electrostatic interactions and weakens
470 enzyme association with colloids. The competition mechanism explains the decrease of enzyme
471 activity but does not explain its increase within the IS range 0.2-1.3 mM. It also does not explain
472 the decrease in activity of the enzymes not associated with CMC (Fig. 6 c,d) which were likely
473 partly associated with unprecipitated colloids and partly were in a free form in the effluents.

474 The second mechanism, affecting the relationship between the enzyme activity and IS, is
475 based on enzyme activity association with pH, which commonly has a bell-curve shape with the
476 maximum activity at an optimal pH level (Leprince and Quiquampoix, 1996; Turner, 2010; Kedi
477 et al., 2013). Importantly, the optimal pH level of enzyme activity narrows and shifts toward the
478 alkaline pH when enzymes are adsorbed on mineral surfaces of soil particles (McLaren and
479 Estermann, 1957; Aliev et al., 1976; Leprince and Quiquampoix, 1996). Furthermore, the
480 enzymes activity peaks shift towards more acid pH values with increasing IS in the suspensions
481 (Goldstein et al., 1964). Given almost one unit difference in the pH between the surface of the
482 colloids and the soil solution and changing chemical composition of the effluents, the pH optima
483 for particular enzyme species, as well as the activity of the colloid associated enzymes, can vary
484 strongly and deviate from that in chemically clean laboratory solution. Therefore, less adequate
485 reproduction of our experimental curves (Fig. 6) by Eq.(5) than those derived from Leprince and
486 Quiquampoix (1996) (Fig. 7) can be attributed to variations in mineral composition of soil
487 colloids and chemical composition of effluents.

488 4.4. *Implications of colloid-facilitated transport.*

489 The colloid-facilitated transport of microbial cells and enzymes is likely part of microbial
490 survival strategy. While microorganisms in soil are primarily attached to the solid surfaces and
491 form associations there (e.g., colonies, films, or flocks), their activity is mainly associated with
492 soil solution. Most organic materials entering the soil (plant residue, dead roots, manure) are
493 insoluble or only partly soluble in water, and barely mobile. Therefore, soil microorganisms must
494 either populate most of available pore space or be highly mobile to access new organic inputs.
495 However, microbial cells and free enzymes are unlikely capable to travel far in soil pores in
496 searching for new energy sources due to: (i) their relatively large size and slow diffusion rates
497 (i.e., for 1-10 $\mu\text{m s}^{-1}$ for enzymes and 7–8 $\mu\text{m s}^{-1}$ microbial cells according to Young et al., 1980;
498 Dechesne et al., 2010; and Zhang and Hess, 2019); (ii) small pore volumes available for
499 diffusion because of pore discontinuity and low thickness of water menisci in partly saturated
500 soils; (iii) overall short presence of free enzymes in soil solutions due to denaturation and
501 proteolysis, or interaction with clay and organic surfaces (Burns, 1982; Ladd, 1985; Sarkar et al.,
502 1989; Nannipieri, 1994, 2002). Moreover, most enzymes are irreversibly attached to soil solids,
503 as evident through low extractability of enzymes from bulk soil (Vepsäläinen, 2001; Štursová
504 and Baldrian 2011). Therefore, only a small fraction of soil enzymes can potentially freely
505 diffuse in soil pores and reach newly added organic materials.

506 Mobilization of colloids and their convective transport through soil macropores during fast
507 water flow events, e.g., irrigation or rainfall, as well as in partly saturated soils, are important for
508 soil microorganisms in exploring new sources of energy. The benefits of colloid-facilitated
509 transport for soil microbes are obvious: (i) much shorter times and longer travel distances as
510 compared with the restricted diffusion; (ii) better protection from protozoa on colloid surfaces
511 (Sarkar et al., 1980; Nannipieri et al., 1982); (iii) energy savings for production of new enzymes,
512 since enzymes can be transported by the colloids; (iv) influx of new partly degraded organic
513 materials from soil surface and their joint transport with microbial cells. Such transport explains
514 appearance of new hotspots of microbial activity commonly observed in soil after heavy rainfalls
515 much better than the diffusion theory does. Soil hotspots and hot moments are defined based on
516 time and rates of microbial activity exceeding the average rates in bulk soil (Kuzyakov and
517 Blagodatskaya, 2015). Input of labile substrates to the hotspots triggers microbial activity and
518 thus drives the hot moments. We suggest that, in addition to the substrate-triggered hot spot

519 activation, the colloid-facilitated transport enables enzymes and microbial cells to move quickly
520 and in relatively large quantities to or with the labile substrates, thus forming new transport-
521 triggered hotspots during high precipitation and preferential flow events.

522 The results of this study and communication with anonymous reviewers have risen research
523 questions that require further in-depth exploration. Among them there are: (i) how to separate the
524 release of colloids and enzymes from soil matrix and their transport through pores; (ii) how sizes
525 and mineralogy of colloids present in soil affect enzymes association with them; (iii) how sizes
526 and mineralogy of colloids are changing in the effluents, and to what extent these changes affect
527 enzyme association and transport with colloids; (iv) what drives the differences between
528 enzymes in their associations with mineral and biological soil colloids and subsequent transport;
529 (v) are enzymes transported with their producers; (vi) how soil structure affects enzyme locations
530 within the soil matrix and their transport with colloids; (vii) which factors are the dominant
531 drivers of enzyme transport in soils?

532

533 **CONCLUSIONS**

534 This study revealed a possibility for hydrolytic enzymes (i.e. β -glucosidase, acid-
535 phosphatase, xylosidase and cellobiohydrolase) to be transported through soil pores by water
536 fluxes. Strong association of hydrolytic enzymes with fine soil particles and mobility of soil
537 colloids results in their joint convective transport. This transport is affected by ionic strength in
538 pore solution via dissociation and release of soil colloids from soil and alteration of enzyme
539 activity in the transported suspensions. The former effect can be attributed to the shift of optimal
540 pH of enzyme activity near the surface of soil colloids. It remains to be seen how soil texture and
541 structure, colloid size and composition, enzyme properties and location in soil pores contribute to
542 their release and transport in the field conditions.

543

544 **Declaration of competing interest**

545 The authors declare that they have no known competing financial interests or personal
546 relationships that could have appeared to influence the work reported in this paper.

547

548 **Acknowledgement**

549 Support for this research was provided in part by the NSF DEB Program (Award #
550 1904267); by the NSF LTER Program (DEB 1027253) at the Kellogg Biological Station; by the
551 Great Lakes Bioenergy Research Center, U.S. Department of Energy, Office of Science, Office
552 of Biological and Environmental Research under Award Number DE-SC0018409; by
553 Foundational Program “Microbial Communities in Soil” under Award Number 2022-67019-
554 36104 from the USDA National Institute of Food and Agriculture; and by the German Research
555 Foundation (DFG; project number 403664478) in context of Priority Programme (2089)
556 “Rhizosphere spatiotemporal organization”. We would also like to acknowledge four anonymous
557 reviewers for constructive comments and suggestions during preparation of this manuscript.
558

559 REFERENCES

- 560 Ahmed, M., Oades, J.M., 1984. Distribution of organic matter and adenosine triphosphate after
561 fractionation of soils by physical procedures. *Soil Biology and Biochemistry* 16, 465-470.
- 562 Aliev, R.A., Guzev, V.S., Zvyagintsev, D.G., 1976. Influence of adsorbents on the optimum pH
563 of catalase. *Moscow University Soil Science Bulletin* 31(2): 32-34.
- 564 Allison, S.D., Weintraub, M.N., Gartner, T.B., Waldrop, M.P., 2011. Evolutionary- economic
565 principles as regulators of soil enzyme production and ecosystem function. In: Shukla, G.,
566 Varma, A. (Eds.), *Soil Enzymology*. Springer-Verlag, Berlin, Germany, pp. 229-243.
- 567 Bradford, S.A., Simunek, J., Bettahar, M., Van Genuchten, M.T., Yates, S.R., 2003. Modeling
568 colloid attachment, straining, and exclusion in saturated porous media. *Environmental*
569 *Science and Technology* 37(10), 2242–2250. <https://doi.org/10.1021/es025899u>
- 570 Bradford, S.A., Yates, S.R., Bettahar, M., & Simunek, J., 2002. Physical factors affecting the
571 transport and fate of colloids in saturated porous media. *Water Resources Research* 38(12),
572 63-1–63-12. <https://doi.org/10.1029/2002wr001340>
- 573 Buffle, J., and Leppard, G.G., 1995. Characterization of aquatic colloids and macromolecules. 1.
574 Structure and behavior of colloidal material. *Environmental Science & Technology* 29, 2169–
575 2175.
- 576 Burns, R.G., 1982. Enzyme activity in soil: Location and a possible role in microbial ecology.
577 *Soil Biology and Biochemistry* 14, 423–427.
- 578 Burns, R.G., 1986. Interaction of Enzymes with Soil Mineral and Organic Colloids. *Interactions*
579 *of Soil Minerals with Natural Organics and Microbes*, Volume 17. Madison, WI United

580 States: Soil Science Society of America, pp.429-451.
581 <https://doi.org/10.2136/sssaspecpub17.c11>

582 Burns, R.G., DeForest, J.L., Marxsen, J., Sinsabaugh, R.L., Stromberger, M.E., Wallenstein,
583 M.D., Weintraub, M.N., Zoppini, A., 2013. Soil enzymes in a changing environment: Current
584 knowledge and future directions. *Soil Biology and Biochemistry* 58, 216–234.
585 <https://doi.org/10.1016/j.soilbio.2012.11.009>

586 Cambardella, C.A., Elliot, E.T., 1992. Particulate soil organic matter changes across a grassland
587 cultivation sequence. *Soil Science Society of America Journal* 56:777-83.

588 Christensen, B.T., Bech-Andersen, S., 1989. Influence of straw disposal on distribution of amino
589 acids in soil particle-size fractions. *Soil Biology and Biochemistry* 21, 35-40.

590 de Jonge, L.W., Kjaergaard, C., Moldrup, P., 2004. Colloids and Colloid-Facilitated Transport of
591 Contaminants in Soils: An Introduction. *Vadose Zone Journal* 3(2), 321–325.
592 <https://doi.org/10.2136/vzj2004.0321>

593 Dechesne, A., Wang, G., Gulez, G., Or, D., Smets, B.F., 2010. Hydration-controlled bacterial
594 motility and dispersal on surfaces. *Proceedings of the National Academy of Sciences of the*
595 *United States of America* 107, 14369–14372.

596 Deng, S., Kang, H., Freeman, C., 2011. Microplate fluorimetric assay of soil enzymes. In Dick,
597 R.P. (Ed.), *Methods of soil enzymology*. Soil Science Society of America, Madison, WI. pp.
598 311–318.

599 Feller, C., Frossard, E. et Brossard, M., 1994. Activité phosphatasique de quelques sols tropicaux
600 à argile 1:1. Répartition dans les fractions granulométriques. *Canadian Journal of Soil Science*
601 74, 121-129.

602 Fornasier, F., Dudal, Y., Quiquampoix, H., 2011. Enzyme extraction from soil. In Dick R.P.
603 (Ed.), *Methods of soil enzymology*. Soil Science Society of America, Madison, WI, pp. 371–
604 384.

605 Flury, M., Flühler, H., 1994. Susceptibility of soils to preferential flow of water: a field study.
606 *Water Resources Research* 30, 1945-1954.

607 Flury, M., Mathison, J.B., Harsh, J.B., 2002. In situ mobilization of colloids and transport of
608 cesium in Hanford sediments. *Environmental Science & Technology* 36, 5335–5341.

609 Gee, G.W., Or, D., 2002. Particle Size Analysis. In: Dane, J.H. and Topp, G.C. (Eds.). Methods
610 of Soil Analysis, Part 4, Physical Methods, Soils Science Society of America, Book Series
611 No. 5, Madison, pp.255-293.

612 Gianfreda, L., Bollag, J-M., 1994. Effect of Soils on the Behavior of Immobilized Enzymes. Soil
613 Science Society America Journal 58, 1672-1681.

614 Goldstein, L., Levin, Y., Katchalski, E., 1964. A water-insoluble polyanionic derivative of
615 trypsin. 11. Effect of the polyelectrolyte carrier on the kinetic behavior of the bound trypsin.
616 Biochemistry 3, 1913-1919.

617 Grolimund, D., Borkovec, M., 1999. Long-term release kinetics of colloidal particles from
618 natural porous media. Environmental Science & Technology 33, 4054–4060.

619 Günther, J.P., Börsch, M., Fischer, P. 2018. Diffusion measurements of swimming enzymes with
620 fluorescence correlation spectroscopy. Accounts of Chemical Research 51, 1911–1920.

621 Haska, G. 1975. Influence of clay minerals on sorption of bacteriolytic enzymes. Microbial
622 Ecology 1, 234-245.

623 Jaber, M., Lambert, J. F., Balme, S., 2018. Protein adsorption on clay minerals. Developments in
624 Clay Science 9, 255-288. <https://doi.org/10.1016/B978-0-08-102432-4.00008-1>

625 Jee, A-Y.; Dutta, S.; Cho, Y-K.; Tlustý, T.; Granick, S., 2017. Enzyme leaps fuel
626 antichemotaxis. Proceedings of the National Academy of Sciences. 115. 201717844.
627 10.1073/pnas.1717844115.

628 Jocteur Monrozier, L., Ladd, J.N., Fitzpatrick, R.W., Foster, R.C., Raupach, M., 1991.
629 Components and microbial biomass content of size fractions in soil of contrasting
630 aggregation. Geoderma 49, 37-62.

631 Kandeler, E., Palli, S., Stemmer, M., Gerzabek, M.H., 1999a. Tillage changes microbial biomass
632 and enzyme activities in particle-size fractions of a Haplic Chernozem. Soil Biology and
633 Biochemistry 31(9), 1253–1264. [https://doi.org/10.1016/S0038-0717\(99\)00041-3](https://doi.org/10.1016/S0038-0717(99)00041-3)

634 Kandeler, E., Stemmer, M., Klimanek, E.M., 1999b. Response of soil microbial biomass, urease
635 and xylanase within particle size fractions to long-term soil management. Soil Biology and
636 Biochemistry 31, 261-273.

637 Kedi, B., Abadie, J., Sei, J., Quiquampoix, H., Staunton, S., 2013. Diversity of adsorption
638 affinity and catalytic activity of fungal phosphatases adsorbed on some tropical soils. Soil
639 Biology and Biochemistry 56, 13–20. <https://doi.org/10.1016/j.soilbio.2012.02.006>

640 Kuzyakov, Y., Blagodatskaya, E. 2015. Microbial hotspots and hot moments in soil: Concept &
641 review. *Soil Biology and Biochemistry*, 83, 184–199.
642 <https://doi.org/10.1016/j.soilbio.2015.01.025>

643 Ladd, J.N. 1985. Soil enzymes. In: Vaughan, D., Malcom, R.E., (Eds). *Soil Organic Matter and*
644 *Biological Activity*. Dordrecht, Netherlands: Martinus Nijhoff, pp 175–221.

645 Lähdesmäki, P., Piispanen, R., 1992. Soil enzymology - role of protective colloid systems in the
646 preservation of exoenzyme activities in soil. *Soil Biology and Biochemistry* 24, 1173–1177.

647 Lehmann, K., Lehmann, R., Totsche, K.U., 2021. Event-driven dynamics of the total mobile
648 inventory in undisturbed soil account for significant fluxes of particulate organic carbon.
649 *Science of the Total Environment* 756, 143774.
650 <https://doi.org/10.1016/j.scitotenv.2020.143774> 143774.

651 Leprince, F., Quiquampoix, H. 1996. Extracellular enzyme activity in soil: Effect of pH and
652 ionic strength on the interaction with montmorillonite of two acid phosphatases secreted by
653 the ectomycorrhizal fungus *Hebeloma cylindrosporum*. *European Journal of Soil Science*,
654 47(4), 511–522. <https://doi.org/10.1111/j.1365-2389.1996.tb01851.x>

655 Makboul, H. E., Ottow, J.C.C., 1979. Alkaline phosphatase activity and the Michaelis constant in
656 the presence of different clay minerals. *Soil Science* 128, 129-135.

657 McCarthy, J.F., Zachara, J.M., 1989. Subsurface transport of contaminants. *Environmental*
658 *Science and Technology* 23(5), 496-502.

659 McDowell-Boyer, L.M., Hunt, J.R., Sitar, N., 1986. Particle transport through porous media,
660 *Water Resources Research*, 22, 1901-1921.

661 McLaren, A.D, Estermann, E.F., 1957. Influence of pH on the activity of chymotrypsin at a
662 solid-liquid interface. *Archives of Biochemistry and Biophysics* 68, 157-160.

663 Miller, W.P., and Baharuddin, M.K., 1986. Relationship of soil dispersibility to infiltration and
664 erosion of southeastern soils. *Soil Science* 142, 235–240.

665 Muddana, H.S., Sengupta, S., Mallouk, T.E., Sen, A., Butler, P.J., 2010. Substrate catalysis
666 enhances single-enzyme diffusion. *Journal of the American Chemical Society* 132, 2110-
667 2111.

668 Nannipieri, P., 1994. The potential use of soil enzymes as indicators of productivity,
669 sustainability and pollution. In: Pankhurst, C.E., Doube, B.M., Gupta, V.V.S.R., Grace, P.R.,

670 (Eds.). *Soil Biota: Management in Sustainable Farming Systems*, Adelaide CSIRO, pp. 238-
671 244.

672 Nannipieri, P., Ascher, J., Ceccherini, M., Landi, L., Pietramellara, G., Renella, G., 2003.
673 Microbial diversity and soil functions. *European Journal of Soil Science* 54, 655-670.

674 Nannipieri, P., Giagnoni, L., Renella, G., Puglisi, E., Ceccanti, B., Masciandaro, G., Fornasier,
675 F., Moscatelli, M.C., Marinari, S., 2012. Soil enzymology: classical and molecular
676 approaches. *Biology and Fertility of Soils* 48, 743–762.

677 Nannipieri, P., Kandeler, E., Ruggiero, P., 2002. Enzyme activities and microbiological and
678 biochemical processes in soil. In: Burns, R.G., Dick, R. (Eds.), *Enzymes in the Environment.*
679 *Activity, ecology and applications*. Marcel Dekker, New York, pp. 1-33.

680 Nannipieri, P., Sequi, P., Fusi, P., 1996. Humus and enzyme activity. In: Alessandro, P. (Ed.),
681 *Humic Substances in Terrestrial Ecosystems*. Elsevier Science B.V., Amsterdam, pp. 293-
682 328.

683 Natsch, A., Keel, C., Troxler, J., Zala, M., Albertini, N., Défago, G., 1996. Importance of
684 preferential flow and soil management in vertical transport of biocontrol strains of
685 *Pseudomonas fluorescens* in structured field soil. *Applied and Environmental Microbiology*
686 62, 33-40.

687 Or, D., Smets, B.F., Wraith, J.M., Dechesne, A., Friedman, S.P., 2007. Physical constraints
688 affecting bacterial and activity in unsaturated porous media - a review. *Advances in Water*
689 *Resources* 30, 1505–1527.

690 Pflug, W. 1982. Effect of clay minerals on the activity of polysaccharide cleaving soil enzymes.
691 *Journal of Plant Nutrition and Soil Science* 145, 493-502.

692 Quiquampoix, H., Staunton, S., Baron, M.H., Ratcliffe, R.G., 1993. Interpretation of the pH
693 dependence of protein adsorption on clay mineral surfaces and its relevance to the
694 understanding of extracellular enzyme activity in soil. *Colloids and Surfaces A:*
695 *Physicochemical and Engineering Aspects*, 75(C), 85–93. [https://doi.org/10.1016/0927-](https://doi.org/10.1016/0927-7757(93)80419-F)
696 [7757\(93\)80419-F](https://doi.org/10.1016/0927-7757(93)80419-F)

697 Riedel, C., Gabizon, R., Wilson, C.A.M., Hamadani, K., Tsekouras, K., Marqusee, S., Presse, S.,
698 Bustamante, C., 2015. The heat released during catalytic turnover enhances the diffusion of an
699 enzyme. *Nature* 517, 227-230.

700 Ross, D. J., 1983. Invertase and amylase activities as influenced by clay minerals, soil-clay
701 fractions and topsoils under grassland. *Soil Biology and Biochemistry* 15, 287-293.

702 Ryan, J. N., Elimelech, M., 1996. Colloid mobilization and transport in groundwater. *Colloids*
703 *and Surfaces A: Physicochemical and Engineering Aspects* 107(95), 1-56.
704 [https://doi.org/10.1016/0927-7757\(95\)03384-X](https://doi.org/10.1016/0927-7757(95)03384-X)

705 Saiya-Cork, K.R., Sinsabaugh, R.L., Zak, D.R., 2002. The effects of long term nitrogen
706 deposition on extracellular enzyme activity in an *Acer saccharum* forest soil. *Soil Biology and*
707 *Biochemistry* 34, 1309-1315. doi:10.1016/s0038-0717(02)00074-3.

708 Sarkar, J.M., Batistic, L., Mayaudon, J., 1980. Les hydrolases du sol et leur association avec les
709 hydrates de carbone. *Soil Biology and Biochemistry* 12, 325-328.

710 Sarkar, J.M., Burns, R.G., 1984. Synthesis and properties of B-D-glucosidase-phenolic
711 copolymers as analogues of soil humicenzyme complexes. *Soil Biology and Biochemistry* 16,
712 619-625

713 Sarkar, J.M., Leonowicz, A., Bollag, J.M., 1989. Immobilization of enzymes on clays and soils.
714 *Soil Biology and Biochemistry* 21(2), 223–230. [https://doi.org/10.1016/0038-0717\(89\)90098-](https://doi.org/10.1016/0038-0717(89)90098-9)
715 9

716 Schulten, H.-R., Leinweber P., Sorge C., 1993. Composition of organic matter in particle-size
717 fractions of an agricultural soil. *Journal of Soil Science* 44, 611-691.

718 Singh, S., Singh, J.S., 1995. Microbial biomass associated with water-stable aggregates in forest,
719 savanna and cropland soils of a seasonally dry tropical region, India. *Soil Biology and*
720 *Biochemistry* 27, 1027-1033.

721 Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw, C.,
722 Contosta, A.R., Cusack, D., Frey, S., Gallo, M.E., Gartner, T.B., Hobbie, S.E., Holland, K.,
723 Keeler, B.L., Powers, J.S., Stursova, M., Takacs-Vesbach, C., Waldrop, M.P., Wallenstein,
724 M.D., Zak, D.R., Zeglin, L.H., 2008. Stoichiometry of soil enzyme activity at global scale.
725 *Ecology Letters* 11, 1252-1264.

726 Smith, M.S., Thomas, G.W., White, R.E., Ritonga, D., 1985. Transport of *Escherichia coli*
727 through intact and disturbed soil columns. *Journal of Environmental Quality* 14, 87-91.

728 Sposito, G., 2016. *The Chemistry of Soils*, 3rd Edition, Oxford University Press, New York, NY

729 Stemmer, M., Gerzabek, M.H., Kandeler, E., 1998. Organic matter and enzyme activity in
730 particle-size fractions of soils obtained after low-energy sonication. *Soil Biology and*
731 *Biochemistry* 30(1), 9-17. [https://doi.org/10.1016/S0038-0717\(97\)00093-X](https://doi.org/10.1016/S0038-0717(97)00093-X)

732 Štursová, M., Baldrian, P., 2011. Effects of soil properties and management on the activity of
733 soil organic matter transforming enzymes and the quantification of soil-bound and free
734 activity. *Plant Soil* 338, 99–110. doi:10.1007/s11104-010-0296-3

735 Turner, B.L., Hopkins, D.W., Haygarth, P.M., Ostle, N., 2002. β -Glucosidase activity in pasture
736 soils. *Applied Soil Ecology* 20, 157-162.

737 Turner, B.L., 2010. Variation in pH optima of hydrolytic enzyme activities in tropical rain forest
738 soils. *Applied and Environmental Microbiology* 76, 6485-6493.

739 Vepsäläinen, M. 2001. Poor enzyme recovery by extraction from soils. *Soil Biology and*
740 *Biochemistry* 33,1131-1135.

741 Yan, J., Pan, G., Ding, C., & Quan, G., 2010a. Kinetic and thermodynamic parameters of β -
742 glucosidase immobilized on various colloidal particles from a paddy soil. *Colloids and*
743 *Surfaces B: Biointerfaces* 79(1), 298–303. <https://doi.org/10.1016/j.colsurfb.2010.04.015>

744 Yan, J., Pan, G., Li, L., Quan, G., Ding, C., & Luo, A., 2010b. Adsorption, immobilization, and
745 activity of β -glucosidase on different soil colloids. *Journal of Colloid and Interface Science*,
746 348(2), 565–570. <https://doi.org/10.1016/j.jcis.2010.04.044>

747 Young, M.E., Carroad, P.A., Bell, R.L., 1980. Estimation of diffusion coefficients of proteins.
748 *Biotechnology and Bioengineering* 22, 947–955.

749 Yu, H., Jo, K., Kounovsky, K.L., de Pablo, J. J., Schwartz, D.C., 2009. Molecular propulsion:
750 chemical sensing and chemotaxis of DNA driven by RNA polymerase. *Journal of the*
751 *American Chemical Society* 131, 5722-5723.

752 Zhang, Y., Hess, H., 2019. Enhanced Diffusion of Catalytically Active Enzymes. *ACS Central*
753 *Science*, 5(6), 939–948. <https://doi.org/10.1021/acscentsci.9b00228>.

754

755 Table 1. Selected properties of the soils collected from the four studied experimental sites.

Soil texture	Sand	Loam	Sandy loam (SL-S)	Sandy loam (SL-D)
Sand %	87±2* a **	38±2 b	58±8 c	56±2 c
Silt %	8±3 a	48±1 c	29±7 b	31±3 b
Clay %	5±2 a	14±1 b	13±1 b	13±1 b
SMC [‡] %	7±2 a	29±2 c	24±1 b	26±1 b
POM %	0.31±0.07 a	0.15±0.09 ab	0.15±0.10 ab	0.11±0.01 b
BD, g cm ⁻³	1.41±0.09 a	1.62±0.08 b	1.73±0.05 bc	1.80±0.05 c
pH	5.7±0.2	6.0±0.2	5.4±0.3	5.4±0.2
EC, μS cm ⁻¹	75±19 a	86±24 a	148±28 b	247±37 c
TN %	0.03±0.01 a	0.14±0.03 c	0.11±0.01 bc	0.10±0.01 b
TC %	0.49±0.11 a	1.53±0.34 c	1.13±0.01 cb	0.98±0.01 ab
Enzyme activity in soil, pmol min ⁻¹ g ⁻¹				
<i>β</i> -glucosidase	2.23±0.42 aA	6.53±1.01 cA	6.90±1.05 cA	4.94±0.18 bA
Phosphatase	1.09±0.26 aB	3.41±0.31 cB	2.77±0.16 bB	2.46±0.06 bB
Xylosidase	0.20±0.04 aC	0.30±0.08 aC	0.82±0.12 bC	0.66±0.12 bC
Cellobiohydrolase	0.23±0.03 aC	0.46±0.29 aC	0.89±0.04 bC	0.55±0.13 aC

756 * Data presented as mean ± one standard deviation.

757 ** Means within the same row followed by the same low case letter are not significantly different
 758 from each other (p<0.05); means of enzyme activity within the same column followed by the
 759 same upper-case letter are not significantly different from each other (p<0.05).

760 [‡] Particle size < 50 μm

761 SMC denotes the soil mineral colloids, POM denotes the particulate organic matter, BD denotes
 762 the soil bulk density, EC denotes the electrical conductivity, TN denoted the total nitrogen, and
 763 TC denotes total carbon.

764

765

766 Table 2. Selected properties of the applied suspensions generated from the soils from the four
 767 studied experimental sites

Soil texture	Sand	Loam	Sandy loam (SL-S)	Sandy loam (SL-D)
SMC, g l ⁻¹	0.68±0.16* a **	2.89±0.15 c	2.42±0.09 b	2.28±0.11 b
pH	5.9	6.7	6.2	6.4
EC, µS cm ⁻¹	6	7	8	9
Enzyme activity in the applied suspensions, pmol min ⁻¹ ml ⁻¹				
<i>β</i> -glucosidase	6.03±1.13 aA	9.66±1.22 bA	14.95±2.28 cA	14.06±0.52 cA
Phosphatase	3.13±0.75 aB	10.63±1.23 bA	10.26±0.60 bB	12.61±0.30 cB
Xylosidase	1.09±0.24 bC	0.28±0.08 aB	1.81±0.26 cD	1.60±0.30 cC
Cellobiohydrolase	0.69±0.09 aC	0.48±0.30 aB	3.43±0.16 cC	1.77±0.41 bC
Average enzyme activity associated with coarse mineral colloids (CMC) in the applied suspensions, %				
<i>β</i> -glucosidase	85.7 aA	100.0 aA	94.1 aA	97.3 aA
Phosphatase	97.6 bA	100.0 bA	57.4 aB	99.5 bA
Xylosidase	33.1 aB	100.0 bA	65.4 abB	57.3 abB
Cellobiohydrolase	32.9 bB	100.0 aA	99.5 aA	78.0 aAB
SMC recovered in the column experiment				
SMC recovered, % of applied	49.1±11.8 c	7.3±2.5 a	31.3±3.3 b	40.5±15.0 bc

768

769 * Data presented as mean ± one standard deviation.

770 ** Means within the same row followed by the same letter are not significantly different from
 771 each other (p<0.05); means of enzyme activity within the same column followed by the same
 772 upper-case letter are not significantly different from each other (p<0.05).

773

774


775 Table 3. Estimated enzyme activities in the effluent from the column experiments in Sand and
 776 Loam soils obtained for the eluent amount that replaced 1 pore volume in the columns. Shown
 777 are averages from all columns for the total enzyme activity in the effluent and the proportion of
 778 the total activity from the enzymes associated with CMC.

Enzymes	Soil	Total enzyme activity in the effluent, pmol min ⁻¹ ml ⁻¹	Activity of the enzymes associated with CMC, % of total
<i>β</i> -glucosidase	Loam	1.4	66
	Sand	3.0	52
Phosphatase	Loam	1.8	79
	Sand	7.6	88
Xylosidase	Loam	0.0	-
	Sand	1.0	50
Cellobiohydrolase	Loam	0.0	-
	Sand	0.9	30

779

780

787




788

789 Figure 2. Concentrations of soil mineral colloids (SMC) and electrical conductivity (EC) in the
790 effluents relative to those in the applied solutions for Sand (a) and Loam (b) soil columns for
791 eluent volumes that passed the columns relative to the pore volumes. Data are combined from 3
792 columns from each soil. Horizontal line (a) marks the relative SMC in the effluent corresponding
793 to that in the eluent. Different scales are used in (a) and (b) for better data visibility.


794

795




797
 798 Figure 3. Activity of the four enzymes in the effluents from Sand and Loam soil columns relative
 799 to those in the applied solutions for eluent volumes that passed the columns relative to the soil
 800 pore volumes. Zero values indicate enzyme activities below the detection limit. Data are
 801 combined from 3 columns of each soil. Horizontal lines mark the relative activity in the effluent
 802 corresponding to that in the eluent.

803



804
 805 Figure 4. Activity of enzymes associated (filled bars) and not associated (open bars) with CMC
 806 in the effluents from Sand, Loam, SL-S and SL-D soil columns for the four enzymes and
 807 different eluent volumes passing the columns. Sum of open and filled bars represents the total
 808 enzyme activity in the effluents.


809
 810
 811
 812



814

815 Figure 5. The relationships between activity of the four enzymes associated with CMC and SMC
 816 contents in the effluents from Sand and Loam soil columns. Zero values indicate concentration
 817 of colloids and enzyme activities below the detection limit. Lines denote linear regressions
 818 between enzyme activity and SMC contents. Data are combined from 3 columns for each soil.
 819 The numbers show the Pearson correlation coefficients at $p < 0.1$ (*) and $p < 0.05$ (**),
 820 respectively.

821



822

823 Figure 6. β -glucosidase and phosphatase activities associated and not associated with CMC as a
 824 function of ionic strength in the effluents from four soils. The activities are expressed per 1 g of
 825 SMC dry mass (a,b) and per 1 ml of effluent (c,d). Black lines show Eq.(5) fitted to the
 826 experimental data.

827


828

829

830

831

832




833

834 Figure 7. Rate of the catalytic reaction (V) of phosphatase adsorbed on montmorillonite as
835 function of ionic strength in solutions with different pH values (symbols) reconstructed from
836 Leprince and Quiquampoix (1996) (Fig. 2, Central column). Lines show lognormal distribution
837 fit to the experimental data.

838


839

840 **Supplemental materials**




841

842 Figure S1. Concentrations of soil mineral colloids (SMC) and electrical conductivity (EC) in the
843 effluents relative to those in the applied solutions for SL-S (a) and SL-D (b) soil columns for
844 eluent volumes that passed the columns relative to the pore volumes. Data are combined from all
845 columns of each soil.




846

847 Figure S2. Activity of the four enzymes in the effluents from SL-1 and SL-2 soil columns
 848 relative to those in the applied solutions for eluent volumes that passed the columns relative to
 849 the pore volumes. Data are combined from all columns of each soil. Horizontal lines mark the
 850 relative activity in the effluent corresponding to that in the eluent.



851
 852 Figure S3. The relationships between activity of the four enzymes associated with CMC and
 853 contents of SMC in the effluents from SL-S and SL-D soil columns. Zero values indicate
 854 concentration of colloids and enzyme activities below the detection limit. Lines denote linear
 855 regressions of enzyme activity on SMC contents. Data are combined from all columns of each
 856 soil. The numbers show the Pearson correlation coefficients at $p < 0.1$ (*) and $p < 0.05$ (**),
 857 respectively.



858
 859 Figure S4. Xylosidase and cellobiohydrolase activities of enzymes associated and not associated
 860 with CMC at different ionic strength in effluents from four soils. The activities are expressed per
 861 1 g of SMC dry mass (a,b) and per 1 ml of effluent (c,d).

862

863