

This is the final draft of the contribution published as:

Tartally, A., Thomas, J.A., **Anton, C.**, Balletto, E., Barbero, F., Bonelli, S., Bräu, M., Casacci, L.P., Csősz, S., Czekes, Z., Dolek, M., Dziekańska, I., Elmes, G., Fürst, M.A., **Glinka, U.**, Hochberg, M.E., Höttinger, H., Hula, V., Maes, D., Munguira, M.L., **Musche, M.**, Stadel Nielsen, P., Nowicki, P., Oliveira, P.S., Peregovits, L., **Ritter, S.**, Schlick-Steiner, B.C., **Settele, J.**, Sielezniew, M., Simcox, D.J., Stankiewicz, A.M., Steiner, F.M., Švitra, G., Ugelvig, L.V., Van Dyck, H., Varga, Z., Witek, M., Woyciechowski, M., Wynhoff, I., Nash, D.R. (2019):

Patterns of host use by brood parasitic *Maculinea* butterflies across Europe

Philos. Trans. R. Soc. B-Biol. Sci. **374** , art. 20180202

The publisher's version is available at:

<http://dx.doi.org/10.1098/rstb.2018.0202>

Host ant use and specificity of ant-brood-parasitic *Maculinea* butterflies across Europe

András Tartally^{1,2}(0000-0002-3284-8239), Jeremy A. Thomas³(0000-0002-5913-4041), Christian Anton⁴, Emilio Balletto⁵(0000-0003-1168-2791), Francesca Barbero⁵(0000-0003-2667-0435), Simona Bonelli⁵(0000-0001-5185-8136), Markus Bräu⁶, Luca Pietro Casacci^{5,7}(0000-0003-3083-7357), Sándor Csősz⁸, Zolt Czekeš⁹(0000-0002-5615-5883), Matthias Dolek¹⁰(0000-0003-3778-9482), Izabela Dziekańska¹¹, Graham Elmes¹², Matthias A. Fürst², Uta Glinka⁴, Michael E. Hochberg¹³, Helmut Höttinger¹⁴, Vladimir Hula¹⁵(0000-0001-7023-7087), Dirk Maes¹⁶(0000-0002-7947-3788), Miguel L. Munguira¹⁷(0000-0002-1554-4380), Martin Musche⁴, Per Stadel Nielsen¹⁸, Piotr Nowicki¹⁹(0000-0002-9797-886X), Paula S. Oliveira²⁰(0000-0003-0063-089X), Pavel Pech²¹, László Peregovits²², Sylvia Ritter¹⁹, Birgit C. Schlick-Steiner²³, Josef Settele^{4,24}(0000-0002-8624-4983), Marcin Sielezniew¹¹, David J. Simcox^{3,12}, Anna M. Stankiewicz-Fiedurek⁷, Florian M. Steiner²³(0000-0003-2414-4650), Giedrius Švitra²⁵, Line V. Ugelvig²(0000-0003-1832-8883), Marianna Vályi Nagy²², Hans Van Dyck²⁶, Zoltán Varga¹, Magdalena Witek⁷(0000-0002-6172-8981), Michal Woyciechowski¹⁹, Irma Wynhoff²⁷(0000-0002-3003-897X), David R. Nash²(0000-0002-0462-6794)*

¹University of Debrecen, Department of Evolutionary Zoology and Human Biology, H-4032, Debrecen, Egyetem tér 1 Hungary

²Centre for Social Evolution, Department of Biology, University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen, Denmark

³Department of Zoology, University of Oxford, Oxford, OX1 3PS, U.K.

⁴UFZ-Helmholtz Centre for Environmental Research, Department of Community Ecology, Theodor-Lieser-Strasse 4, 06120 Halle, Germany

⁵Department of Life Sciences and Systems Biology, University of Turin, Turin, 10123, Italy

⁶Bavarian Academy for Nature Conservation and Landscape Management, Seethalerstr. 6, Laufen/Salzach, 83410, Germany

⁷Laboratory of Social and Myrmecophilous Insects, Museum and Institute of Zoology, Polish Academy of Sciences, Wilcza 64, 00-679 Warszawa, Poland

⁸MTA-ELTE-MTM Ecology Research Group, Pázmány Péter sétány 1/C, Budapest, H1117, Hungary

⁹Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Clinicilor St 5–7, 400006 Cluj-Napoca, Romania

¹⁰Büro Geyer und Dolek, Alpenblick 12, 82237 Wörthsee, Germany

¹¹Institute of Biology, University of Białystok, Ciołkowskiego 1J, 15-245 Białystok, Poland

¹²Centre for Ecology & Hydrology Wallingford, Maclean Building, Benson Lane, Wallingford, OX10 8BB, UK

¹³Institut des Sciences de l'Evolution – CNRS UMR 5554, Université de Montpellier – CC 065, 34095 Montpellier Cedex 05, France Cedex 05, France

¹⁴Institute of Zoology, Department of Integrative Biology and Biodiversity Research, University of Natural Resources and Life Sciences, Gregor-Mendel-Straße 33, 1180 Vienna, Austria

¹⁵Department of Zoology, Hydrobiology, Fishery and Apiculture, Faculty of AgriScience, Mendel University Brno, Zemedelska 1, Brno, CZ-61300, Czech Republic

¹⁶Research Institute for Nature and Forest (INBO), Herman Teirlinckgebouw, Havenlaan 88 bus 73, B-1000 Brussels, Belgium

¹⁷Universidad Autónoma de Madrid, Facultad de Ciencias, Departamento de Biología (Zoología), Cantoblanco, ES-28049 Madrid, Spain

¹⁸Gevninge Bygade 10, 4000 Roskilde, Denmark

¹⁹Institute of Environmental Sciences, Jagiellonian University, Gronostajowa 7, 30-387 Kraków, Poland

- ²⁰Center for the Research and Technology of Agro-Environmental and Biological Sciences (CITAB) / University of Trás-os-Montes and Alto Douro, Department of Forest Science and Landscape, 5000-911 Vila Real, Portugal
- ²¹University of Hradec Králové, Faculty of Science, Hradec Králové, Czech Republic
- ²²Department of Zoology, Hungarian Natural History Museum, H-1088 Budapest Baross u. 13., Hungary
- ²³Molecular Ecology Group, Department of Ecology, University of Innsbruck, Technikerstrasse 25, 6020 Innsbruck, Austria
- ²⁴German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany
- ²⁵Lithuanian Entomological Society, Akademijos 2, LT-08412 Vilnius, Lithuania
- ²⁶Behavioural Ecology & Conservation Group, Biodiversity Research Centre, Earth and Life Institute, Université catholique de Louvain (UCL), Louvain-la-Neuve, Belgium
- ²⁷Dutch Butterfly Conservation & Butterfly Conservation Europe, PO Box 506, 6700 AM Wageningen, The Netherlands

Keywords: Coevolution, Geographic Mosaic, Chemical mimicry, Local adaptation, *Phengaris*

Summary

The range of hosts exploited by a parasite is determined by several factors, including host availability, infectivity and exploitability. Each of these can be the target of natural selection on both host and parasite, which will determine the local outcome of interactions, and potentially lead to coevolution. However, geographical variation in host use and specificity has rarely been investigated. *Maculinea* (= *Phengaris*) butterflies are brood parasites of *Myrmica* ants that are patchily distributed across the Palæarctic and have been studied extensively in Europe. Here we review the published records of ant host use by the European *Maculinea* species, as well as providing new host ant records for more than 100 sites across Europe. This comprehensive survey demonstrates that while all but one of the *Myrmica* species found on *Maculinea* sites have been recorded as hosts, the most common is usually that exploited most. Host sharing and host switching are both relatively common, but there is evidence of specialization at many sites, which varies among *Maculinea* species. We show that most *Maculinea* display the features expected for coevolution to occur in a geographic mosaic, which has probably allowed these rare butterflies to persist in Europe.

1. Introduction

(a) Specificity in exploitative interactions

The vast majority of organisms have evolved to exploit the resources of other organisms, whether via herbivory, predation or parasitism [1]. Specialization on a small number of organisms can increase the efficiency of exploitation at the expense of generality [2], although there is still debate on the evolutionary mechanisms leading to such specialization [3, 4]. For parasites, which show long-term association with their hosts, the precise hosts that they exploit can be regarded as being determined by a series of proximate filters [5]. In turn, hosts must be encountered, infected and exploited [5, 6]. These filters can be the product of distribution, ecology and evolution. For example, hosts can only be encountered if their geographic distribution is included in the dispersal range of the parasite, and, all else being equal, more abundant hosts are more likely to be encountered than rare hosts. On the other hand, if parasites have evolved search strategies for particular hosts, this can change the “apparency” of hosts in the environment [7]. Whether a host can be infected may depend on general traits of the host unrelated to the parasite (such as physical or chemical barriers), or the evolution of particular parasite infection mechanisms [8]. Finally, exploitation of the host’s resources to produce new parasites may also be determined by general traits, such as immunocompetence and host vigour [9], or by specific interactions between the host and parasite phenotypes [10]. While some aspects of these filters may be entirely environmentally determined, others will have genetic components that are open to natural selection, and potentially coevolution. Genetic responses of both host and parasite, mediated by the environment (i.e. Gene × Gene × Environment interactions) are likely to lead to coevolution of parasites and hosts in a geographic mosaic [11]. One likely outcome of this mosaic coevolution is that which and how many hosts a parasite exploits varies among populations, although this is an aspect of host-parasite interactions that is rarely explored.

To examine the geographical mosaic of host specificity, we examine the variation in host use and specificity across Europe of *Maculinea* (= *Phengaris*) butterflies, which are well-known brood parasites of *Myrmica* ant colonies. The reliance of these butterflies on the sequential exploitation of both specific host plants and host ants means that they naturally occur in small, patchy populations that provide the ideal background for examining geographic mosaics.

(b) The infection and exploitation of *Myrmica* colonies by *Maculinea* butterflies

Maculinea (abbreviated as “*Ma.*” hereafter) species are obligate brood parasitic lycaenid butterflies, whose caterpillars initially feed on the developing seeds of specific host plants, but are taken in and raised by *Myrmica* (“*My.*”) colonies in their final larval instar [12, 13]. The caterpillars develop within the host ant nest during the autumn and spring, exploiting the resources of the colony, and will then either pupate there, or continue development for an additional year[14].

*Author for correspondence (DRNash@bio.ku.dk). ORCID 0000-0002-0462-6794.
†We dedicate this paper to the memory of Graham Elmes, who inspired so many *Maculinea* biologists to also consider the ants.

Although many ant species have associations with lycaenid butterflies [15], the only confirmed host ants of *Maculinea* butterflies are all from the genus *Myrmica*. Thomas et al. [12, 16] gave an overview of misidentifications of the ant species and misinterpretations of captive observations and field data in some initial research. Yamaguchi [17] mentions *Aphaenogaster japonica* as a host of *Ma. teleius* and *Ma. arionides* in Japan, but these have subsequently been shown to be misidentifications of *Myrmica kotokui* [18-20].

Of the four currently recognized species of *Maculinea* found in Europe [21] (see also below), *Ma. nausithous* and *Ma. teleius* initially develop on the flowers of great burnet (*Sanguisorba officinalis*), albeit in slightly different successional stages [15, 22] and positions on the plant [23-25], but often co-occur on the same sites. *Ma.alcon* develops on gentians (*Gentiana* spp. and *Gentianella* spp.), and *Ma. arion* develops on either thyme (*Thymus* spp.) or wild marjoram (*Origanum vulgare*) [13, 26]. After developing through three larval instars on these plants, the newly-moulted fourth instar exits the plant and descends to the soil surface where it awaits discovery by a foraging *Myrmica* worker [13, 27, 28].

Infection of *Myrmica* colonies (usually referred to as “adoption” [27, 29]) involves the larva being picked up by a worker ant and taken back to the ant nest, where it is placed amongst the ant brood. While the first foraging worker of any species of *Myrmica* to encounter a *Maculinea* larva under its food plant will almost always pick it up and take it back to the nest [27, 30], transfer to the brood chamber and initial integration may be highly species specific [29]. This specificity is largely mediated by the cuticular hydrocarbons that the fourth instar larvae produce, which mimic those of their host ants [31-34], with better-matching mimics achieving greater infection [34, 35]. However, in some populations that use multiple host ant species, this mimicry is imperfect, and seems to represent a compromise between mimicking different host *Myrmica* [34, 36]. Prior to adoption *Ma. arion* and *Ma. teleius* show a prolonged stereotypical sequence of behaviours before being picked up [25, 28, 30, 37], which may also convey non-chemical (e.g. tactile [28] or acoustic [38]) information, while *Ma.alcon* and *Ma. nausithous* are picked up quickly and with simpler behavioural sequences [25, 27, 29, 37].

Once inside the nest, *Maculinea* caterpillars have two *Myrmica* ant exploitation strategies. *Ma. teleius* and *Ma. arion* larvae feed as predators on the ant brood, while larvae of the *Ma.alcon* group (see below) have a more efficient “cuckoo” strategy where they are mainly fed by trophallaxis by the worker ants as if they were ant larvae [27, 39], but are also able to prey directly on the ant brood. While the larvae of *Ma. nausithous* are also predatory, they share some characteristics with the cuckoo species, feeding mostly on *Myrmica* eggs and small larvae [40]. Exploitation by *Maculinea* caterpillars severely reduces host ant fitness [34, 41-43], so there is strong selection on ants to identify these virulent parasites, which has in turn led to sophisticated strategies to misdirect the ants. Much closer chemical mimicry [44], involving additional species-specific secretions, is achieved after 4-6 days by caterpillars in the nests of their primary host species, while those adopted by secondary or non-host *Myrmica* species suppress this second phase of secretions and depend, usually unsuccessfully, on the chemical camouflage [44] afforded by simply acquiring nest odours from the host [32, 36, 45]. Caterpillars and pupae also mimic sounds produced by *Myrmica* queens [46], which increase their perceived value to the ant colony, although these signals are not specific within the genus *Myrmica* [47]. Despite these sophistications, *Maculinea* larvae and pupae are frequently killed by their host ants [6, 39, 48], especially under food stress [49].

(c) Previous studies on host ant specificity of *Maculinea* and the scope of this study

Initial results from Western Europe [12] suggested that each *Maculinea* species is adapted to exploit a different *Myrmica* host ant species. However, further work in the region quickly demonstrated that *Ma.alcon* shows geographical shifts in host ant specificity [50]. Over the last twenty years there have been a plethora of publications describing host ant use by *Maculinea* butterflies in Europe (table S1), which has more than doubled the number of *Myrmica* species that are known to be exploited and blurred the clear patterns that were initially thought to exist, leading some to question whether there really was any specificity [51].

The establishment of the MacMan EU research network [52] provided a unique opportunity to study host ant specificity at a pan-European scale, and to implement some more standardized methods of data collection, including the avoidance of artefacts that bedevilled certain early reports [16]. This study therefore aims to both synthesize the published information on host ant use by these butterflies, and to add original data collected during and after the MacMan project. To our knowledge, this study represents the most comprehensive analysis of geographic variation in host use for any parasite, and many of our results should be widely applicable to other host parasite systems.

(d) Conservation considerations

The occurrence of *Maculinea* butterflies in small, patchy populations, which are naturally vulnerable to extinction [53], together with changes in land use over the last century, have led to rapid declines of many populations and extinctions in several European countries [54, 55]. All *Maculinea* species in Europe were considered as vulnerable to global extinction in the first European data book for butterflies [56] and although

many have been reclassified in the latest version [57], declines have continued in many countries [58]. An understanding of the ant host use of *Ma. arion* in the U.K. has proven critical in both working out why it became extinct [54] and in its successful reintroduction [59]. However, the type of geographical variation in host specificity that we examine here means that the lessons from the populations in the U.K. may not be directly applicable elsewhere in Europe, and other reintroduction programs have been less immediately successful [55]. We hope that by highlighting the geographic mosaic of coevolution between *Maculinea* butterflies and their host ants we will aid future conservation efforts for these butterflies across Europe.

2. Methods

In order to investigate how different populations of *Maculinea* parasites have adapted to exploit their *Myrmica* ant hosts, ideally one should be able to examine how the three filters on host use (encounter, infection and exploitation) differ among sites. However, distinguishing between infection and exploitation in the field has proved impractical on a large scale, so the combined effect of these two filters was examined.

(a) The encounter filter

The use of specific host plants by the first three larval instars of *Maculinea* butterflies provides a constraint for the encounter filter. Only *Myrmica* colonies that are within ant worker foraging range of the larval food plants will be encountered by the fourth instar *Maculinea* larvae. As a general rule of thumb, it is usually assumed that the foraging range of *Myrmica* workers is largely restricted to within two meters of the nest [60], so that for this study we defined the encounter filter as the community of *Myrmica* ants that is present within two meters of host plants. The encounter filter is, of course, also open to change if either the parasites have evolved mechanisms to preferentially exploit host plants closer to host ant nests, or if *Myrmica* ants do not forage randomly with respect to host plants. Many other lycaenid butterflies that associate with specific ants use these ants as an oviposition cue [61-63], but the evidence for the use of ant-dependent oviposition in *Maculinea* has been controversial [24, 64-68]. The most recent evidence suggests that while *Maculinea* butterflies may lay their eggs in response to the presence of *Myrmica* ants, they do not distinguish between different species of *Myrmica* [69]. Hence our use of *Myrmica* distribution relative to overall host plant distribution as a measure of the encounter filter may underestimate the rate of encounters [35], but not the potential hosts that are encountered. Ovipositing female *Maculinea* butterflies do, however, select the buds of their food plants that are in particular growth-forms or phenological stages [24, 64, 70], and these may occur in microhabitats that may be associated with particular *Myrmica* species [24]. Ideally, therefore, the encounter filter should be the *Myrmica* community found around food plants on which eggs have been laid [35], although this effect is likely to be small, especially in comparison with difference in *Myrmica* communities between areas with and without food plants [60, 71, 72]. On the other hand, females may avoid plants on which others have laid eggs [73], which will lead to a more even distribution of eggs (and larvae) relative to *Myrmica* nests [65, 68]. Larvae leaving plants may also time their exit to maximize the chances of being encountered by *Myrmica* workers rather than any other ants, but this does not influence which *Myrmica* species they encounter [30].

For this study, the main method of assessing the *Myrmica* community close to food plants was the excavation of nests. Although there were small variations in methods used between groups, the majority of data was collected in the late spring or early summer, when food plants could be reliably identified and when fully grown *Maculinea* larvae and pupae were present in nests (see also the following section). Patches of food plants were identified (taking into account any information on where butterflies were flying and laying eggs the previous year), plants randomly selected, and then a circular area with a radius of 2 m searched for all *Myrmica* nests, which were subsequently excavated. To examine the wider geographical variation in the encounter filter (and because excavation of nests was not always practical or permitted on some sites), it was also measured on additional sites by setting out ant baits (sugar and/or protein) in areas within two meters of host plants. Baits were always placed more than 2 m, and usually more than 4 m, apart, so each bait that attracted *Myrmica* workers was assumed to correspond to a separate nest.

Myrmica ants were generally identified following [74, 75] or [76], but if an identification was doubtful, specimens were sent to *Myrmica* experts (Sándor Csősz, Graham Elmes, Alexander Radchenko or Bernhard Seifert) for identification. *Myrmica* species names follow the revision published in [77]. Where latitude, longitude or altitude of sites were not provided directly, they were estimated using www.mapcoordinates.net.

(b) The infection and exploitation filters

If *Maculinea* larvae successfully infect and exploit a *Myrmica* colony, they will develop within the nest during the autumn and winter. We therefore assessed the success of infection and exploitation by examining the presence and number of live fully grown larvae and pupae in the excavated nests. Comparison of the community of *Myrmica* ants within 2 m of host plants that housed overwintered *Maculinea* larvae and pupae with the total community was used as a measure of the specificity of these infection and exploitation filters.

There are several potential pitfalls to measuring host use by examining the presence of parasites within nests [16]. Critically, we restricted sampling to full-grown larvae, pupae, or emerging adults to measure only those individuals that had successfully exploited a *Myrmica* colony. However, we will still have recorded caterpillars that survived occasionally in rare benign nests of a host species to which they are ill-adapted [59]. Our datasets will inevitably also contain a few false positives, especially for predatory species of *Maculinea* [78], where the final depletion of ant brood caused a host ant colony to desert its parasitized nest site to be replaced by an offshoot from a different but neighbouring *Myrmica* species before the butterfly emerged [16, 41]. However, our sampling was so extensive, both geographically and quantitatively, that we are confident that the broad patterns that emerge are unequivocal. In addition to our original data, a comprehensive review of the literature on host ant use by *Maculinea* butterflies across Europe was conducted and combined with our newly collected data where appropriate (table S1).

For this study, the unit of replication was the site, defined as a distinct area in which a discrete population of one of the European *Maculinea* species was present, and specifically the part of that area where its host plant occurred. This was straightforward for *Ma. teleius* and *Ma. nausithous*, since patches of *Sanguisorba officinalis* are usually well defined. For *Ma. arion*, both *Thymus* and *Origanum* occur on many areas where the butterfly is found, so these were not distinguished. However, for *Ma. alcon* there are at least two distinct types of habitat used by the butterfly, primarily associated with two different host plants in the genus *Gentiana*. Some populations are found on wet heathlands and boggy meadows, and mostly use the marsh gentian, *G. pneumonanthe* as a host, while others are found on much drier areas, often at higher elevations, and mostly use the cross-leaved gentian *G. cruciata*. These two types of populations of *Ma. alcon* have been considered as two separate species in the past (*Ma. alcon* and *Ma. rebeli* respectively), but recent molecular studies have found no phylogenetic differentiation [79-83]. Nevertheless, we treat them separately here, since they occur in very different habitats, which are likely to have very different *Myrmica* communities, and because where examined each type secretes a distinctive pre-adoption chemical profile regardless of the local host ant species [84]. We will refer to the mostly *G. pneumonanthe*-using, hygrophilic form, as *Ma. alcon* H, and the mostly *G. cruciata*-using, xerophilic form as *Ma. alcon* X. Both these forms of *Ma. alcon* may also lay eggs on other gentian plants, particularly in central Europe and the Pyrenees, where the willow gentian, *Gentiana asclepiadea*, and various *Gentianella* spp. may be used as initial food plants [80, 85-87], but the distinction between wet and dry sites is usually clear. It has also become apparent in recent years that the specific name *Ma. rebeli* (originally described as a subspecific form of *Ma. alcon* [88]) has been incorrectly applied to the *G. cruciata*-using form of *Ma. alcon* [89], and should be associated with *Ma. alcon* from another distinct habitat - above the treeline (>1500m. above sea level) in the Alps. Some data are now available on the host ant use of such alpine populations (putative *Ma. alcon* form *rebeli*), which have been found to lay their eggs exclusively on *Gentianella rhaetica* [90], so we will discuss these separately where appropriate, and we also treat one record from the literature as belonging to this form (see below). These sites were not included in the main analyses. On sites where multiple *Maculinea* were known to fly, identification of *Maculinea* larvae found in nests followed [91], although it is not possible to distinguish between pupae of *Ma. nausithous* and *Ma. teleius* based on morphology. Unless their identities were confirmed by rearing to adulthood, the small number of pupae found on sites that were known to support both of these *Maculinea* were therefore excluded from the database and our analysis.

Methods used for the analysis of the contents of the database are described where appropriate below. Statistical analyses were conducted using R version 3.5 [92], JMP version 13.2 (©SAS institute 2016) and PAST version 3.16 [93].

3. Results and discussion

In total we collected data on *Myrmica* host ant availability from 419 sites (figure 1, table S1), of which 214 also provided direct records of host ant use. Most sites (83.3%) only supported a single *Maculinea*, but 70 sites supported up to four different *Maculinea* (figure 1), most notably *Ma. teleius* and *Ma. nausithous*, which co-occurred on 51 of these sites. The sites on which each *Maculinea* occurred accorded provided a good sample of their known European distributions (figure S1). All *Maculinea* were generally found at lower altitudes towards the north, and in more mountainous areas towards the south (ANCOVA on log(altitude); latitude: $F_{1,487} = 249.78$, $p < 0.0001$; figure S2), but the characteristic altitude for each *Maculinea* varied ($F_{4,487} = 9.94$, $p < 0.0001$, *Maculinea* × latitude interaction: $F_{1,487} = 5.90$, $p = 0.0001$), with *Ma. alcon* X being found at significantly higher altitudes (Predicted altitude at mean latitude (49.46° N): *Ma. alcon* X = 377 m, *Ma. alcon* H = 199 m, *Ma. arion* = 248 m, *Ma. nausithous* = 199 m, *Ma. teleius* = 257 m).

(c) The encounter filter – *Myrmica* communities on *Maculinea* sites

A total of 17 different *Myrmica* species were found on *Maculinea* sites across Europe (figure 2, figure S3), with the number on any particular site varying from 1 to 9 (figure 2). These belonged to four of the species groups currently recognized within *Myrmica* (the *rubra*, *scabrinodis*, *lobicornis* and *schencki* groups) [77] and well

supported by recent phylogenetic analyses [94-96]. Members of the *lobicornis* group tended to be rare, but the other three groups all included very common species (figure 2). The most common species across sites was *My. scabrinodis* (more than 10,000 nests examined), and the rarest was *My. tulinae* (6 nests among over 2,400 *Myrmica* nests examined at Krakow in Poland) although the identification of this latter species should be treated with caution [77] and molecular data suggest it may not be a good species [96].

In order to get a reliable estimate of the total *Myrmica* community on a site, a minimum number of samples of *Myrmica* nests is necessary. We estimated this minimum number by calculating the difference between the bias-corrected Chao-1 estimate of true *Myrmica* species richness [97] and the observed *Myrmica* species richness for each site and comparing this with the number of nests examined (either the number excavated or the number of baits that attracted *Myrmica* workers). This showed that above a sample of 24 nests, most sites had reached a stable estimate of the *Myrmica* community (figure S4), so a minimum sample of 25 nests was used to compare ant communities (which reduced the number of sites with sufficient data to 199). Somewhat surprisingly given the known European distribution of *Myrmica* species [77, 98] and their overall distribution across sites (figure S3), the number of *Myrmica* species recorded on a site in this dataset did not vary significantly with latitude, longitude or altitude (Poisson GLM: Likelihood-ratio (L-R) $\chi^2 = 0.541$, $d.f. = 1$, $p = 0.462$; L-R $\chi^2 = 0.018$, $d.f. = 1$, $p = 0.893$; L-R $\chi^2 = 0.067$, $d.f. = 1$, $p = 0.795$ respectively). The size of the *Myrmica* community was, however, significantly different among sites occupied by the different *Maculinea* (L-R $\chi^2 = 20.87$, $d.f. = 4$, $p = 0.0003$, figure 2).

To compare the composition of ant communities, we used Non-metric Multi-Dimensional Scaling (NMDS) ordination based on Bray-Curtis dissimilarities in the R-package *vegan* [99], which successfully (stress value of 0.134) reduced the data down to three axes. Discriminant analyses based on these axes showed that the *Myrmica* community on sites occupied by different *Maculinea* could be split into two major divisions by the first NMDS axis: those found on drier sites occupied by *Ma. arion* and *Ma.alcon* X, and those found on the wetter sites occupied by *Ma. nausithous*, *Ma. teleius* and *Ma.alcon* H (Wilks' $\lambda_{328} = 0.360$, $p < 0.0001$, 91.3% correctly classified), with no difference within the xerophilic group (Wilks' $\lambda_{328} = 0.997$, $p = 0.977$, 56.0% correctly classified) and a small but significant difference between *Ma. nausithous* and the other *Maculinea* in the hygrophilic group (Wilks' $\lambda_{328} = 0.786$, $p < 0.0001$, 50.8% correctly classified), although there was some overlap of all groups (figure 2). There was a significant change in *Myrmica* communities with latitude (ANCOVA; $F_{1,244} = 7.025$, $p = 0.009$), but this varied for the different *Maculinea* (figure S5; Among *Maculinea*: $F_{4,244} = 99.14$, $p < 0.001$; *Maculinea* \times latitude interaction: $F_{4,244} = 5.11$, $p < 0.001$).

(d) The infection and exploitation filters – *Myrmica* specificity

Of the 17 *Myrmica* species found on *Maculinea* sites, overwintered larvae or pupae were found in the nests of all but one (*My. lobulicornis*; figure S6). However, the number of hosts used on any site was generally low (figure 2), with 70% of sites examined having only a single host ant, and *My. scabrinodis* is the only species that has been recorded as a host for all five *Maculinea*. There was significant variation among *Maculinea* in the number of hosts used (Poisson GLM; L-R $\chi^2 = 15.27$, $d.f. = 4$, $p = 0.004$), and patterns of host use differed both geographically and phylogenetically. To examine geographical similarity, spatial autocorrelation analysis was carried out in GenoDive 2.0b23 [100] separately for each *Maculinea* by comparing a matrix of pairwise distances between sites that hosted that *Maculinea* with a matrix of Jaccard similarities in their community of host ants. Distances were divided into ten classes for each *Maculinea* so that each contained 10% of the sample pairs. Moran's r was then used to examine how the similarity of hosts changed with geographical distance (figure 3).

As has been suggested previously, *Ma. nausithous* has the smallest range of hosts, being restricted to a single host (either *My. rubra* or *My. scabrinodis*) on all but one site, where it was found in the nests of four *Myrmica* species (one being *My. tulinae*, so this record should be treated with caution). However, there was little spatial structure in this pattern, with the less common host (*My. scabrinodis*) being used across Europe.

Ma.alcon H was only found in the nests of more than two *Myrmica* species on a single site (figure 3) and was generally split into two geographically distinct groups (figure 3). Those in the north-west of Europe exploited species within the *rubra* species group (*My. rubra* and *My. ruginodis*), while in the remainder of Europe, ants from the *scabrinodis* species group (*My. scabrinodis*, *My. aloba*, *My. slovacae* and *My. vandeli*) were hosts. The only exception to this was one site in Belgium where *My. scabrinodis* was used as a host on a site where *My. rubra* and *My. ruginodis* were the main hosts. This pattern was reflected in the spatial autocorrelation analysis which shows that hosts were very similar among sites up to 1000 km apart, but significantly different beyond this. There has been one report of a *My. schencki* nest housing the only *Ma.alcon* found on a Polish site in 2013 [101], but in a more extensive survey at the same site the following year only *My. scabrinodis* nests were found to be exploited, and in larger numbers [101], so we consider that this record (of a prepupa) was probably a case of nest-takeover [16] and have not included it in the database.

Ma.alcon X generally exploited *My. sabuleti* or *My. schencki*, with all other recorded hosts being exploited on sites where one of these two species was also a host (except for two sites where *My. specioides* was the only host). Host communities were less similar beyond a distance of around 800 km, but sites towards the extreme south west and north east all used *My. schencki* as a host. An outwardly similar pattern was found in *Ma. teleius*, except in this case the two main hosts were *My. rubra* and *My. scabrinodis*, and there were two sites where *My. slovaca* was a host (in one case shared with *My. specioides*), but there was no real spatial structure in host use.

Ma. arion sites tended to fall into two groups which exploited *Scabrinodis* group ants (*My. lonae*, *My. sabuleti*, *My. scabrinodis* or *My. specioides*) in most of Europe, and *My. schencki* (sometimes in combination with other species) in north-east Europe (although one site where *My. schencki* is used as a host is known from Italy). The relatively small area where *My. schencki* dominates is reflected in the spatial autocorrelation results, where communities go from being significantly similar to significantly dissimilar over a distance of around 400 km, but whether this is representative of the whole range of *Ma. arion* is unclear, as the *My. schencki*-using populations of this butterfly are on the eastern border of the area we covered. Distinguishing between the closely related *My. sabuleti* and *My. lonae* can be problematic in northern Europe [77] and sharing of mitochondrial haplotypes suggests that they may not be separate species. Hence it is unclear whether the apparent switch from using *My. sabuleti* to using *My. lonae* observed in the most northern sites is a genuine host switch.

The putative *Ma.alcon* form *rebeli* populations from above the tree-line in the Austrian Alps have been shown to use *My. sulcinodis* as a host [90], which is the most common *Myrmica* species recorded on the two sites examined (table S1). It is notable that another record of an *Ma.alcon* group caterpillar found in a *My. sulcinodis* nest was made by David Jutzeler in the 1980's in the Swiss Alps at a site above the tree-line, and where eggs were found on plants in the *Gentianella germanica* complex [102, 103]. This site was subsequently destroyed in a landslide (Jutzeler, personal communication), but we consider it highly likely that this was also the putative form *rebeli*.

To investigate whether there was any evidence of specificity (i.e. that *Myrmica* hosts were not simply exploited in the proportion that they were available on each site), the distribution of the number of nests that were found to contain *Maculinea* larvae or pupae was compared with the distribution of nests that did not contain *Maculinea* using contingency tables. For sites with only a single potential host ant species present, host ant specificity cannot be examined. For sites with two or more potential host ant species present, the significance of the deviation from homogeneity in the contingency table was tested using a χ^2 statistic, the probability of which was tested by random reassignment of the number of nests to each cell in the table 100,000 times, with the constraint that marginal total were retained. The observed value of the χ^2 statistic was then compared with the distribution of χ^2 statistics generated from the 100,000 permutations. This analysis showed that the probability that the pattern of host ant use found simply reflected host ant availability was often low, being below the conventional $\alpha = 0.05$ in 55 out of 160 cases where it could be calculated (figure 4). The (log-transformed) probability that *Myrmica* nests were infected and exploited in the proportion encountered was negatively correlated with the log of the number of *Myrmica* nests in which *Maculinea* larvae or pupae were found on a site (ANCOVA; $F_{1,150} = 54.6$, $p < 0.0001$) and differed among the different *Maculinea* (among *Maculinea*: $F_{4,150} = 6.05$, $p = 0.0002$; *Maculinea* \times log(number of infected nests): $F_{4,150} = 2.33$, $p = 0.058$; figure 4), with *Ma.alcon* X and *Ma. arion* showing a greater proportion of sites showing specificity than the other four *Maculinea*. However, for all *Maculinea*, the proportion of sites where significant ($\alpha = 0.05$) specificity was found was higher than the 5% predicted by chance unless only a single infested nest was found (figure 4).

A recent theoretical model of the evolution of host use in *Maculinea* [104], which explored the roles of *Myrmica* species abundance and similarity in host phenotypes on the evolution of specificity. This concluded that two stable strategies are likely to exist: 1) specialization on a single, abundant host, or 2) use of multiple hosts when host abundance is lower and hosts at least partially share phenotypes related to *Maculinea* infection ability.

(e) Local adaptation

If specialization on one or a few hosts is favoured by natural selection, as appears to be the case in *Maculinea* brood parasites, then all else being equal, butterflies will benefit most by specializing on the most abundant host. This implies that they should exploit common hosts more frequently than expected based on their encounter rate and fail to successfully exploit less abundant hosts [34, 105] – a form of frequency-dependent selection. To examine this, we plotted the proportional exploitation of each host used (i.e. the proportion of nests containing overwintered *Maculinea* larvae or pupae for each site that belonged to that host) against the proportional availability of that host (i.e. the proportion of *Myrmica* nests excavated that belonged to that host) (figure 5). If host *Myrmica* were exploited in the proportion available, sites would lie along the 1:1 line. To test whether there were deviations from the 1:1 line, 1-proportion Z- tests were carried out comparing the proportion of points that were above or below the 1:1 line for each *Maculinea* with the null hypothesis that

there would be equally many above and below the line, subdividing the x-axis into those *Myrmica* that were more abundant (> 50% of the nests available) and those that were less abundant (<50% of available nests) at each site. This showed that the expected pattern was indeed seen for *Ma. alcon H* and *Ma. arion*, with over-exploitation of common hosts (Single proportion Z-test; $Z = +3.92, p < 0.001$; $Z = +3.15, p < 0.001$ respectively) and under-exploitation of rare hosts ($Z = -5.82, p < 0.001$; $Z = -6.67, p < 0.001$ respectively). *Ma. alcon X*, *Ma. nausithous* and *Ma. teleius* all under-exploited rare hosts ($Z = -3.86, p < 0.001$; $Z = -2.97, p = 0.002$; $Z = -3.05, p = 0.001$ respectively) but did not overexploit common hosts ($Z = -0.816, p = 0.793$; $Z = +0.242, p = 0.404$; $Z = +0.756, p = 0.775$ respectively). The general under-exploitation of rare hosts across all *Maculinea* is not unexpected, since populations that continue to exploit only rare hosts are unlikely to persist, particularly in the predatory species *Ma. arion* and *Ma. teleius*, which have the greatest effect on host colony fitness [39]. It has been shown that that a minimum of 50% [39] to 68% [59] of the larval population of these *Maculinea* must be adopted by the main host (in host-specific populations) for intrinsic growth rates to be positive. Nevertheless, some apparently maladapted populations are found, for example at one isolated Polish site [106], the rare *My. lobicornis* is the main host, despite only making up 13% of the available *Myrmica* nests [107]. In the more efficient and less virulent cuckoo-feeding species *Ma. alcon H* and *Ma. alcon X* this threshold is expected to be considerably lower and has been shown empirically to be as low as 13% [39]. It must be remembered that which *Myrmica* species are rare or abundant close to food plants is likely to change over time, both in response to environmental changes and in response to the selection pressure exerted by *Maculinea* [71, 108, 109]. The failure of *Ma. alcon X* and *Ma. nausithous* to adapt to use the most common hosts therefore could indicate that host switching may be more difficult for these *Maculinea*, or alternatively that their host ants respond more rapidly to exploitation. Even if, overall, the most commonly available hosts tend to be overexploited and the less abundant hosts avoid parasitism, suggesting local adaptation by *Maculinea*, there are still several cases where local maladaptation (specialization on a locally rare host) is evident for all *Maculinea*.

(f) Host sharing and host switching

In cases where multiple host ants are used on the same site, there may still be specificity that reflects coevolution with hosts, particularly if hosts share similar parasite defence mechanisms. All else being equal, we might expect more phylogenetically related hosts to have more similar defence mechanisms, so that host sharing and switching between hosts may be modulated by *Myrmica* phylogeny. To test this hypothesis we calculated the Phylogenetic Species Evenness (PSE) index [110] for each site on which multiple *Myrmica* hosts were used both for the complete *Myrmica* community and the community of *Myrmica* nests that supported overwintered *Maculinea* larvae or pupae, using the R package *picante* [111]. This index provides a measure of phylogenetic diversity within a community that takes species abundances into account and should be independent of species richness, and ranges between 0 (species are highly related and/or there is very high skew in their abundances) to 1 (all species are unrelated and equally abundant). For the analysis, which requires a phylogenetic tree with branch lengths, a modified version of the tree produced by [95] was used, with the species that they did not include added based on relationships inferred from [96]. The values of this index for the encountered and exploited *Myrmica* communities at each site were then compared with a paired t-test separately for every *Maculinea* which had multiple sites where more than one *Myrmica* was exploited simultaneously (i.e. all except *Ma. nausithous*). This showed that PSE was significantly lower for the host than the general *Myrmica* community for *Ma. alcon H* (Paired t-test; $t = -3.55, d.f. = 11, p = 0.0053$), but not for the other *Maculinea* (*Ma. alcon X*: $t = +0.432, d.f. = 17, p = 0.671$; *Ma. arion*: $t = -0.180, d.f. = 8, p = 0.862$; *Ma. teleius*: $t = +1.34, d.f. = 13, p = 0.206$). This suggests that shared hosts are not generally more closely related for three of the four *Maculinea* where it could be tested. This pattern is also seen when species groups are compared. In *Ma. alcon H* and to some extent in *Ma. arion*, shared hosts are usually from the same *Myrmica* species group, but in *Ma. alcon X* and *Ma. teleius* there are many sites where hosts are shared among *Myrmica* from different species groups, although this is dominated by sharing between two particular pairs of species, *My. sabuleti* and *My. schencki* in *Ma. alcon X*, and *My. scabrinodis* and *My. rubra* in *Ma. teleius*. This latter pairing in particular contrasts with *Ma. alcon H*, where these two species tend not to be shared. One possible reason for this could be that the very common *My. scabrinodis* appears to consist of at least two cryptic species that are morphologically indistinguishable, but are genetically quite distinct [96, 112] and differ in their ecology and behaviour [60].

Infection and exploitation of *Myrmica* colonies by *Maculinea* butterflies critically depends on circumventing the ants' self-/non-self-recognition system by mimicking their cuticular hydrocarbons [31, 33-35, 45]. These cuticular recognition cues are generally species-specific in *Myrmica*, but also show some similarities among species [33-35, 113]. Hence it is expected that sharing of, or switching between, certain hosts may be more easily achieved if they have more similar sets of hydrocarbons, even if some degree of specificity is maintained. For example, different populations of *Ma. alcon H* in Denmark exploit *My. rubra* or *My. ruginodis*, or both species simultaneously, but never exploit *My. scabrinodis*. Although it is relatively common [114] and is exploited almost exclusively in most of Europe (figure 3). This is because the closely related *My. rubra* and *My. ruginodis* have much more similar cuticular hydrocarbons than either has with *My. scabrinodis*. Although producing a hydrocarbon profile that is a reasonable mimic of both *My. rubra* and *My. ruginodis* may have costs in terms of somewhat reduced infectivity [34], this step is still within the levels of natural variability

found in Danish *Ma. alcon* *H* populations, whereas the hydrocarbons of *My. scabrinodis* are not [34]. Does this pattern hold for the other *Maculinea* butterflies across Europe? Pre-adoption larvae of *Ma. alcon* *X* in eastern Austria that simultaneously exploit *My. sabuleti* and *My. schencki* have been shown to produce a set of cuticular hydrocarbons that simultaneously match different portions of the cuticular profile of these rather distantly related *Myrmica* [36]. Although, in general, cuticular hydrocarbons of ants tend to evolve rather slowly [115], when the cuticular hydrocarbons of different *Myrmica* are compared, they generally show rather little phylogenetic similarity [113]. This may be because of the additional roles of ant cuticular hydrocarbons in species [116] and possibly mate [117, 118] recognition, which may lead to selection on their divergence during speciation. Whether host-sharing and -switching is related to cuticular hydrocarbon similarity therefore remains an open question, but an interesting example is provided by *My. vandeli* which has cuticular hydrocarbons practically indistinguishable from *My. scabrinodis* [113]. It is thought that *My. vandeli* may act as a temporary social parasite of *My. scabrinodis* [77], and therefore has evolved a similar set of hydrocarbons to itself integrate into colonies of this host. Hence it is notable that we only found *My. vandeli* being exploited by *Maculinea* on sites on which they also exploit *My. scabrinodis* (figure 3).

(g) Fitting the patterns together – The geographic mosaic of coevolution

There is considerable variation in host use of all the European *Maculinea*, but this does not imply lack of specificity in their interactions with *Myrmica* ants at a population level [51]. The different initial host plants used by the different *Maculinea* are found in different types of habitat, which correlate with the different *Myrmica* communities found there (figure 2), and set the boundaries of the encounter filter for the *Maculinea* butterflies. However, there is still a lot of variation and overlap in *Myrmica* communities, so that even very close sites rarely have precisely the same community present. This fulfils one of the three main tenets of the geographical mosaic theory of coevolution [11], that there should be a mosaic of hot-spots and cold-spots for coevolution. Coevolution between *Maculinea* butterflies and any particular *Myrmica* species can only take place when that *Myrmica* species is present at a particular site. The second tenet of geographic mosaic theory is that there should be geographical variation in the outcome of coevolution. At the majority of sites, *Maculinea* butterflies appear to be locally adapted to their host *Myrmica*, in that they are effectively exploiting the most abundant *Myrmica* host. The identity of this abundant host varies geographically, however, and overall, but with a fair number of exceptions, we found that each *Maculinea* exploits a constant host ant species on multiple sites across areas of thousands rather than hundreds of km² in Europe, with major host switches occurring in each species at greater distances (figure 3): small, population-scale switches as found for *M. alcon* *H* in Denmark [34] and for *Ma. alcon* *X* in Poland, Italy and parts of central Europe [32, 35, 80, 105] are comparatively unusual, but quite apparent within the snapshot of time sampled during this study. However, there are a substantial minority of sites where *Maculinea* populations seem to be maladapted and specialize on a rare *Myrmica*. This is precisely what is expected if there is ongoing coevolution between *Maculinea* butterflies and their *Myrmica* hosts, because there is also expected to be selection on hosts to avoid infection and exploitation, and the lack of parasitism of a common *Myrmica* species can be seen as local adaptation of the host to avoid parasitism. The third component of the geographic mosaic theory is (limited) trait-remixing among populations. This is also a characteristic of *Maculinea* populations. Although *Maculinea* are quite limited in their dispersal when examined using mark-release-recapture techniques or behavioural observations [119–122], their genetic structure suggests occasional long-distance dispersal [123–126], so allowing the maintenance of the genetic diversity required for coevolution across a wide geographic area.

4. Conclusion

The life cycle of *Maculinea* butterflies makes them entirely dependent on *Myrmica* ants, which suffer serious fitness losses as a consequence. Both sides in the interaction are therefore expected to impose severe selection pressure on the other, so providing the basic conditions for coevolution. In this paper we have looked at *Maculinea* evolution in terms of how they exploit and are adapted to different *Myrmica* species, however it must also be borne in mind that coevolution within species pairs is likely to be the norm [34], and that the patterns we examine here can only be interpreted in the framework of (albeit very tight) diffuse coevolution. The variety of exploitation strategies used, however, mean that whether and how coevolution proceeds is likely to be different for the different *Maculinea*, which is reflected in our results.

The “cuckoo”-feeding *Ma. alcon* group are more dependent on integration within the *Myrmica* colony, as they need to constantly interact with the ants that feed them [39], which requires sustained deception [32, 33]. This seems to have led to large scale geographic mosaic of exploitation, in which the same or related hosts are used over distances of a thousand kilometres or so. Specificity is generally high within sites, but local adaptation is variable. It is frequent in *M. alcon* *H*, but much less so in *Ma. alcon* *X*, which generally interacts with a larger potential host community.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The best-known predatory *Maculinea*, *Ma. arion*, shows a somewhat similar pattern, but on a smaller spatial scale, which tends to lead to local adaptation to the most common *Myrmica* host. However, it also seems to be able to exploit the nests of other *Myrmica* on the same sites, albeit so inefficiently that a population cannot be supported by the presence of the mal-adapted host species alone [59], which leads to generally lower apparent specificity. This is taken to a further extreme by *Ma. teleius*, which has the largest host range, largely exploits *Myrmica* in much the same proportion as they are encountered and shows no apparent geographical structure in its host use, giving little evidence of ongoing coevolution. It should be recalled, however, that our data will contain more false positives for these latter species, caused by ants switching nest sites during the pupal stage following the disproportionate damage inflicted through carnivory on the typically small nests of their main hosts. For example, roughly half the instances for *Ma. arion* emerging from nests occupied by *My. scabrinodis* in the UK resulted from nest-takeover and half from genuine survival with this normally unsuitable host [41, 59]

Ma. nausithous is unusual in many ways. Its hosts seem to be very limited compared with the other *Maculinea*, which has led to an apparent lack of local adaptation in its host use. Despite using the same major hosts as *Ma. alcon* *H*, it does not seem to be able to use the related *Myrmica* species in the same way, suggesting that it does not rely on the same type of cuticular mimicry to gain access to its food [127]. This is possibly because its apparent need, when small, to eat ant eggs in the heavily protected host chambers that contain queens [40] requires closer host-specific mimicry of its main host that is simply ineffective with other *Myrmica* species. Essentially there is also little evidence of coevolution between this species and its *Myrmica* hosts, but for completely different reasons to *Ma. teleius*, which are at opposite ends of the host specificity spectrum. In the case of *Ma. nausithous*, there is little geographic variation in outcomes, while in the case of *Ma. teleius*, we seem to lack coevolutionary hotspots.

It is clear that the idea of “one *Myrmica* for one *Maculinea*” does not hold across Europe as a whole but is often true in smaller regions within the continent. It is equally clear that different *Maculinea* have different propensities for using multiple *Myrmica* hosts or shifting host. Such alternation in the use of a network of hosts by parasites is expected to be a common outcome of antagonistic coevolution [128]. Similar patterns to those shown by *Maculinea* butterflies and their *Myrmica* hosts have also been suggested for other brood parasites [129], but not documented in such detail. Host defences and in what ways parasites overcome them is critical to how such antagonistic coevolution proceeds. The variability found within the genus *Maculinea* is potentially very useful for exploring this aspect of coevolution, but we still need to learn more about exploitation and defence strategies of both partners. In particular, the difference between the patterns of host use by *Ma. alcon* that occur in dry and wet habitats is an intriguing pattern that merits further investigation, although it seems that somewhat different chemical deception strategies may be involved [84]. What leads to the very limited host range of *Ma. nausithous* compared to the closely related *Ma. teleius* is also unclear and should be investigated in more detail.

Within the lycaenid butterflies, ant brood parasitism is a very uncommon strategy [130], despite most lycaenid butterflies having some sort of symbiotic relationship with ants [131]. Despite its rarity, brood parasitism has evolved multiple times within the *Lycaenidae*, but has rarely led to diverse lineages [130, 131], suggesting that it is a strategy that does not persist long over evolutionary time spans. The genus *Maculinea* is relatively young (ca. 2.5 MYA) compared with other lycaenid genera [79], and the conservation concern over all its constituent species reflects its natural rarity and vulnerability to local extinction. However, such local extinctions are also likely to be a natural part of the geographic mosaic in which these butterflies coevolve with *Myrmica* ants. The propensity for host ant switching shown by all *Maculinea* is likely to be the key to their persistence on regional and continental scales. Whether this continues to be the case under increased pressure from human activities and climate change remains to be seen.

5. Figure legends

Figure 1. Summary of the 419 *Maculinea* sites examined in this study. *a)* Map showing the distribution of the sites colour-coded according to the *Maculinea* present on each site. Where more than one *Maculinea* occurs on a site, symbols are dissected according to the *Maculinea* present. *b)* UpSet [132] showing the intersecting sets of *Maculinea* across all sites.

Figure 2. Diversity of the *Myrmica* species community found on *Maculinea* sites. *a)* The seventeen *Myrmica* species, belonging to four *Myrmica* species groups found on *Maculinea* sites, showing their phylogenetic relatedness [95, 96] and overall abundance (total number of nests) across all sites. *b)* Histograms showing the numbers of *Myrmica* species recorded from sites that supported each *Maculinea*. Counts are based on sites where at least 25 nests were excavated, or 25 baits attracted *Myrmica* ants. Sites supporting more than one *Maculinea* are counted separately for each. *c)* NMDS ordination plot of the *Myrmica* communities found on sites supporting the different *Maculinea* based on Bray-Curtis dissimilarities for sites where more than 25 nests were excavated, or 25 baits attracted *Myrmica* ants. Convex hulls define the extent of the communities associated with each *Maculinea*. Sites supporting more than one *Maculinea* are counted separately for each. The ordination scores of the *Myrmica* species are also plotted. *d)* Histograms showing the numbers of *Myrmica* species whose nests contained overwintered *Maculinea* larvae or pupae recorded from sites that supported each *Maculinea*. Sites supporting more than one *Maculinea* are counted separately for each.

Figure 3. Host use across the five main *Maculinea* groups. For each *Maculinea* the top panel shows the geographical distribution all sites where overwintered larvae or pupae were found in excavated nests ("host nests"). Each symbol is a pie diagram showing the proportion of host nests that belonged to each *Myrmica* species, with size proportional to the number of host nests examined. The central panel shows an Euler diagram with the set of hosts *Myrmica* used for each *Maculinea*. The area assigned to each *Myrmica* species is proportional to the number of sites where that ant was a host. The bottom panel shows a spatial autocorrelogram showing how the similarity between the host community used on different sites (based on Jaccard similarity indices) varies with distance between sites. Pairs of sites are grouped into 10 distance classes with equal sample size (which therefore are different for each *Maculinea*). Markers are placed at the maximum distance for each class and are coloured depending on whether the correlation among communities (Moran's r) is significantly different from zero ($p < 0.05$; filled circles) or not (open circles).

Figure 4. Relationship between specificity and sampling effort for each *Maculinea*. Specificity was tested by comparing the distribution of *Myrmica* nests of each species found close to food plants with that of the nests of those species found to contain overwintered larvae or pupae of *Maculinea* for all sites where more than one *Myrmica* species was recorded. Each point in the graph represents the estimated probability that hosts are used in the proportion available for each *Maculinea* at each site, coloured according to the *Maculinea* present. Least-squared regression lines are shown in the same colour for each *Maculinea*. Note that for clarity, the y-axis is reversed so that sites showing greater specificity are further from the x-axis. The conventional level of statistical significance ($\alpha = 0.05$) is indicated by a dashed line, and all points above this can be considered to show significant specificity. Where each regression line crosses this threshold gives an estimate of the number of nests containing overwintered *Maculinea* caterpillars or pupae needed to achieve a 50% probability of detecting significant specificity.

Figure 5. Comparison of proportional availability and proportional exploitation of *Myrmica* nests for each *Maculinea*. Each point represents one *Myrmica* species on one site where more than one potential host *Myrmica* species was recorded. The x-axis represents the relative abundance of that *Myrmica* species at the site, and the y-axis is the proportion of all host nests at that site of that *Myrmica* species. If *Myrmica* species are exploited in the proportion that they are available, points should be evenly distributed above and below the 1:1 diagonal (shown as a dashed line). Points below this line (in the grey area) represent under-exploitation, while points above the line represent over-exploitation. The relationship is visualized for each *Maculinea* with a LOWESS regression line (smoothing parameter $\alpha = 0.4$).

Additional Information

Acknowledgments

We thank Raldi Bakker, Birgit Binzenhöfer, Maria da Conceição Rodrigues, Márta Ferencz, Krzysztof Frackiel, Anja Freese-Hager, Adi Geyer, Adam Górnicki, Patrick Gros, Mariusz Gwardian, Przemysław Klimczuk, Bernadette Lázár, Andreas Lomborg, Gyöngyvér Molnár, Enikő Német, Andreas Nunner, Bas Oteman, Krzysztof Pałka, Bernd Reiser, Tobias Schiefer, Martina Sasic, Markus Schwibinger, Piotr Skórka, Ewa B. Śliwińska, Vizauer Tibor-Csaba, Robert Völkl and Adam Warecki for field assistance. Koos Boomsma, Joanna Kajzer, Bálint Markó, Christian Stettmer and Michał Włostowski provided valuable support for the study.

Ethics

All novel research for this project was carried out in accordance with the regulations operating in the specific European counties where sites were located, including obtaining local permits to work in protected areas where appropriate.

Data Accessibility

The dataset supporting this article has been uploaded as part of the Supplementary Material.

Authors' Contributions

AT, JAT and DRN conceived the study. AT compiled the database, with input from all other authors. DRN carried out the analysis and wrote the manuscript, with input from JAT and AT. All authors read, corrected and approved the submitted version of the manuscript. Original data for the manuscript was provided by all authors, whose contributions are given in detail in table S1.

Competing Interests

The authors declare no competing interests

Funding

Most of the data for this project was collected with support from the EU-project MacMan (EVK2-CT-2001-00126). The database was developed as part of the EU Marie-Curie fellowship InDifferAnt (Project No: 237116) awarded to AT and a Danish National Research Foundation grant to the Centre for Social Evolution, Copenhagen (DNRF57). Subsequent development of the database was supported by the EU Marie-Curie Career Integration fellowship AntLab, and by two Bolyai János scholarships from the Hungarian Academy of Sciences (MTA). Other support for the fieldwork incorporated into the database came from the EU BiodivERsA-Project CLIMIT, The Bayerisches Landesamt für Umwelt (LfU), The VolkswagenStiftung, The Deutsche Bundesstiftung Umwelt (DBU). PSO was supported by grants from the European Investment Funds in the FEDER/COMPETE/POCI– Operacional Competitiveness and Internacionalization Programme, under Project POCI-01-0145-FEDER-006958, National Funds from the Portuguese Foundation for Science and Technology (FCT), under the project UID/AGR/04033/2013 and the Municipality of Vila Real (Biodiversity Program), Portugal. PN was supported by the Polish National Science Centre grant DEC-2013/11/B/NZ8/00912. The State Committee for Scientific Research (grants no.3 P04G 026 24 and 2 P04G024 30), Poland, Polish Ministry of Science and Higher Education (subsidy for maintaining the research potential of the Faculty of Biology and Chemistry, University of Bialystok).

References

- [1] Lafferty, K. D. & Kuris, A. M. 2002 Trophic strategies, animal diversity and body size. *Trends Ecol Evol* **17**, 507-513. (DOI:10.1016/S0169-5347(02)02615-0).
- [2] Whitlock, M. C. 1996 The red queen beats the jack-of-all-trades: The limitations on the evolution of phenotypic plasticity and niche breadth. *Am Nat* **148**, S65-S77. (DOI:10.1086/285902).
- [3] Forister, M. L., Dyer, L. A., Singer, M. S., Stireman, J. O., 3rd & Lill, J. T. 2012 Revisiting the evolution of ecological specialization, with emphasis on insect-plant interactions. *Ecology* **93**, 981-991. (DOI:10.1890/11-0650.1).
- [4] Futuyma, D. J. & Moreno, G. 1988 The Evolution of Ecological Specialization. *Annu Rev Ecol Syst* **19**, 207-233. (DOI:10.1146/annurev.es.19.110188.001231).
- [5] Combes, C. 2001 *Parasitism: the ecology and evolution of intimate interactions*, University of Chicago Press.
- [6] Nash, D. R., Als, T. D. & Boomsma, J. J. 2011 Survival and growth of parasitic *Maculinea alcon* caterpillars (Lepidoptera, Lycaenidae) in laboratory nests of three *Myrmica* ant species. *Insectes Soc* **58**, 391-401. (DOI:10.1007/s00040-011-0157-y).
- [7] Feeny, P. 1976 Plant apparency and chemical defense. In *Biochemical interaction between plants and insects: Proceedings of the fifteenth annual meeting of the Phytochemical Society of North America* (eds. J. W. Wallace & R. L. Mansell), pp. 1-40. New York, Plenum Press.
- [8] Schmid Hempel, P. 2011 *Evolutionary parasitology - the integrated study of infections, immunology, ecology, and genetics*. Oxford, Oxford University Press.
- [9] Beldomenico, P. M. & Begon, M. 2010 Disease spread, susceptibility and infection intensity: vicious circles? *Trends Ecol Evol* **25**, 21-27. (DOI:10.1016/j.tree.2009.06.015).
- [10] Little, T. J., Watt, K. & Ebert, D. 2006 Parasite-host specificity: experimental studies on the basis of parasite adaptation. *Evolution* **60**, 31-38. (DOI:10.1111/j.0014-3820.2006.tb01079.x).
- [11] Thompson, J. N. 1999 Specific hypotheses on the geographic mosaic of coevolution. *Am Nat* **153**, S1-S14. (DOI:10.1086/303208).
- [12] Thomas, J. A., Elmes, G. W., Wardlaw, J. C. & Woyciechowski, M. 1989 Host specificity among *Maculinea* butterflies in *Myrmica* ant nests. *Oecologia* **79**, 452-457. (DOI:10.1007/Bf00378660).
- [13] Elmes, G. & Thomas, J. A. 1987 Die Gattung *Maculinea*. In *Tagfalter und ihre Lebensräume: Arten, Gefährdung und Schutz* (ed. W. Geiger), pp. 354-368. Basel, Schweizerischer Bund für Naturschutz.
- [14] Thomas, J. A., Elmes, G. W. & Wardlaw, J. C. 1998 Polymorphic growth in larvae of the butterfly *Maculinea rebeli*, a social parasite of *Myrmica* ant colonies. *Proc Biol Sci* **265**, 1895-1901. (DOI:10.1098/rspb.1998.0517).
- [15] Fiedler, K. 2006 Ant-associates of Palaearctic lycaenid butterfly larvae (Hymenoptera: Formicidae; Lepidoptera: Lycaenidae) – a review. *Myrmecological News* **9**, 77-87.
- [16] Thomas, J. A., Elmes, G. W., Schönrogge, K., Simcox, D. J. & Settele, J. 2005 Primary hosts, secondary hosts and non-hosts: common confusions in the interpretation of host specificity in *Maculinea* butterflies and other social parasites of ants. In *Studies on the Ecology and Conservation of Butterflies in Europe* (eds. J. Settele, E. Kühn & J. A. Thomas), pp. 99-104, Pensoft Publisher Sofia-Moscow.
- [17] Yamaguchi, S. 1988 *The life histories of five myrmecophilous lycaenid butterflies of Japan*. Kodansha, Japan.
- [18] Hiraga, S. 2003 Reidentifying of host ants of *Maculinea arionides*. *Yadoriga* **196**, 31-34.
- [19] Maruyama, M., Komatsu, T., Kudo, S., Shimada, T. & Kinomura, K. 2013 *The guests of Japanese ants*. Minamiyana, Japan, Tokai University Press.
- [20] Ueda, S., Komatsu, T., Itino, T., Arai, R. & Sakamoto, H. 2016 Host-ant specificity of endangered large blue butterflies (*Phengaris* spp., Lepidoptera: Lycaenidae) in Japan. *Sci Rep* **6**, 36364. (DOI:10.1038/srep36364).
- [21] Fric, Z., Wahlberg, N., Pech, P. & Zrzavy, J. 2007 Phylogeny and classification of the *Phengaris*-*Maculinea* clade (Lepidoptera : Lycaenidae): total evidence and phylogenetic species concepts. *Syst Entomol* **32**, 558-567. (DOI:10.1111/j.1365-3113.2006.00387.x).
- [22] Kőrösi, Á., Örvössi, N., Batáry, P., Harnos, A. & Peregovits, L. 2012 Different habitat selection by two sympatric *Maculinea* butterflies at small spatial scale. *Insect Conserv Divers* **5**, 118-126. (DOI:10.1111/j.1752-4598.2011.00138.x).
- [23] Figurny, E. & Woyciechowski, M. 1998 Flowerhead selection for oviposition by females of the sympatric butterfly species *Maculinea teleius* and *M. nausithous* (Lepidoptera : Lycaenidae). *Entomol Gen* **23**, 215-222. (DOI:10.1127/entom.gen/23/1998/215).
- [24] Thomas, J. A. & Elmes, G. W. 2001 Food-plant niche selection rather than the presence of ant nests explains oviposition patterns in the myrmecophilous butterfly genus *Maculinea*. *Proc Biol Sci* **268**, 471-477. (DOI:10.1098/rspb.2000.1398).
- [25] Thomas, J. A. 1984 The behaviour and habitat requirements of *Maculinea nausithous* (the dusky large blue butterfly) and *M. teleius* (the scarce large blue) in France. *Biol Conserv* **28**, 325-347. (DOI:10.1016/0006-3207(84)90040-5).
- [26] Thomas, J. A. 1995 The ecology and conservation of *Maculinea arion* and other European species of large blue butterfly. In *Ecology and conservation of butterflies* (ed. A. S. Pullin), pp. 180-197, Springer.
- [27] Elmes, G. W., Thomas, J. A. & Wardlaw, J. C. 1991 Larvae of *Maculinea rebeli*, a Large-Blue Butterfly, and Their *Myrmica* Host Ants - Wild Adoption and Behavior in Ant-Nests. *J Zool* **223**, 447-460. (DOI:10.1111/j.1469-7998.1991.tb04775.x).
- [28] Frohawk, F. W. 1916 Further observations on the last stage of the larva of *Lycaena arion*. *Trans Entomol Soc London* **1915**, 313-316.
- [29] Als, T. D., Nash, D. R. & Boomsma, J. J. 2001 Adoption of parasitic *Maculinea alcon* caterpillars (Lepidoptera : Lycaenidae) by three *Myrmica* ant species. *Anim Behav* **62**, 99-106. (DOI:10.1006/anbe.2001.1716).
- [30] Thomas, J. A. 2002 Larval niche selection and evening exposure enhance adoption of a predacious social parasite, *Maculinea arion* (large blue butterfly), by *Myrmica* ants. *Oecologia* **132**, 531-537. (DOI:10.1007/s00442-002-1002-9).
- [31] Akino, T., Knapp, J. J., Thomas, J. A. & Elmes, G. W. 1999 Chemical mimicry and host specificity in the butterfly *Maculinea rebeli*, a social parasite of *Myrmica* ant colonies. *Proc Biol Sci* **266**, 1419-1426. (DOI:10.1098/rspb.1999.0796).


- [32] Thomas, J. A., Elmes, G. W., Sielezniew, M., Stankiewicz-Fiedurek, A., Simcox, D. J., Settele, J. & Schönrogge, K. 2013 Mimetic host shifts in an endangered social parasite of ants. *Proc Biol Sci* **280**, 20122336. (DOI:10.1098/rspb.2012.2336).
- [33] Elmes, G., Akino, T., Thomas, J., Clarke, R. & Knapp, J. 2002 Interspecific differences in cuticular hydrocarbon profiles of *Myrmica* ants are sufficiently consistent to explain host specificity by *Maculinea* (large blue) butterflies. *Oecologia* **130**, 525-535. (DOI:10.1007/s00442-001-0857-5).
- [34] Nash, D. R., Als, T. D., Maile, R., Jones, G. R. & Boomsma, J. J. 2008 A mosaic of chemical coevolution in a large blue butterfly. *Science* **319**, 88-90. (DOI:10.1126/science.1149180).
- [35] Casacci, L. P., Schönrogge, K., Thomas, J. A., Balletto, E., Bonelli, S. & Barbero, F. in review Host specificity pattern and chemical deception in a social parasite of ants. *Sci Rep*.
- [36] Schlick-Steiner, B. C., Steiner, F. M., Höttinger, H., Nikiforov, A., Mistrik, R., Schafellner, C., Baier, P. & Christian, E. 2004 A butterfly's chemical key to various ant forts: intersection-odour or aggregate-odour multi-host mimicry? *Naturwissenschaften* **91**, 209-214. (DOI:10.1007/s00114-004-0518-8).
- [37] Fiedler, K. 1990 New information on the biology of *Maculinea nausithous* and *M. teleius* (Lepidoptera: Lycaenidae). *Nota lepid* **12**, 246-256.
- [38] Sala, M., Casacci, L. P., Balletto, E., Bonelli, S. & Barbero, F. 2014 Variation in butterfly larval acoustics as a strategy to infiltrate and exploit host ant colony resources. *PLoS One* **9**, e94341. (DOI:10.1371/journal.pone.0094341).
- [39] Thomas, J. A. & Elmes, G. W. 1998 Higher productivity at the cost of increased host-specificity when *Maculinea* butterfly larvae exploit ant colonies through trophallaxis rather than by predation. *Ecol Entomol* **23**, 457-464. (DOI:10.1046/j.1365-2311.1998.00153.x).
- [40] Elfferich, N. W. 1998 New facts on the life history of the dusky large blue *Maculinea nausithous* (Lepidoptera: Lycaenidae) obtained by breeding with *Myrmica* ants in plaster nests. *Deinsea* **4**, 97-102.
- [41] Thomas, J. A. & Wardlaw, J. C. 1992 The capacity of a *Myrmica* ant nest to support a predacious species of *Maculinea* butterfly. *Oecologia* **91**, 101-109. (DOI:10.1007/BF00317247).
- [42] Wardlaw, J. C., Thomas, J. A. & Elmes, G. W. 2000 Do *Maculinea rebeli* caterpillars provide vestigial mutualistic benefits to ants when living as social parasites inside *Myrmica* ant nests? *Entomol Exp Appl* **95**, 97-103. (DOI:10.1046/j.1570-7458.2000.00646.x).
- [43] Witek, M., Slipinski, P., Naumiec, K., Krupski, A., Babik, H., Walter, B., Symonowicz, B. & Dubiec, A. 2016 Performance of *Myrmica* ant colonies is correlated with the presence of social parasites. *Ecol Entomol* **41**, 284-291. (DOI:10.1111/een.12299).
- [44] Nash, D. R. & Boomsma, J. J. 2008 Communication between hosts and social parasites. In *Sociobiology of Communication - an interdisciplinary approach* (eds. P. d'Ettorre & D. P. Hughes), pp. 55-79. Oxford, Oxford University Press.
- [45] Schönrogge, K., Wardlaw, J. C., Peters, A. J., Everett, S., Thomas, J. A. & Elmes, G. W. 2004 Changes in chemical signature and host specificity from larval retrieval to full social integration in the myrmecophilous butterfly *Maculinea rebeli*. *J Chem Ecol* **30**, 91-107. (DOI:10.1023/B:JOEC.0000013184.18176.a9).
- [46] Barbero, F., Thomas, J. A., Bonelli, S., Balletto, E. & Schönrogge, K. 2009 Queen ants make distinctive sounds that are mimicked by a butterfly social parasite. *Science* **323**, 782-785. (DOI:10.1126/science.1163583).
- [47] Thomas, J. A., Schönrogge, K., Bonelli, S., Barbero, F. & Balletto, E. 2010 Corruption of ant acoustical signals by mimetic social parasites: *Maculinea* butterflies achieve elevated status in host societies by mimicking the acoustics of queen ants. *Commun Integr Biol* **3**, 169-171. (DOI:10.4161/cib.3.2.10603).
- [48] Witek, M., Skorka, P., Sliwiska, E. B., Nowicki, P., Moron, D., Settele, J. & Woyciechowski, M. 2011 Development of parasitic *Maculinea teleius* (Lepidoptera, Lycaenidae) larvae in laboratory nests of four *Myrmica* ant host species. *Insectes Soc* **58**, 403-411. (DOI:10.1007/s00040-011-0156-z).
- [49] Elmes, G. W., Wardlaw, J. C., Schönrogge, K., Thomas, J. A. & Clarke, R. T. 2004 Food stress causes differential survival of socially parasitic caterpillars of *Maculinea rebeli* integrated in colonies of host and non-host *Myrmica* ant species. *Entomol Exp Appl* **110**, 53-63. (DOI:10.1111/j.0013-8703.2004.00121.x).
- [50] Elmes, G. W., Thomas, J. A., Hammarstedt, O., Munguira, M. L., Martín, J. & Van Der Made, J. 1994 Differences in host-ant specificity between Spanish, Dutch and Swedish populations of the endangered butterfly, *Maculinea alcon* (Denis et Schiff.) (Lepidoptera). *Memorabilia Zool* **48**, 55-68.
- [51] Pech, P., Fric, Z. & Konvicka, M. 2007 Species-specificity of the *Phengaris* (*Maculinea*)-*Myrmica* host system: Fact or myth? (Lepidoptera : Lycaenidae; Hymenoptera : Formicidae). *Sociobiology* **50**, 983-1003.
- [52] Settele, J., Thomas, J. A., Boomsma, J. J., Kuehn, E., Nash, D. R., Anton, C., Woyciechowski, M. & Varga, Z. 2002 *MACULINEA* butterflies of the habitats directive and European red list as indicators and tools for conservation and MANagment (MacMan). *Verh Ges Ökol* **32**, 63.
- [53] Saccheri, I., Kuussaari, M., Kankare, M., Vikman, P., Fortelius, W. & Hanski, I. 1998 Inbreeding and extinction in a butterfly metapopulation. *Nature* **392**, 491-494. (DOI:10.1038/33136).
- [54] Thomas, J. 1980 Why Did the Large Blue Become Extinct in Britain? *Oryx* **15**, 243. (DOI:10.1017/s0030605300024625).
- [55] Wynhoff, I. 1998 Lessons from the reintroduction of *Maculinea teleius* and *M. nausithous* in the Netherlands. *J Insect Conserv* **2**, 47-57. (DOI:10.1023/A:1009692723056).
- [56] Van Swaay, C. A. M. & Warren, M. S. 1999 *Red data book of European butterflies (Rhopalocera)*. Strasbourg, Council of Europe Publishing.
- [57] van Swaay, C., Cuttelod, A., Collins, S., Maes, D., Munguira, M. L., Šašić, M., Settele, J., Verovnik, R., Verstrael, T., Warren, M., et al. 2010 *European Red List of Butterflies*, &IUCN (International Union for Conservation of Nature)&IUCN.
- [58] van Swaay, C., Maes, D., Collins, S., Munguira, M. L., Sasic, M., Settele, J., Verovnik, R., Warren, M., Wiemers, M., Wynhoff, I., et al. 2011 Applying IUCN criteria to invertebrates: How red is the Red List of European butterflies? *Biol Conserv* **144**, 470-478. (DOI:10.1016/j.biocon.2010.09.034).
- [59] Thomas, J. A., Simcox, D. J. & Clarke, R. T. 2009 Successful conservation of a threatened *Maculinea* butterfly. *Science* **325**, 80-83. (DOI:10.1126/science.1175726).
- [60] Elmes, G. W., Thomas, J. A., Wardlaw, J. C., Hochberg, M. E., Clarke, R. T. & Simcox, D. J. 1998 The ecology of *Myrmica* ants in relation to the conservation of *Maculinea* butterflies. *J Insect Conserv* **2**, 67-78. (DOI:10.1023/A:1009696823965).
- [61] Fraser, A. M., Tregenza, T., Wedell, N., Elgar, M. A. & Pierce, N. E. 2002 Oviposition tests of ant preference in a myrmecophilous butterfly. *J Evolution Biol* **15**, 861-870. (DOI:10.1046/j.1420-9101.2002.00434.x).
- [62] Pierce, N. E. & Elgar, M. A. 1985 The Influence of Ants on Host Plant-Selection by *Jalmenus evagoras*, a Myrmecophilous Lycaenid Butterfly. *Behav Ecol Sociobiol* **16**, 209-222. (DOI:10.1007/Bf00310983).
- [63] Atsatt, P. R. 1981 Ant-dependent food plant selection by the mistletoe butterfly *Ogyris amaryllis* (Lycaenidae). *Oecologia* **48**, 60-63. (DOI:10.1007/BF00346988).

- [64] Fürst, M. A. & Nash, D. R. 2010 Host ant independent oviposition in the parasitic butterfly *Maculinea alcon*. *Biol Lett* **6**, 174-176. (DOI:10.1098/rsbl.2009.0730).
- [65] van Dyck, H., Talloen, W., Feenstra, V. & Wynhoff, I. 2000 Does the presence of ant nests matter for oviposition to a specialized myrmecophilous *Maculinea* butterfly? *Proc Biol Sci* **267**, 861-866. (DOI:10.1098/rspb.2000.1082).
- [66] Musche, M., Anton, C., Worgan, A. & Settele, J. 2006 No Experimental Evidence for Host Ant Related Oviposition in a Parasitic Butterfly. *J Insect Behav* **19**, 631-643. (DOI:10.1007/s10905-006-9053-0).
- [67] Patricelli, D., Barbero, F., La Morgia, V., Casacci, L. P., Witek, M., Balletto, E. & Bonelli, S. 2011 To lay or not to lay: oviposition of *Maculinea arion* in relation to *Myrmica* ant presence and host plant phenology. *Anim Behav* **82**, 791-799. (DOI:10.1016/j.anbehav.2011.07.011).
- [68] Wynhoff, I., Grutters, M. & van Langevelde, F. 2008 Looking for the ants: selection of oviposition sites by two myrmecophilous butterfly species *Anim Biol* **58**, 371-388. (DOI:10.1163/157075608X383683).
- [69] Patricelli, D., Barbero, F., Occhipinti, A., Berte, C. M., Bonelli, S., Casacci, L. P., Zebelo, S. A., Crocoll, C., Gershenzon, J., Maffei, M. E., et al. 2015 Plant defences against ants provide a pathway to social parasitism in butterflies. *Proc Biol Sci* **282**. (DOI:10.1098/rspb.2015.1111).
- [70] Van Dyck, H. & Regniers, S. 2010 Egg spreading in the ant-parasitic butterfly, *Maculinea alcon*: from individual behaviour to egg distribution pattern. *Anim Behav* **80**, 621-627. (DOI:10.1016/j.anbehav.2010.06.021).
- [71] Nash, D. R. & Andersen, A. 2015 *Maculinea* sommerfugle og stikmyrer på danske heder - coevolution i tid og rum. *Flora og Fauna* **121**, 133-141.
- [72] Soares, P. O., Crespi, A. L., Rodrigues, M. C. & Arnaldo, P. S. 2012 The habitat vegetational structure and the success of the blue alcon, *Maculinea alcon* (Denis & Schiffermuller). *Plant Biosyst* **146**, 1-6. (DOI:10.1080/11263504.2011.558671).
- [73] Sielezniew, M. & Stankiewicz-Fiedurek, A. M. 2013 Behavioural evidence for a putative oviposition-detering pheromone in the butterfly, *Phengaris* (*Maculinea*) *teleius* (Lepidoptera: Lycaenidae). *Eur J Entomol* **110**, 71-80. (DOI:10.14411/eje.2013.009).
- [74] Wardlaw, J. C., Elmes, G. W. & Thomas, J. A. 1998 Techniques for studying *Maculinea* butterflies: II. Identification guide to *Myrmica* ants found on *Maculinea* sites in Europe. *J Insect Conserv* **2**, 119-127. (DOI:10.1023/a:1009621522615).
- [75] Seifert, B. 1996 *Ameisen: beobachten, bestimmen*, Naturbuch Verlag.
- [76] Czechowski, W., Radchenko, A. & Czechowska, W. 2002 *The ants (Hymenoptera, Formicidae) of Poland*. Warszawa, Museum and Institute of Zoology PAS 200 p.
- [77] Radchenko, A. G. & Elmes, G. W. 2010 *Myrmica ants (Hymenoptera: Formicidae) of the Old World*. Warszawa, Poland, Natura optima dux Foundation.
- [78] Thomas, J. A., Schönrogge, K. & Elmes, G. W. 2005 Specializations and host associations of social parasites of ants. In *Insect Evolutionary Ecology: Proceedings of the Royal Entomological Society's 22nd Symposium* (eds. M. D. E. Fellowes, G. J. Holloway & J. Rolff), pp. 479-518. London, Royal Entomological Society.
- [79] Als, T. D., Vila, R., Kandul, N. P., Nash, D. R., Yen, S. H., Hsu, Y. F., Mignault, A. A., Boomsma, J. J. & Pierce, N. E. 2004 The evolution of alternative parasitic life histories in large blue butterflies. *Nature* **432**, 386-390. (DOI:10.1038/nature03020).
- [80] Sielezniew, M., Rutkowski, R., Ponikwicka-Tyszko, D., Dzienkańska, I. & Švitra, G. 2012 Differences in genetic variability between two ecotypes of the endangered myrmecophilous butterfly *Phengaris* (= *Maculinea*) *alcon* - the setting of conservation priorities. *Insect Conserv Divers* **5**, 223-236. (DOI:10.1111/j.1752-4598.2011.00163.x).
- [81] Koubínová, D., Dinca, V., Dapporto, L., Voda, R., Suchan, T., Vila, R. & Alvarez, N. 2017 Genomics of extreme ecological specialists: multiple convergent evolution but no genetic divergence between ecotypes of *Maculinea alcon* butterflies. *Sci Rep* **7**, 13752. (DOI:10.1038/s41598-017-12938-8).
- [82] Tartally, A., Kelager, A., Fürst, M. A. & Nash, D. R. 2016 Host plant use drives genetic differentiation in syntopic populations of *Maculinea alcon*. *PeerJ* **4**, e1865. (DOI:10.7717/peerj.1865).
- [83] Ugelvig, L. V., Vila, R., Pierce, N. E. & Nash, D. R. 2011 A phylogenetic revision of the *Glaucopsyche* section (Lepidoptera: Lycaenidae), with special focus on the *Phengaris-Maculinea* clade. *Mol Phylogenet Evol* **61**, 237-243. (DOI:10.1016/j.ympev.2011.05.016).
- [84] Schönrogge, K., Barbero, F., Bonelli, S., Casacci, L. P., Tartally, A., Sielezniew, M., Nash, D. R., Maes, D., Settele, J. & Thomas, J. A. in prep A tale of two species: chemical similarities and differences between two forms of *Maculinea alcon*. *Insect Conserv Divers*.
- [85] Bolt, D., Neumeyer, R., Rey, A. & Sohni, V. 2010 Kleiner Moorbläuling (Lepidoptera: Lycaenidae: *Phengaris alcon*) und Echte Knotenameisen (Hymenoptera: Formicidae: *Myrmica*) am Pfannenstiel (Kanton Zürich, Schweiz). *Entomol Helvet* **3**, 27-48.
- [86] Gros, P. 2002 Nachweis von *Maculinea rebeli* (HIRSCHKE, 1904) aus dem Bundesland Salzburg (Lepidoptera: Lycaenidae). *Z Arb Gem Öst Ent* **54**, 13-23.
- [87] Tessier, M. 2012 Inventaire de l'Azuré des mouillères *Maculinea alcon* (DENIS & SCHIFFERMÜLLER, 1775) (Lepidoptera Lycaenidae) en Ariège. I. Résultats préliminaires. *Bull Soc Linn Bordeaux* **147**, 129-139.
- [88] Hirschke, H. 1905 Eine neue hochalpine Form der *Lycaena alcon* F. aus den steirischen Alpen. *Jahresber Wiener Entomol Ver* **15**, 1-3.
- [89] Kudrna, O. & Fric, Z. F. 2013 On the identity and taxonomic status of *Lycaena alcon rebeli* HIRSCHKE, 1905 — a long story of confusion and ignorance resulting in the fabrication of a "ghost species" (Lepidoptera: Lycaenidae). *Nachr Entomol Ver Apollo* **34**, 117-124.
- [90] Tartally, A., Koschuh, A. & Varga, Z. 2014 The re-discovered *Maculinea rebeli* (Hirschke, 1904): Host ant usage, parasitoid and initial food plant around the type locality with taxonomical aspects (Lepidoptera, Lycaenidae). *ZooKeys*, 25-40. (DOI:10.3897/zookeys.406.7124).
- [91] Sliwiska, E. B., Nowicki, P., Nash, D. R., Witek, M., Settele, J. & Woyciechowski, M. 2006 Morphology of caterpillars and pupae of European *Maculinea* species (Lepidoptera : Lycaenidae) with an identification table. *Entomol Fennica* **17**, 351-358.
- [92] R Core Team. 2018 R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. (3.5.0 ed).
- [93] Hammer, Ø., Harper, D. A. T. & Ryan, P. D. 2001 PAST: Paleontological Statistics Software Package for education and data analysis. *Palaeontol Electr* **4**, 1-9.
- [94] Jansen, G., Savolainen, R. & Vepsäläinen, K. 2010 Phylogeny, divergence-time estimation, biogeography and social parasite-host relationships of the Holarctic ant genus *Myrmica* (Hymenoptera: Formicidae). *Mol Phylogenet Evol* **56**, 294-304. (DOI:10.1016/j.ympev.2010.01.029).
- [95] Jansen, G., Vepsäläinen, K. & Savolainen, R. 2011 A phylogenetic test of the parasite-host associations between *Maculinea* butterflies (Lepidoptera: Lycaenidae) and *Myrmica* ants (Hymenoptera: Formicidae). *Eur J Entomol* **108**, 53-62. (DOI:10.14411/eje.2011.007).
- [96] Ebsen, J. R., Boomsma, J. J. & Nash, D. R. in preparation Phylogeography and cryptic speciation in the *Myrmica scabrinodis* (Nylander, 1846) species complex (Hymenoptera: Formicidae). *Insect Conserv Divers*.
- [97] Chao, A. 1984 Nonparametric-Estimation of the Number of Classes in a Population. *Scand J Stat* **11**, 265-270.

- [98] Cushman, J. H., Lawton, J. H. & Manly, B. F. 1993 Latitudinal patterns in European ant assemblages: variation in species richness and body size. *Oecologia* **95**, 30-37. (DOI:10.1007/BF00649503).
- [99] Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., et al. 2018 vegan: Community Ecology Package. R package version 2.5-2. <https://CRAN.R-project.org/package=vegan>.
- [100] Meirmans, P. G. & Van Tienderen, P. H. 2004 GENOTYPE and GENODIVE: two programs for the analysis of genetic diversity of asexual organisms. *Mol Ecol Notes* **4**, 792-794.
- [101] Sielezniew, M., Bystrowski, C., Deoniziak, K. & Da Costa, J. M. 2015 An unexpected record of *Myrmica schencki* EMERY, 1895 as a secondary host ant of the hygrophilous form of a small and isolated population of the Alcon Blue butterfly *Phengaris* (= *Maculinea*) *alcon* (DENIS et SCHIFFERMÜLLER, 1775) (Lepidoptera, Lycaenidae) in NE Poland. *Polish J Entomol* **84**, 49-59. (DOI:10.1515/pjen-2015-0005).
- [102] Jutzeler, D. 1988 Fund von *Maculinea rebeli* (Hirschke, 1904) im Glarnerland (Lepidoptera, Lycaenidae). *Mitt Entomol Ges Basel* **384**, 124-125.
- [103] Jutzeler, D. 1989 *Maculinea rebeli* (HIRSCHKE): Ein Raupenfund im Glarnerland (Lepidoptera: Lycaenidae). *Mitt Entomol Ges Basel* **39**, 92-93.
- [104] Assis, R. A. d., Cecconello, M. d. S., Casacci, L. P., Barbero, F., Assis, L. M. E. d., Venturino, E. & Bonelli, S. 2018 A theory and a mathematical model for the evolution of single and multiple host behavior in a parasite-host system (*Maculinea*-*Myrmica*). *Ecol Complex* **34**, 178-187. (DOI:10.1016/j.ecocom.2017.11.007).
- [105] Tartally, A., Nash, D. R., Lengyel, S. & Varga, Z. 2008 Patterns of host ant use by sympatric populations of *Maculinea alcon* and *M. 'rebeli'* in the Carpathian Basin. *Insectes Soc* **55**, 370-381. (DOI:10.1007/s00040-008-1015-4).
- [106] Sielezniew, M. & Rutkowski, R. 2012 Population isolation rather than ecological variation explains the genetic structure of endangered myrmecophilous butterfly *Phengaris* (= *Maculinea*) *arion*. *J Insect Conserv* **16**, 39-50. (DOI:10.1007/s10841-011-9392-9).
- [107] Sielezniew, M., Dziekańska, I. & Stankiewicz-Fiedurek, A. M. 2010a Multiple host-ant use by the predatory social parasite *Phengaris* (= *Maculinea*) *arion* (Lepidoptera, Lycaenidae). *J Insect Conserv* **14**, 141-149. (DOI:10.1007/s10841-009-9235-0).
- [108] Hochberg, M. E., Clarke, R. T., Elmes, G. W. & Thomas, J. A. 1994 Population dynamic consequences of direct and indirect interactions involving a large blue butterfly and its plant and red ant hosts. *J Anim Ecol* **63**, 375-391. (DOI:10.2307/5555).
- [109] Thomas, J. A., Elmes, G. W., Clarke, R. T., Kim, K. G., Munguira, M. L. & Hochberg, M. E. 1997 Field evidence and model predictions of butterfly-mediated apparent competition between gentian plants and red ants. *Acta Oecol* **18**, 671-684. (DOI:10.1016/S1146-609x(97)80050-1).
- [110] Helmus, M. R., Bland, T. J., Williams, C. K. & Ives, A. R. 2007 Phylogenetic measures of biodiversity. *Am Nat* **169**, E68-E83. (DOI:10.1086/511334).
- [111] Kembel, S. W., Cowan, P. D., Helmus, M. R., Cornwell, W. K., Morlon, H., Ackerly, D. D., Blomberg, S. P. & Webb, C. O. 2010 Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* **26**, 1463-1464. (DOI:10.1093/bioinformatics/btq166).
- [112] Ebsen, J. R., Nash, D. R. & Boomsma, J. J. 2005 Cryptic *Myrmica* species among the hosts of *Maculinea* butterflies. In *Studies on the ecology and conservation of butterflies in Europe - Vol. 2: species ecology along a European gradient: Maculinea butterflies as a model* (eds. J. Settele, J. Thomas & E. Kühn), p. 171. Sofia, Pensoft.
- [113] Guillem, R. M., Drijfhout, F. P. & Martin, S. J. 2016 Species-Specific Cuticular Hydrocarbon Stability within European *Myrmica* Ants. *J Chem Ecol* **42**, 1052-1062. (DOI:10.1007/s10886-016-0784-x).
- [114] Als, T. D., Nash, D. R. & Boomsma, J. J. 2002 Geographical variation in host-ant specificity of the parasitic butterfly *Maculinea alcon* in Denmark. *Ecol Entomol* **27**, 403-414. (DOI:10.1046/j.1365-2311.2002.00427.x).
- [115] van Wilgenburg, E., Symonds, M. R. E. & Elgar, M. A. 2011 Evolution of cuticular hydrocarbon diversity in ants. *J Evolution Biol* **24**, 1188-1198. (DOI:10.1111/j.1420-9101.2011.02248.x).
- [116] Oldham, N. J., Morgan, E. D., Agosti, D. & Wehner, R. 1999 Species recognition from postpharyngeal gland contents of ants of the *Cataglyphis bicolor* group. *J Chem Ecol* **25**, 1383-1393. (DOI:10.1023/A:1020935011325).
- [117] Beibl, J., D'Ettore, P. & Heinze, J. 2007 Cuticular profiles and mating preference in a slave-making ant. *Insectes Soc* **54**, 174-182. (DOI:10.1007/s00040-007-0929-6).
- [118] Oppelt, A., Spitzenpfeil, N., Kroiss, J. & Heinze, J. 2008 The significance of intercolonial variation of cuticular hydrocarbons for inbreeding avoidance in ant sexuals. *Anim Behav* **76**, 1029-1034. (DOI:10.1016/j.anbehav.2008.05.020).
- [119] Nowicki, P. & Vrabec, V. 2011 Evidence for positive density-dependent emigration in butterfly metapopulations. *Oecologia* **167**, 657-665. (DOI:10.1007/s00442-011-2025-x).
- [120] Bonelli, S., Vrabec, V., Witek, M., Barbero, F., Patricelli, D. & Nowicki, P. 2013 Selection on dispersal in isolated butterfly metapopulations. *Popul Ecol* **55**, 469-478. (DOI:10.1007/s10144-013-0377-2).
- [121] Timuş, N., Craioveanu, C., Sitaru, C., Rus, A. & Rákossy, L. 2013 Differences in adult phenology, demography, mobility and distribution in two syntopic ecotypes of *Maculinea alcon* (*cruciata* vs. *pneumonanthe*) (Lepidoptera: Lycaenidae) from Transilvania (Romania). *Entomol Rom* **18**, 21-30.
- [122] Kőrösi, Á., Örvössy, N., Batáry, P., Kover, S. & Peregovits, L. 2008 Restricted within-habitat movement and time-constrained egg laying of female *Maculinea rebeli* butterflies. *Oecologia* **156**, 455-464. (DOI:10.1007/s00442-008-0986-1).
- [123] Sielezniew, M., Patricelli, D., Rutkowski, R., Witek, M., Bonelli, S. & Bus, M. M. 2015 Population genetics of the endangered obligatorily myrmecophilous butterfly *Phengaris* (= *Maculinea*) *arion* in two areas of its European range. *Insect Conserv Divers* **8**, 505-516. (DOI:10.1111/icad.12129).
- [124] Andersen, A., Simcox, D. J., Thomas, J. A. & Nash, D. R. 2014 Assessing reintroduction schemes by comparing genetic diversity of reintroduced and source populations: A case study of the globally threatened large blue butterfly (*Maculinea arion*). *Biol Conserv* **175**, 34-41. (DOI:10.1016/j.biocon.2014.04.009).
- [125] Ugelvig, L. V., Andersen, A., Boomsma, J. J. & Nash, D. R. 2012 Dispersal and gene flow in the rare, parasitic Large Blue butterfly *Maculinea arion*. *Mol Ecol* **21**, 3224-3236. (DOI:10.1111/j.1365-294X.2012.05592.x).
- [126] Vanden Broeck, A., Maes, D., Kelager, A., Wynhoff, I., WallisDeVries, M. F., Nash, D. R., Oostermeijer, J. G. B., Van Dyck, H. & Mergeay, J. 2017 Gene flow and effective population sizes of the butterfly *Maculinea alcon* in a highly fragmented, anthropogenic landscape. *Biol Conserv* **209**, 89-97. (DOI:10.1016/j.biocon.2017.02.001).
- [127] Witek, M., Casacci, L. P., Barbero, F., Patricelli, D., Sala, M., Bossi, S., Maffei, M., Woyciechowski, M., Balletto, E. & Bonelli, S. 2013 Interspecific relationships in co-occurring populations of social parasites and their host ants. *Biol J Linn Soc* **109**, 699-709. (DOI:10.1111/bij.12074).
- [128] Nuismer, S. L. & Thompson, J. N. 2006 Coevolutionary alternation in antagonistic interactions. *Evolution* **60**, 2207-2217. (DOI:10.1111/j.0014-3820.2006.tb01858.x).

- [129] Feeney, W. E., Welbergen, J. A. & Langmore, N. E. 2014 Advances in the Study of Coevolution Between Avian Brood Parasites and Their Hosts. *Annu Rev Ecol Evol Syst* **45**, 227-246. (DOI:10.1146/annurev-ecolsys-120213-091603).
- [130] Pierce, N. E. 1995 Predatory and parasitic Lepidoptera: carnivores living on plants. *J Lepid Soc* **49**, 412-453.
- [131] Pierce, N. E., Braby, M. F., Heath, A., Lohman, D. J., Mathew, J., Rand, D. B. & Travassos, M. A. 2002 The ecology and evolution of ant association in the Lycaenidae (Lepidoptera). *Annu Rev Entomol* **47**, 733-771. (DOI:10.1146/annurev.ento.47.091201.145257).
- [132] Lex, A., Gehlenborg, N., Strobel, H., Vuilleumot, R. & Pfister, H. 2014 UpSet: Visualization of Intersecting Sets. *IEEE Trans Vis Comput Graph* **20**, 1983-1992. (DOI:10.1109/TVCG.2014.2346248).

Figure 1






Figure 2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46




Figure 3




Figure 4




Figure 5